

1 **REVIEW**

2 **Distribution, status and recent population dynamics of Alpine ibex *Capra ibex* in Europe**

3

4 Alice BRAMBILLA* *Department of Evolutionary Biology and Environmental Studies, University*
5 *of Zurich, Winterthurerstrasse 190, 8057 Zurich (ZH), Switzerland; Alpine Wildlife Research*
6 *Centre, Gran Paradiso National Park, Frazione Jamonin 5, 10080 Noasca (TO), Italy. Email:*
7 [*alicebrambilla1@gmail.com*](mailto:alicebrambilla1@gmail.com)

8 Achaz VON HARDENBERG *Conservation Biology Research Group, Department of Biological*
9 *Sciences, University of Chester, Parkgate Road, CH2 4BJ, Chester, UK. Email:*
10 [*a.vonhardenberg@chester.ac.uk*](mailto:a.vonhardenberg@chester.ac.uk)

11 Luca NELLI *Institute of Biodiversity, Animal Health and Comparative Medicine, University of*
12 *Glasgow, Graham Kerr Building, G12 8QQ Glasgow, UK. Email: [*luca.nelli@glasgow.ac.uk*](mailto:luca.nelli@glasgow.ac.uk)*

13 Bruno BASSANO *Alpine Wildlife Research Centre, Gran Paradiso National Park, Frazione*
14 *Jamonin 5, 10080 Noasca (TO), Italy. Email: [*bruno.bassano@pngp.it*](mailto:bruno.bassano@pngp.it)*

15

16 *Correspondence author

17 **Running head:** Current distribution of Alpine ibex in Europe

18 **Keywords:** Alps, Alpine ibex, *Capra ibex*, current distribution, distribution map, population
19 dynamics, ungulates

20

21 Received: 6 September 2018

22 Accepted: 16 January 2020

23 Editor: DR

24

25 **ABSTRACT**

26 1. Despite its recent successful and well-documented reintroduction history, a comprehensive and

current update of the distribution and status of the Alpine ibex *Capra ibex* is lacking. As some concerns persist about its conservation, a status update appears essential for future conservation and management strategies on a large scale.

2. We provide an exhaustive update of the geographic range of the species, alongside estimates of its current abundance and population trends from 2004 to 2015.

3. We gathered census and distribution data for all the Alpine ibex colonies from management authorities and research groups that monitor them in different countries, and from the literature and publicly available reports. We produced a distribution map, reported the number of individuals observed in the most recent censuses, and estimated global, national, and local population trends using Bayesian hierarchical models.

4. Our model estimated that there were a total of 55297 Alpine ibex in the Alps in 2015 (lower 95% Credible Interval [CrI]: 51157; upper 95% CrI: 62710). The total number of individuals appears to have increased slightly over the last 10 years from the 47000-51000 estimated in previous reports. Positive population trends were observed in Switzerland and Italy, while no trend was apparent in France. For Austria, Germany, and Slovenia, there were insufficient data to estimate a trend. The slopes of the colonies' trends were positively correlated with the year of colony foundation.

5. The geographic range of the Alpine ibex does not seem to have increased in size in recent years, although the accuracy of the spatial data varies among countries.

6. The periodic and standardised collection of census data for all colonies and a common policy of data-sharing at a European level appear essential for monitoring the global trend of this species and for planning balanced conservation and management actions.

INTRODUCTION

The Alpine ibex *Capra ibex* is a charismatic mountain ungulate endemic to the European Alps, where it occurs in all Alpine countries (France, Italy, Switzerland, Lichtenstein, Austria, Germany, Slovenia). Although its elevational range goes from 750 m above sea level (asl) in the Vercors

53 Regional National Park, France, up to more than 3000 m asl in the Western Alps, most Alpine ibex
54 populations are found between 1500 and 3000 m asl, and the most suitable habitat for this species
55 consists of the alpine meadows and rocky cliffs found at this elevation (Grignolio et al. 2003, 2007).
56 The current distribution is, however, a consequence of the recent history of the species which, after
57 suffering almost total extinction in the 19th century, survived only in a restricted area in the north-
58 western Italian Alps and was the object of extensive reintroductions throughout the whole Alpine arc
59 in the 20th century (Stüwe & Nievergelt 1991).

60 The origin of Alpine ibex as a species is still debated (Manceau et al. 1999, Pidancier et al.
61 2006, Kazanskaya et al. 2007), but the ancestors of Alpine ibex are likely to have migrated to Europe
62 from Central Asia around 300000 years ago (Cregut-Bonnoure 1992). At that time, the species
63 occupied its largest range, also outside the Alpine region. At the end of the Riss Glaciation, as a result
64 of reforestation of low-elevation areas, the Alpine ibex became restricted to the Alpine region. In the
65 Middle Ages, the Alpine ibex was still spread throughout the Alps, but intensive hunting, following
66 the development of firearms, brought most populations to extinction (Grodinsky & Stüwe 1987). The
67 first signs of the decline of the species are dated to the late Middle Ages and, in the following centuries
68 (16th – 18th), Alpine ibex gradually disappeared from all Alpine countries except Italy. In Austria and
69 Slovenia, Alpine ibex began disappearing from the beginning of the 18th century. In Switzerland and
70 France, the decline was more progressive, but the last signs of Alpine ibex presence were at the
71 beginning of the 19th century in the Wallis canton. At the beginning of the 19th century, there were
72 no more than a hundred individuals left in a restricted area surrounding the Gran Paradiso Massif,
73 Italy (Grodinsky & Stüwe 1987). Hunting was then prohibited by a Royal decree in 1821 and, thanks
74 to active protection of the species and the institution of the Gran Paradiso Royal Hunting Reserve by
75 King Vittorio Emanuele II in 1856, the population increased to approximately 3000 individuals at the
76 beginning of the 20th century. In the first half of the 20th century, around 90 ibex were captured in the
77 Gran Paradiso area and brought into captive-breeding programs at two Swiss wildlife parks, St.
78 Gallen and Interlaken (Stüwe & Scribner 1989). In the last century, the species was reintroduced first

79 in Switzerland and then in all the other countries of the Alps (Tosi et al. 1986, Wiersema 1990,
80 Giacometti 1991, Stüwe & Nievergelt 1991). Reintroductions were performed from 1911 to 2014,
81 with most of the colonies (82%) founded between 1950 and 2000. 12% of the colonies were founded
82 before 1950, mostly in Switzerland and Italy (Giacometti 1991) plus two in France and one in
83 Germany, and only 6% were founded after 2000. Today, the species is present on the entire Alpine
84 arc, although its range is still fragmented and there are some suitable areas that are not yet occupied
85 (Gruppo Stambecco Europa, Alpine ibex European Specialist Group [GSE-AIESG] unpublished
86 data).

87 The Alpine ibex is included in the Bern Convention (Convention on the Conservation of
88 European Wildlife and Natural Habitats, Appendix III – Protected Fauna Species, 1979) and in the
89 European Directive 43/92/CEE “Habitat”, Annex V (Updated with Directive 97/62/CE, 27 October
90 1997).

91 A map of the geographic range of the species is included in the International Union for
92 Conservation of Nature’s (IUCN) Red List of threatened species (Aulagnier et al. 2008). The IUCN
93 assessment classifies the Alpine ibex as a species of Least Concern “in view of its wide distribution,
94 presumed large population, and because it is not declining at nearly the rate required to qualify for
95 listing in a threatened category”, but also declares that “the species needs conservation action to
96 prevent future decline” (Aulagnier et al. 2008). Indeed, in the last few years, several concerns have
97 arisen about the conservation status of the species. The strong bottleneck that occurred in the 19th
98 century dramatically decreased the genetic variability of the species. The average heterozygosity of
99 Alpine ibex is one of the lowest registered in wild mammals (Biebach & Keller 2010), and the
100 presence of inbreeding depression has been shown in the Gran Paradiso colony, the only remnant (not
101 reintroduced) Alpine ibex population in the Alps (Brambilla et al. 2015). Furthermore, inbreeding
102 reduced the intrinsic per-capita population growth rate in Alpine ibex colonies (Bozzuto et al. 2019).
103 Alpine ibex can interbreed with domestic goats *Capra hircus*, and genetic analyses of the major
104 histocompatibility complex region revealed that successful hybridisation events between the two

105 species, with consequent introgression, have occurred in the past (Grossen et al. 2014). Moreover,
106 despite the general recovery of Alpine ibex in the recent decades on a global scale, the remnant Gran
107 Paradiso colony has shown strong declines (Jacobson et al. 2004, Mignatti et al. 2012). Furthermore,
108 trends in the abundance of the species in different countries reported during the GSE-AIESG meetings
109 in 2012 and 2015 indicate that its status and population dynamics are not consistent throughout the
110 Alps. Finally, direct and indirect consequences of epidemic diseases have affected several recently
111 reintroduced colonies, as well as established ones. Examples are sarcoptic mange in the eastern Alps
112 (Carmignola et al. 2006), brucellosis in the Bargy Massif, France (Mick et al. 2014), respiratory
113 diseases in the Vanoise National Park, France (Garnier et al. 2016), and infectious kerato-
114 conjunctivitis in several areas, including the Gran Paradiso where infectious kerato-conjunctivitis was
115 found to be related to low genetic variation at the major histocompatibility complex region (Brambilla
116 et al. 2018).

117 The rescue and restoration of Alpine ibex represents one of the most successful conservation
118 efforts in Europe. The reintroduction history is well documented, and information on population size
119 is available for all the reintroduced colonies, as most of them were monitored, although with varying
120 effort, at least in the first years after the releases. However, the different management strategies in
121 the European countries (for a brief description of the management regulations in the European
122 countries hosting Alpine ibex populations, see Appendix S1) and the lack of a communal policy on
123 data sharing have resulted in fragmented information about the current status of the species. Some
124 information is published in regional and national reports (also reviewed by De Danieli & Sarasa
125 2015), but much remains unpublished. Despite its recent successful reintroduction history and the
126 strong interest in this charismatic species, a comprehensive and current update of the distribution and
127 status of Alpine ibex is still lacking. Considering the concerns about the conservation of the species,
128 such an update appears to be timely and essential for planning future conservation and management
129 strategies on a continental scale.

130 We provide an exhaustive update of the current geographical range, as well as estimates of

131 abundance and local and global population trends, of the Alpine ibex in the Alps from 2004 to 2015.

132

133 **METHODS**

134 **Definition of distinct populations and census method**

135 For this project, we use a working definition of the term ‘colony’ to describe the distribution of
136 populations of Alpine ibex. When possible, we considered the different reintroduced nuclei, which
137 are historically known and are under different management authorities, as distinct. However, this was
138 possible only for some areas where the colonies are isolated, and there is a clear spatial separation
139 among them. For this reason, we considered differently 1) areas where the oldest and largest colonies
140 are located; 2) areas where the distribution of the animals is continuous and difficult to separate into
141 distinct nuclei; and 3) trans-boundary colonies. For 2) and 3), we defined the colonies following
142 administrative boundaries of the different management authorities that perform censuses.

143 Block counts, also referred to as ground counts, are the most common method used to assess
144 population size in Alpine ibex colonies. The areas occupied by each colony are divided into sectors,
145 and expert observers with good knowledge of the territory count ibex in each sector by means of
146 binoculars or telescopes from footpaths or vantage points. Within an area, the count for all sectors
147 should be conducted in a short time (however, this is not always possible due to observers’ availability
148 or environmental constraints). During the surveys, the total number of observed ibex is recorded. In
149 most of the areas, group size, location and age and sex classes are also recorded. As different
150 authorities do not always perform simultaneous censuses, overlaps in counts are possible due to
151 animal movement across administrative boundaries. However, the risk of double counts is low, as
152 neighbouring authorities tend to coordinate the census. Errors in counts due to missing animals may
153 not be negligible, and for this reason, the numbers presented in this study have to be considered as
154 minimum counts (Largo et al. 2008). For a summary of the colonies considered in the study and their
155 locations, see Appendix S2.

156

157 **Data collection**

158 Data on the abundance and distribution of Alpine ibex were directly requested from management
159 authorities or research groups that monitor colonies in different countries after informing them about
160 the aims of the project. When available, data published in the literature or publicly available online
161 were also gathered. All individuals, public bodies, and research groups that provided data are listed
162 in the Acknowledgements section. A European-wide dataset was built including data from all
163 colonies. When available, the data collected for each colony were: a) the most recent census data; b)
164 number of individuals counted in previous censuses; c) year of colony foundation; d) season of the
165 census; e) information about the area used by ibex during the year (winter and summer range) and
166 any geographical information available; f) sex and age structure composition of the colony. The latter
167 information (d-e-f) was not explicitly used for the analysis of this study, as complete and reliable data
168 were not available for all countries, but was nevertheless stored in the database, which is constantly
169 being updated by the GSE-AIESG with new data and information. The minimum information
170 available for all colonies were: name, country, spatial extent of the occupied area (accuracy differs
171 between colonies), number of individual ibex counted in the last census, and year of the last census.
172 The colony of Gasthofgebirge, Austria, was included in the count data but was not used for models
173 of population dynamics, as its foundation was too recent (2014).

174

175 **Range map**

176 The current geographic range map for Alpine ibex in the European Alps was created using a
177 Geographic Information System (Q GIS 2.18.0 – Las Palmas, QGIS Development Team 2016): the
178 shapefile containing the area occupied by each colony and the associated census data were uploaded
179 to a project containing a digital terrain model of the Alps (reference system WGS 84, UTM 32 N).
180 The area occupied by each colony was defined based on the information received from the
181 management authorities or research groups that monitor colonies (see point e in the previous
182 paragraph). This information was shared as digital shapefiles or as scanned paper maps. In the latter

183 case, the maps were sufficiently accurate to allow us to import them into the QGIS environment,
184 reference them (using the Georeferencer GDAL plugin), and manually digitise them. All the polygons
185 were finally merged into a single shapefile. The density of ibex in each of the different colonies was
186 calculated as the total number of animals counted divided by the area occupied by each colony. In
187 order to increase the accuracy of the area occupied by each colony, we removed unsuitable habitat
188 categories. In order to do this, we intersected the layer of the ibex colonies with the layer of the Corine
189 Land Cover (EEA, 2018) and removed from the colonies layer surfaces belonging to the following
190 categories: “Glaciers and perpetual snow”, “Road and rail networks and associated land”, “Water
191 bodies”, “Urban fabric”. However, as the accuracy of the area considered to be occupied by each
192 colony varies greatly among them, spatial data were not used for further analyses.

193

194 **Count data**

195 In order to obtain the current size of the total Alpine ibex population in Europe and to compare it with
196 estimates obtained from the models, we summed the number of individuals of each colony counted
197 in the last census. We counted the total number of individuals in the Alps and the total for each
198 country. To reduce the risk of double counts or of missing animals due to animal movement across
199 administrative boundaries, when possible, we used censuses performed in the same season (winter or
200 summer counts) for geographically close colonies, as most ibex movements occur seasonally.

201

202 **General model of population dynamics**

203 To model the dynamics of the different colonies, we used a Bayesian hierarchical Poisson model with
204 the year as a fixed effect and the colony (id) and the country as random factors, allowing slopes to
205 vary in each colony. We modelled both random and the fixed effects with a thin-plate spline
206 regression line ($k=10$) allowing for varying variances (Wood 2003). The model was fitted using the
207 MCMCglmm package (Hadfield 2010) in R v.3.3.3 (R core Team 2016). The model was run for
208 130000 iterations with 30000 burn-in and a thinning of 100. We did not include any environmental

209 covariates, as the scope of this analysis was purely to describe the dynamics of the different colonies
210 (and thus of the whole Alpine ibex population), rather than to infer the relative importance of
211 environmental variability in predicting trends.

212 The timespan of the models ranged from 2004 to 2016. However, since several data points
213 were missing for 2016, we chose 2015 as the most recent year with reliable estimations (see Fig. 2),
214 and we present estimates related to 2015 in the Results section. We extracted specific estimates for
215 177 colonies in the Alps (all the nuclei that are present on the map except Gasthofgebirge, Austria)
216 using the *predict* function in MCMCglmm (Hadfield 2010), not marginalising over the random
217 effects to get colony-specific estimates (marginal = NULL). We obtained predictions and lower and
218 upper 95% Credible Intervals (CrI) on the original data scale (counts) using type="response" in the
219 *predict* function. To estimate the total population size per year, we summed up the posterior mcmc
220 samples for each colony and extracted CrIs after back-transforming to the data scale.

221

222 **Countries estimates and effects of the year of foundation on modelled population growth**

223 To evaluate population trends for Alpine ibex in each country, we obtained the slope and confidence
224 intervals (CI) for each country from a linear regression model performed on the yearly estimates
225 extracted from the general Bayesian hierarchical model described in the previous section, with the
226 year as an independent variable. As count data for many of the colonies in Austria, Germany and
227 Slovenia were only available for a few years, the trends were modelled only for Italy, Switzerland
228 and France.

229 The same procedure was used to obtain the population growth rate for each colony (extracting
230 slope and CI from the general model). As more rapid growth is expected in colonies that derive from
231 recent introductions in unoccupied habitats, we tested the effect of the year of colony foundation on
232 population growth rate. This was achieved by modelling individual colony slopes as a function of
233 foundation year with a linear mixed-effects model (lme4 package, Bates et al. 2015), with the country
234 as a random term.

235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260

RESULTS

Range map

A geographic range map showing the current distribution of the Alpine ibex colonies is presented in Fig. 1. From a visual inspection of the map, particularly observing the detail of the borders of the area occupied by each colony, it is possible to assess that the spatial resolution greatly varies among colonies and among countries.

An interactive map with the density of Alpine ibex in each colony as well as a graphical representation of the number of ibex counted in each of them from 2004 to 2015 is presented in Appendix S2.

Count data and models of population dynamics

Based on the information gathered from the management authorities or research groups that monitor Alpine ibex, a total of 178 Alpine ibex colonies were identified over the entire Alpine arc. Alpine ibex colonies are present in all the six major Alpine countries: France, Italy, Switzerland, Germany, Austria, and Slovenia plus Liechtenstein with large differences in the number of individuals and colonies for each country. A total of 55297 Alpine ibex (lower 95% CrI: 51157; upper 95% CrI: 62710) was estimated for 2015 by our Bayesian hierarchical model. Summing up all the individuals from the last censuses conducted in all colonies, we obtained a total count of 53154 individuals which falls within the 95% CrIs of our model. Despite an apparent slight increase of the number of individuals, the slope and CI of the linear regression fitted on the estimated yearly counts do not provide evidence of an increasing trend in the whole Alpine ibex population of the Alps in the years covered by this study (2004-2015, $\beta = 662.8$; lower 95% CI: -232.1; upper 95% CI: 1607.0). Count data and model estimates describing the current status of the species in each country (from west to east) are summarised in Table 1.

The recent trend of the species as a whole and in Italy, France and Switzerland, obtained by

the Bayesian hierarchical Poisson model are presented below. Graphics representing the number of ibex estimated by the model are reported in Fig. 2 and 3(a-f).

FRANCE

In France, there are 30 Alpine ibex colonies, with a total number of 9002 individuals counted (the last census of different colonies ranged from 2008 to 2016). Few counts were available in recent years, and consequently, CrIs for the yearly estimates for the French colonies are relatively wide (Bayesian model median estimate for 2015: 7775 individuals; lower 95% CrI: 5955; upper 95% CrI: 12286). Despite Alpine ibex abundance in France appearing to show a declining trend over the last 10 years (Fig. 3a), the slope of the linear model does not provide evidence of a decline, since the CIs around the estimate broadly overlap zero ($\beta = -118.8$; lower 95% CI: -386.9; upper 95% CI: 404.1).

SWITZERLAND

In Switzerland, there are 45 Alpine ibex colonies, with a total of 17875 individuals counted in 2016. Model estimates indicate 17664 individuals in 2015 (lower 95% CrI: 17398; upper 95% CrI: 17923). Overall, the abundance of Alpine ibex in Switzerland (Fig. 3b) shows an increasing trend over the last decade ($\beta = 306.1$; lower 95% CI: 269.7; upper 95% CI: 347.9).

ITALY

In Italy, there are 67 Alpine ibex colonies, with a total of 16471 individuals counted in the years 2012-2017. The Bayesian model estimates 19872 individuals in 2015 (lower 95% CrI: 16847; upper 95% CrI: 25373). The total number of Alpine ibex in Italy (Fig. 3c) seems to show an increasing trend in the last 10 years ($\beta = 464.6$; lower 95% CI: 107.8; upper 95% CI: 1,194.0). However, although they do not overlap zero, CIs around the estimated slope for the trend are rather wide.

AUSTRIA

287 In Austria, 9013 Alpine ibex assigned to 27 colonies were counted in 2015. The boundaries of the
288 areas occupied by the colonies do not seem to be very accurately known (except for the Hohe Tauern
289 colony), possibly because Alpine ibex counts are done by different authorities that do not always
290 collect spatial information. Moreover, different parts of the area occupied by some colonies might be
291 under the authority of different hunting districts, so it is difficult to obtain a precise number of
292 colonies. Count data are not available for each year in each colony, and therefore CrIs around model
293 estimates are relatively wide (model estimates for 2015: 8813; lower 95% CrI: 8491; upper 95% CrI:
294 9186).

295

296 **GERMANY**

297 In Germany, 516 individual Alpine ibex assigned to five colonies were counted in the years 2014-
298 2017. The model estimates 549 individuals in 2015 (lower 95% CrI: 327; upper 95% CrI: 1003). The
299 figure seems to highlight a declining trend (Fig. 3e) in the last decade, although it was not possible
300 to draw a firm conclusion about the trend of the species in Germany.

301

302 **SLOVENIA**

303 The last data from Slovenia, received in 2016, indicated that 277 Alpine ibex have been counted from
304 four colonies in this country (model estimates for 2015: 323; lower 95% CrI: 185; upper 95% CrI:
305 570). The number of ibex was greatly reduced in the last 10 years due to an epidemic of sarcoptic
306 mange (Iztok Koren, personal communication). However, probably due to the timespan of our
307 models, the declining trend was not evident (Fig. 3f).

308

309 **Table 1.** Number of colonies, and number of individual Alpine ibex *Capra ibex* counted (N counted)
310 and estimated (N estimated) in the European countries in the Alpine Arc. “N counted” was obtained
311 by summing the number of individuals counted during the last census conducted in each population.
312 “Year of last counts” specifies the years in which the last census was conducted. As not all the
313 populations in the same country are surveyed every year, for the same country there may be different

314 “Year of last counts”. “N estimated” represents the number of individuals estimated from a
 315 hierarchical Poisson model for the year 2015. “U95CrI” and “L95CrI” are the upper and lower 95%
 316 Credible Intervals of the Poisson model for 2015. For Austria, the number of colonies was obtained
 317 from the sum of different counting units. “Trend” indicates whether the slopes and CIs provided
 318 evidence of positive trends (+) or no trends (=); NA indicates the countries for which it was not
 319 possible to model the trend because of insufficient data.

320

Country	N colonies	Year of last counts	N counted	N estimated	L95CrI	U95CrI	Trend
France	30	2008-2016	9002	7775	5955	12286	=
Switzerland	45	2016	17875	17664	17398	17923	+
Italy	67	2012-2017	16471	19872	16847	25373	+
Austria	27	2016-2018	9013	8813	8491	9186	NA
Germany	5	2014-2017	516	549	327	1003	NA
Slovenia	4	2016	277	323	185	570	NA
Alpine arc	178	2008-2018	53154	55297	51157	62710	=

321

322

323 **Effects of the year of colony foundation on population growth**

324 The linear mixed effect model for population trends was performed on the slopes of 142 colonies
 325 from three countries (Switzerland N=43; Italy N=65; France N=30). The results of the models
 326 indicated that the year of colony foundation was positively related with the growth rate of the colony
 327 ($\beta \pm SE = 0.087 \pm 0.034$, 95% CI range: 0.020-0.154, $R^2=0.215$). The slope of the colonies founded
 328 in recent times is steeper than that of older colonies.

329

330 **DISCUSSION**

331 The European distribution of Alpine ibex presented in this study confirms that the species is currently

332 present on the entire Alpine arc, with 178 colonies and more than 53000 individuals. A previous
333 attempt to estimate the total number of Alpine ibex produced a total number of 50195 ± 1012
334 individuals in 2013 (De Danieli & Sarasa 2015), which is consistent with the estimate of our model
335 for that year.

336 Comparing the total number of individuals estimated in our study for 2015 ($N= 55297$; lower
337 95% CrI: 51157; upper 95% CrI: 62710) with the minimum number reported for 2005-2007
338 (Apollonio et al. 2009, $N= 47000$ individuals), with the unofficial report provided during the GSE-
339 AIESG meeting in 2012 ($N= 49000$ -50000 individuals), and with the estimate by De Danieli and
340 Sarasa (2015) for 2013 ($N= 49000$ -51000 individuals), the overall number of individuals appears to
341 have increased slightly over the last 10 years. Nevertheless, our linear model does not provide clear
342 evidence of a numerical increase in the total number of individuals during the years 2004-2015, due
343 to the high level of uncertainty around the estimates.

344 Alpine ibex are commonly counted using total block counts, although not all areas were
345 surveyed with the same frequency and number of observers. Moreover, most of the colonies were
346 counted in the pre-reproductive spring season, while others were surveyed after the birth of kids, and
347 a few others during winter. Despite the unavoidable uncertainties in these large-scale population
348 counts, the estimated trends at the global and national scales presented in this work are likely to reflect
349 the actual population dynamics, as the methods used to count the animals have remained constant
350 over time (Jacobson et al. 2004).

351 If we consider the population dynamics of Alpine ibex on a larger temporal scale, considering
352 that 150 years ago only a few hundred individuals survived in a single area (Grodinsky & Stüwe
353 1987), it is clear that the species has recovered and its total abundance has increased. However, the
354 dynamics of the recent decades are harder to interpret. Although the global Alpine ibex population
355 appears to have been stable over the last decade, we observed positive trends in Switzerland and Italy,
356 while it was not possible to draw reliable conclusions about the trends for the other European
357 countries (Fig. 3 and Table 1). This does not mean that the number of Alpine ibex in France, Austria,

358 Germany and Slovenia did not change in recent years, but that the lack of available census data in
359 many years, and the consequent uncertainty around the estimates, does not allow us to discuss the
360 dynamics of the species in these countries.

361 In Switzerland and Austria, Alpine ibex are hunted to regulate the population size of colonies.
362 The population dynamics of Alpine ibex in this country are therefore influenced by management
363 decisions. Before our study (1990-2004), the number of Alpine ibex in Switzerland showed a marked
364 decline (Source: Swiss Federal Office for the Environment BAFU-FOEN), probably due to a
365 combination of hunting pressure (hunting is regulated by the ORES Act of 30 April 1990) and harsh
366 winters. The hunting rate was deliberately reduced from 1999-2000 onwards as a consequence of this
367 decline (BAFU-FOEN and Iris Biebach, personal communication), which may explain the increase
368 observed in the timeframe of our study.

369 In Italy, it is more likely that the observed dynamics are the result of natural processes, as
370 Alpine ibex hunting is forbidden (unfortunately, no data on estimated poaching pressure is available).
371 However, the CIs around the estimate of the growth rate of the species in Italy are rather wide, which
372 suggest caution in interpreting this result as a clear sign of population increase. A possible explanation
373 for the increase of the numbers of Alpine ibex in Switzerland and Italy, despite the different
374 management strategies, may be that several of the colonies in those countries have not yet reached
375 carrying capacity. Indeed, studies performed in the Swiss colonies (Sæther et al. 2007, Bozzuto et al.
376 2019) as well as in the Gran Paradiso colony in Italy (Gran Paradiso National Park, unpublished data)
377 have revealed relatively little density dependence. For the Swiss colony, this may partly be related to
378 active management keeping the colonies below carrying capacity, but it may also be a signal that
379 carrying capacity has not yet been reached. A detailed and rigorous analysis of the population
380 dynamics of each colony was beyond the scope of this work. However, we show that the population
381 growth rate of the colonies is positively related to the year of colony foundation (i.e. that the older
382 colonies are growing less rapidly than the newer ones). This is as expected, since recently founded
383 colonies typically grow exponentially in the first few years. However, the fact that Italy and

384 Switzerland showed positive trends, despite hosting some of the oldest colonies in the Alps, further
385 corroborates the hypothesis that there is still potential for the colonies in those countries to grow. The
386 timeframe of our study (i.e. 12 years, constrained by data availability) may be too limited to point out
387 long-term population trends. The reasons driving the dynamics of the species in Italy and Switzerland
388 would be worth exploring further in more detailed analyses.

389 Translocations of Alpine ibex for restocking purposes have taken place in Swiss and Italian
390 colonies. Translocation was common practice in Switzerland: it is estimated that about 700
391 individuals were moved for restocking from 1950 to 2003, excluding founding events. Restocking is
392 still happening sporadically today, although at a lower rate (40 individuals from 2004 to 2015; source:
393 BAFU-FOEN). In Italy, if we exclude the founding of new colonies, translocations for restocking are
394 not common. Given that most of the restocking events were done within the same country and
395 comprised only a few individuals at a time, it is unlikely that they can have affected global abundance
396 directly. Instead, it is possible that successful restocking events might have increased the growth rate
397 of certain colonies, by reducing inbreeding levels (Hogg et al. 2006, Bozzuto et al. 2019). While it is
398 possible that previous translocations have had a positive effect on population growth, it is not possible
399 to disentangle their effects within the framework of this review, as translocations happened in
400 different years for different colonies, and any genetic effects are likely to be delayed by several years
401 from the restocking events.

402 The high heterogeneity in the spatial resolution of the data at our disposal requires caution
403 when visually interpreting our range map and comparing it with what was reported by previous
404 updates (e.g. Aulagnier et al. 2008). However, spontaneous recolonisation of new areas does not seem
405 to have happened in recent years, although the density of many populations has increased. This is in
406 line with the extremely long recolonisation time required by Alpine ibex (Gauthier & Villaret 1990,
407 Scillitani et al. 2012), and with the fact that spontaneous contact between separate Alpine ibex
408 colonies is unlikely to occur without human intervention, particularly for populations that are far
409 apart. On the other hand, in areas where the Alpine ibex population density is higher, for example in

410 the western Alps, several exchanges between populations have been observed (Groupe National
411 Bouquetin France, personal communication), and in Switzerland at least one colony has become
412 established through natural dispersal (Lukas Keller, personal communication).

413

414 **CONCLUSION**

415 In conclusion, the total abundance of the Alpine ibex in the Alps appears to have remained
416 approximately stable or increased slightly in the last 12 years. However, variation in population trends
417 between countries should be monitored in detail, as it may be a signal of different population
418 dynamics in different areas of the species' range. The isolation of the colonies, combined with low
419 recolonisation rates; extremely low genetic variability (Biebach & Keller 2010) that may lead to
420 inbreeding depression (Brambilla et al. 2015), increased disease susceptibility (Brambilla et al. 2018)
421 and reduced growth rate of colonies (Bozzuto et al. 2019); and local crashes or extinctions due to
422 disease outbreaks are closely linked issues and continue to justify conservation action for the Alpine
423 ibex. Moreover, the effect of climate change on this species has not yet been fully understood as
424 different studies have reported different effects on population parameters (e.g. Pettorelli et al. 2007,
425 Büntgen et al. 2014). However, concerns have been raised about increases in temperature and
426 consequent behavioural changes (Mason et al. 2017), and the likely effects of climate change on
427 population dynamics (Jacobson et al. 2004, Pettorelli et al. 2007, Mignatti et al. 2012). A better
428 knowledge of the immunogenetics mechanisms and of the effect of environmental changes on the
429 dynamics of the species are necessary to monitor its status. The underlying base for all these analyses
430 is the availability of continuous data on the dynamics of each population.

431 Our synthesis highlights the fact that efforts to monitor the size and dynamics of Alpine ibex
432 populations are not homogeneous among different Alpine countries, resulting in high levels of
433 uncertainty around population size estimates for many colonies. This uncertainty hinders the reliable
434 estimation of population trends at the colony and national levels, and therefore the correct assessment
435 of the conservation status of the Alpine ibex in important sectors of its Alpine range. We therefore

436 recommend increasing the long-term monitoring efforts on Alpine ibex in all Alpine countries;
437 organising yearly total block counts in all colonies in the same season where possible; and, ideally,
438 agreeing on a common monitoring protocol for all European Alpine ibex colonies. We believe that
439 such a goal, while ambitious, is achievable, following the example of countries such as Switzerland,
440 where standardised yearly counts are already in place. For the ongoing conservation of the Alpine
441 ibex, it may be advantageous to exploit the transalpine collaboration platforms that already exist, such
442 as the GSE-AIESG and the Large Carnivores, Wild Ungulates and Society Working Group (WISO)
443 of the Alpine Convention for its implementation and coordination.

444

445 **ACKNOWLEDGEMENTS**

446 This work is dedicated to the memory of Dr Vittorio Peracino (1938-2018), former veterinarian of
447 the Gran Paradiso National Park and one of the main actors in the successful re-introduction history
448 of the Alpine ibex. Without his incessant efforts and commitment during his whole career, the
449 geographic range of the species would not be as wide as it is today.

450 In line with the aims of the GSE-AIESG and the WISO platform, the Gran Paradiso National
451 Park (as secretariat of the GSE-AIESG) and the Swiss Ministry for the Environment (BAFU-FOEN),
452 holding the presidency of the WISO platform for the mandate 2013-2014, co-financed this project
453 with the objective of increasing knowledge about the status of the species and the size of the existing
454 colonies. We deeply thank all the people that collaborated on this large project: the scientific service
455 of the Gran Paradiso National Park; Iris Biebach and Lukas Keller of the University of Zurich for
456 helpful discussion on the manuscript; Giuseppe Bogliani from the University of Pavia; the Swiss
457 Federal Office for the Environment including Reinhard Schnidrig and Caroline Nienhuis; Claudio
458 Groff of the WISO platform; and Kelsey Horvath and Matt Geary for revision of the English
459 language. Most of all, we are indebted to all the people who provided data and information on
460 different countries and populations and who are listed below.

461 France: Jerome Cavailhes and Michael Delorme (Parc National de la Vanoise, Groupe

462 National Boquetin), Ludovic Imberdis (Parc National des Ecrins), Patrick Orméa (Parc National du
463 Mercantour), Carole Toïgo (Office National de la Chasse et de la Faune Sauvage), and all the
464 members of Groupe National Bouquetins.

465 Switzerland: Nicolas Bourquin (Centre Suisse de Cartographie de la Faune), Nicole Imesch
466 and Jürg Schenker (Swiss Bundesamt für Umwelt), Iris Biebach (University of Zurich).

467 Italy: Radames Bionda (Aree Protette dell'Ossola), Anna Bonettini (Parco dell'Adamello),
468 Maria Santa Calabrese (Provincia Autonoma di Trento), Eugenio Carlini (Istituto Oikos), Diego Corti
469 (Provincia di Torino), Christian Chioso (Corpo Forestale della Valle d'Aosta), Renato Dotta
470 (Comprensorio Alpino TO4), Marco Favalli (Parco Naturale Dolomiti Friulane), Maria Ferloni
471 (Provincia di Sondrio), Fulvio Genero (Parco Naturale Regionale delle Prealpi Giulie), Omar
472 Giordano (Comprensorio Alpino CN2), Marco Giovo (Comprensorio Alpino TO1), Elena Lux
473 (Provincia di Verbania), Laura Martinelli (Parco Naturale Alpi Marittime), Luca Maurino (Ente di
474 gestione delle Alpi Cozie), Paolo Molinari (Wildlife Consulting), Piergiorgio Partel (Parco Naturale
475 Paneveggio Pale di San Martino), Luca Pedrotti (Parco Nazionale dello Stelvio), Aurelio Perrone
476 (Wildlife Science s.n.c.), Elisa Ramassa (Ente di gestione delle Alpi Cozie), Gianfranco Ribetto (Ente
477 di gestione delle Alpi Cozie), Giovanni Riccardi (Comprensorio Alpino CN1), Davide Righetti
478 (Provincia Autonoma di Bolzano), Domenico Rosselli (Ente di gestione delle Alpi Cozie), Luciano
479 Rossi (Aree Protette Valle Sesia), Stefano Vendrami (Provincia di Belluno), Ramona Viterbi (Parco
480 Nazionale Gran Paradiso).

481 Austria: Gunther Gressmann (Nationalpark Hohe Tauern), Josef Erber (Hunting Association
482 Salzburg), Armin Deutz (Styrian Hunting Association), Ernst Rudigier and Martin Schwärzler
483 (Tyrolean Hunting Association), Hubert Schatz (Amt der Vorarlberger Landesregierung).

484 Germany: Iris Biebach (University of Zurich), Jochen Grab (Nationalpark Berchtesgaden),
485 Christine Miller, Franz Steger (district office Bad Tölz, Bavaria).

486 Slovenia: Iztok Koren (Slovenia Forest Service).

487
488

489 **REFERENCES**

- 490 Apollonio M, Giacometti M, Lanfranchi P, Lovari S, Meneguz PG, Molinari P et al. (2009) Piano di
491 conservazione, diffusione e gestione dello stambecco sull’arco Alpino italiano. Provincia di Sondrio.
492
- 493 Aulagnier S, Kranz A, Lovari S, Jdeidi T, Masseti M, Nader I, de Smet K, Cuzin F (2008) *Capra*
494 *ibex*. IUCN Red List of Threatened Species. Version 2013.1.
495
- 496 Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4.
497 *Journal of Statistical Software* 67: 1-48.
498
- 499 Biebach I, Keller LF (2010) Inbreeding in reintroduced populations: the effects of early reintroduction
500 history and contemporary processes. *Conservation Genetics* 11: 527-538.
501
- 502 Bozzuto C, Biebach I, Muff S, Ives AR, Keller LF (2019) Inbreeding reduces long-term growth of
503 Alpine ibex populations. *Nature Ecology & Evolution* 3: 1359-1364.
504
- 505 Brambilla A, Biebach I, Bassano B, Bogliani G, von Hardenberg A (2015) Direct and indirect causal
506 effects of heterozygosity on fitness-related traits in Alpine ibex. *Proceedings of the Royal Society of*
507 *London B: Biological Sciences* 282: 20141873.
508
- 509 Brambilla A, Keller L, Bassano B, Grossen C (2018) Heterozygosity–fitness correlation at the
510 major histocompatibility complex despite low variation in Alpine ibex (*Capra ibex*). *Evolutionary*
511 *Applications* 11: 631-644.
512
- 513 Büntgen U, Liebhold A, Jenny H, Myserud A, Egli S, Nievergelt D, Stenseth NC, Bollmann K
514 (2014) European springtime temperature synchronises ibex horn growth across the eastern Swiss
515 Alps. *Ecology Letters* 17: 303-313.
516
- 517 Carmignola G, Stefani P, Gerstgrasser L (2006) 6° Rapporto rogna sarcoptica. Provincia autonoma
518 di Bolzano, Ufficio caccia e pesca.
519
- 520 EEA (2018) Corine Land Cover (CLC) 2018, Version 20b2. Release date: 21-12-2018. European
521 Environment Agency. <https://land.copernicus.eu/pan-european/corine-land-cover/clc2018>
522
- 523 Cregut-Bonnoure E (1992) Dynamics of bovid migration in western Europe during the middle and

524 late Pleistocene. *Courier Forschungsinstitut Senckenberg* 153: 177-185.

525

526 De Danieli C, Sarasa M (2015) Population estimates, density-dependence and the risk of disease
 527 outbreaks in the Alpine ibex *Capra ibex*. *Animal Biodiversity and Conservation* 38: 101-119.

528

529 Garnier A, Gaillard JM, Gauthier D, Besnard A (2016) What shapes fitness costs of reproduction in
 530 long - lived iteroparous species? A case study on the Alpine ibex. *Ecology* 97: 205-214.

531

532 Gauthier D, Villaret JC (1990) La réintroduction en France du bouquetin des Alpes. *Revue d'Écologie*
 533 *(La Terre et La Vie)* 5: 97-120.

534

535 Giacometti M (1991) Beitrag zur Ansiedlungsdynamik und aktuellen Verbreitung des
 536 Alpensteinbockes (*Capra i. ibex* L.) im Alpenraum. *Zeitschrift für Jagdwissenschaft* 37: 157-173.

537

538 Grignolio S, Parrini F, Bassano B, Luccarini S, Apollonio M (2003) Habitat selection of adult males
 539 of Alpine ibex, *Capra ibex ibex*. *Folia Zoologica* 52: 113-120.

540

541 Grignolio S, Rossi I, Bertolotto E, Bassano B, Apollonio M (2007) Influence of the kid on space use
 542 and habitat selection of female Alpine ibex. *Journal of Wildlife Management* 71: 713-719.

543

544 Grodinsky C, Stüwe M (1987) With lots of help alpine ibex return to their mountains. *Smithsonian*
 545 18: 68-77.

546

547 Grossen C, Keller LF, Biebach I, Croll D (2014). Introgression from domestic goat generated
 548 variation at the major histocompatibility complex of Alpine ibex. *PLOS Genetics* 10:e1004438.

549

550 Grossen C, Biebach I, Angelone-Alasaad S, Keller LF, Croll D (2018) Population genomics
 551 analyses of European ibex species show lower diversity and higher inbreeding in reintroduced
 552 populations. *Evolutionary Applications* 11: 123-139.

553

554 Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed models: the
 555 MCMCglmm R package. *Journal of Statistical Software* 33: 1-22. <http://www.jstatsoft.org/v33/i02/>

556

557 Hogg JT, Forbes SH, Steele BM, Luikart G (2006) Genetic rescue of an insular population of large
 558 mammals. *Proceedings of the Royal Society B: Biological Sciences* 273: 1491-1499.

559

560 Jacobson AR, Provenzale A, von Hardenberg A, Bassano B, Festa-Bianchet M (2004) Climate
561 forcing and density dependence in a mountain ungulate population. *Ecology* 85: 1598-1610.

562

563 Kazanskaya EY, Kuznetsova MV, Danilkin AA (2007) Phylogenetic reconstructions in the genus
564 *Capra* (Bovidae, Artiodactyla) based on the mitochondrial DNA analysis. *Russian Journal of*
565 *Genetics* 43: 181-189.

566

567 Largo E, Gaillard JM, Festa-Bianchet M, Toïgo C, Bassano B, Cortot H et al. (2008) Can ground
568 counts reliably monitor ibex *Capra ibex* populations. *Wildlife Biology* 14: 489-500.

569

570 Manceau V, Després L, Bouvet J, Taberlet P (1999) Systematics of the genus *Capra* inferred from
571 mitochondrial DNA sequence data. *Molecular Phylogenetics and Evolution* 13: 504-510.

572

573 Mick V, Le Carrou G, Corde Y, Game Y, Jay M, Garin-Bastuji B (2014) *Brucella melitensis* in
574 France: persistence in wildlife and probable spillover from Alpine ibex to domestic animals. *PLoS*
575 *One* 9(4), e94168.

576

577 Mason TH, Brivio F, Stephens PA, Apollonio M, Grignolio S (2017) The behavioral trade-off
578 between thermoregulation and foraging in a heat-sensitive species. *Behavioral ecology* 28: 908-918.

579

580 Mignatti A, Casagrandi R, Provenzale A, von Hardenberg A, Gatto M (2012) Sex- and age-structured
581 models for Alpine ibex *Capra ibex ibex* population dynamics. *Wildlife Biology* 18: 318-332.

582

583 Pettorelli N, Pelletier F, von Hardenberg A, Festa-Bianchet M, Côté SD (2007) Early onset of
584 vegetation growth vs. rapid green-up: impacts on juvenile mountain ungulates. *Ecology* 88: 381-
585 390.

586

587 Pidancier N, Jordan S, Luikart G, Taberlet P (2006) Evolutionary history of the genus *Capra*
588 (Mammalia, Artiodactyla): discordance between mitochondrial DNA and Y-chromosome
589 phylogenies. *Molecular Phylogenetics and Evolution* 40: 739-749.

590

591 QGIS Development Team (2016) QGIS Geographic Information System. Open Source Geospatial
592 Foundation Project. <http://qgis.osgeo.org>

593

594 R Core Team (2016) *R: a Language and Environment for Statistical Computing*. R Foundation for
 595 Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
 596

597 Scillitani L, Sturaro E, Menzano A, Rossi L, Viale C, Ramanzin M (2012). Post-release spatial and
 598 social behaviour of translocated male Alpine ibexes (*Capra ibex ibex*) in the eastern Italian Alps.
 599 *European Journal of Wildlife Research* 58: 461-472.
 600

601 Sæther BE, Lillegård M, Grøtan V, Filli F, Engen S (2007). Predicting fluctuations of reintroduced
 602 ibex populations: the importance of density dependence, environmental stochasticity and uncertain
 603 population estimates. *Journal of Animal Ecology* 76: 326-336.
 604


605 Stüwe M, Nievergelt B (1991) Recovery of alpine ibex from near extinction: the result of effective
 606 protection, captive breeding, and reintroductions. *Applied Animal Behaviour Science* 29: 379-387.
 607

608 Stüwe M, Scribner KT (1989) Low genetic variability in reintroduced alpine ibex (*Capra ibex ibex*)
 609 populations. *Journal of Mammalogy* 70: 370-373.
 610


611 Tosi G, Scherini G, Apollonio M, Ferrario G, Toso S, Pacchetti G, Guidali F (1986) Modello di
 612 valutazione ambientale per la reintroduzione dello stambecco (*Capra ibex ibex*). *Supplemento alle*
 613 *ricerche di biologia della selvaggina* 77: 5-77.
 614

615 Wiersema G, Gauthier D (1990) Status and aspects of reintroduction and management of the Ibex in
 616 the Alps. *Travaux Scientifiques du Parc National de la Vanoise* 18: 235-252.
 617

618 Wood SN (2003) Thin plate regression splines. *Journal of the Royal Statistical Society: Series B*
 619 *Statistical Methodology* 65: 95-114.




621
622 **Fig. 1.** Geographic range map of the 178 Alpine ibex *Capra ibex* populations in Europe. For a
623 summary of all colonies where numbers of Alpine ibex have been counted, see Appendix S2.
624



625
626 **Fig. 2.** Estimates of the trend in numbers of Alpine ibex present on the Alps from 2004 to 2016. The
627 black line represents the estimated number and the filled area indicates the Credible Intervals
628 extracted by Bayesian hierarchical Poisson models. The y-axis lower limit was set to 40000 for a
629 better view of the trend.

630
631



632

633 **Fig. 3.** Estimates of the trend in numbers of Alpine ibex present in Alpine countries from 2004 to
634 2016. The black line represents the estimated number and the filled area the Credible Intervals
635 extracted by Bayesian hierarchical Poisson models. The y-axis limits differ for each plot to favour
636 readability of the trends.
637

638
639
640
641
642
643
644
645
646
647
648
649
650

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher’s website.

Appendix S1 Description of the different management regulations in the European countries hosting Alpine ibex populations.

Appendix S2 Interactive map showing all the populations considered in the study, the species’ geographic range, and the location, density, and number of Alpine ibex counted from 2004 to 2016.