

NEW BINARY SELF-DUAL CODES VIA A GENERALIZATION OF THE FOUR CIRCULANT CONSTRUCTION

JOE GILDEA, ABİDİN KAYA, AND BAHATTIN YILDIZ*

ABSTRACT. In this work, we generalize the four circulant construction for self-dual codes. By applying the constructions over the alphabets \mathbb{F}_2 , $\mathbb{F}_2 + u\mathbb{F}_2$, $\mathbb{F}_4 + u\mathbb{F}_4$, we were able to obtain extremal binary self-dual codes of lengths 40, 64 including new extremal binary self-dual codes of length 68. More precisely, 43 new extremal binary self-dual codes of length 68, with rare new parameters have been constructed.

1. INTRODUCTION

Binary self-dual codes have generated a considerable amount of interest in the literature for decades for their connections to many other mathematical structures and applications. They have an upper bound on their minimum distance, which is given by Conway and Sloane in [4], and is finalized by Rains in [16] as $d \leq 4\lfloor \frac{n}{24} \rfloor + 6$ when $n \equiv 22 \pmod{24}$ and $d \leq 4\lfloor \frac{n}{24} \rfloor + 4$, otherwise, where n is the length of the self-dual code. Self-dual codes meeting these bounds are called *extremal*.

There is an extensive literature on constructions for extremal binary self-dual codes. One of the main directions of research in the literature has been to construct extremal binary self-dual codes whose weight enumerators have new parameters, that were not known to exist before. This comes from the works by Conway and Sloane in [4] and Dougherty et al. in [5] in which the possible weight enumerators of all extremal self-dual codes of lengths up to 100 were classified.

While the tools in constructing extremal binary self-dual codes may differ from taking a special matrix construction, considering a certain automorphism or the neighboring construction, in all of these cases the final step is to do a computer search over a reduced set of possible inputs. Using the afore-mentioned tools reduce the search field considerably so that the search is now feasible to do in a reasonable time.

For most known constructions for self-dual codes, one of the key concepts is “circulant” matrices. It is well-known that circulant matrices are determined uniquely by their first rows and that they commute in matrix multiplication. The double-circulant, bordered double circulant and four-circulant constructions are some of the well-known construction methods in the literature that make use of circulant matrices. Through these constructions the search field for a self-dual code of length $2n$ usually reduces to a constant multiple of 2^n , which makes it feasible to search for self-dual codes of lengths up to 88 for example.

2010 *Mathematics Subject Classification*. Primary 94B05, 15B33.

Key words and phrases. circulant matrices; extremal self-dual codes; Gray maps; extension theorems; neighboring construction.

*The Corresponding Author.

In this work, we will be considering a generalized version of the four-circulant construction over the alphabets \mathbb{F}_2 , $\mathbb{F}_2 + u\mathbb{F}_2$ and $\mathbb{F}_4 + u\mathbb{F}_4$ to construct extremal binary self-dual codes. Our construction, in general, is different than the four-circulant construction and we will be giving the comparative results. Using this construction, we are able to construct many extremal binary self-dual codes of lengths 40 and 64, and in particular we are able to construct 43 new extremal binary self-dual codes of length 68 with new weight enumerators in $W_{68,2}$. The exact parameters in the weight enumerators are given in section 5.

The rest of the paper is organized as follows. In section 2, we give the preliminaries on the alphabets to be used, special types of matrices that we use in our constructions and the well known four circulant construction. In section 3, we introduce our generalization of the four circulant construction and we give theoretical results about when they lead to self-dual codes as well as their connection to the ordinary four-circulant construction. In section 4 we give the numerical results about extremal binary self-dual codes of lengths 40 and 64 that we obtain by a direct application of our constructions over different alphabets together with a comparison with the usual four-circulant construction. In section 5, we apply the neighboring construction as well as extensions to the codes obtained in section 4 to find new extremal binary self-dual codes of length 68. We finish with concluding remarks and directions for possible future research.

2. PRELIMINARIES

Let \mathcal{R} be a commutative Frobenius ring of characteristic 2. A code \mathcal{C} of length n over \mathcal{R} is an \mathcal{R} -submodule of \mathcal{R}^n . Elements of the code \mathcal{C} are called codewords of \mathcal{C} . Let $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$ be two elements of \mathcal{R}^n . The duality is understood in terms of the Euclidean inner product; $\langle x, y \rangle_E = \sum x_i y_i$. The dual \mathcal{C}^\perp of the code \mathcal{C} is defined as

$$\mathcal{C}^\perp = \{x \in \mathcal{R}^n \mid \langle x, y \rangle_E = 0 \text{ for all } y \in \mathcal{C}\}.$$

We say that \mathcal{C} is self-dual if $\mathcal{C} = \mathcal{C}^\perp$.

Two self-dual binary codes of dimension k are said to be neighbors if their intersection has dimension $k - 1$.

Let $\mathbb{F}_4 = \mathbb{F}_2(\omega)$ be the quadratic field extension of the binary field $\mathbb{F}_2 = \{0, 1\}$, where $\omega^2 + \omega + 1 = 0$. The ring $\mathbb{F}_4 + u\mathbb{F}_4$ defined via $u^2 = 0$ is a commutative binary ring of size 16. We may easily observe that it is isomorphic to $\mathbb{F}_2[\omega, u] / \langle u^2, \omega^2 + \omega + 1 \rangle$. The ring has a unique non-trivial ideal $\langle u \rangle = \{0, u, u\omega, u + u\omega\}$. Note that $\mathbb{F}_4 + u\mathbb{F}_4$ can be viewed as an extension of $\mathbb{F}_2 + u\mathbb{F}_2$ and so we can describe any element of $\mathbb{F}_4 + u\mathbb{F}_4$ in the form $\omega a + \bar{\omega}b$ uniquely, where $a, b \in \mathbb{F}_2 + u\mathbb{F}_2$.

$$\begin{array}{ccc} (\mathbb{F}_4 + u\mathbb{F}_4)^n & \xrightarrow{\psi_{\mathbb{F}_4 + u\mathbb{F}_4}} & (\mathbb{F}_2 + u\mathbb{F}_2)^{2n} \\ \downarrow \varphi_{\mathbb{F}_4 + u\mathbb{F}_4} & & \downarrow \varphi_{\mathbb{F}_2 + u\mathbb{F}_2} \\ \mathbb{F}_4^{2n} & \xrightarrow{\psi_{\mathbb{F}_4}} & \mathbb{F}_2^{4n} \end{array}$$

Let us recall the following Gray Maps from [8, 15] and [6];

$$\begin{aligned}
\psi_{\mathbb{F}_4} &: a\omega + b\bar{\omega} \mapsto (a, b), \quad a, b \in \mathbb{F}_2^n \\
\varphi_{\mathbb{F}_2+u\mathbb{F}_2} &: a + bu \mapsto (b, a + b), \quad a, b \in \mathbb{F}_2^n \\
\psi_{\mathbb{F}_4+u\mathbb{F}_4} &: a\omega + b\bar{\omega} \mapsto (a, b), \quad a, b \in (\mathbb{F}_2 + u\mathbb{F}_2)^n \\
\varphi_{\mathbb{F}_4+u\mathbb{F}_4} &: a + bu \mapsto (b, a + b), \quad a, b \in \mathbb{F}_4^n
\end{aligned}$$

Note that these Gray maps preserve orthogonality in the respective alphabets, for the details we refer to [15]. The binary codes $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ and $\psi_{\mathbb{F}_4} \circ \varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ are equivalent to each other. The Lee weight of an element in $\mathbb{F}_4 + u\mathbb{F}_4$ is defined to be the Hamming weight of its binary image under any of the previously mentioned compositions of the maps. A self-dual code is said to be of Type II if the Lee weights of all codewords are multiples of 4, otherwise it is said to be of Type I.

Proposition 2.1. ([15]) *Let C be a code over $\mathbb{F}_4 + u\mathbb{F}_4$. If C is self-orthogonal, so are $\psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ and $\varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$. C is a Type I (resp. Type II) code over $\mathbb{F}_4 + u\mathbb{F}_4$ if and only if $\varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ is a Type I (resp. Type II) \mathbb{F}_4 -code, if and only if $\psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ is a Type I (resp. Type II) $\mathbb{F}_2 + u\mathbb{F}_2$ -code. Furthermore, the minimum Lee weight of C is the same as the minimum Lee weight of $\psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ and $\varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$.*

Corollary 2.2. *Suppose that C is a self-dual code over $\mathbb{F}_4 + u\mathbb{F}_4$ of length n and minimum Lee distance d . Then $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ is a binary $[4n, 2n, d]$ self-dual code. Moreover, C and $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ have the same weight enumerator. If C is Type I (Type II), then so is $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$.*

In subsequent sections we will be writing tables in which vectors with elements from the rings $\mathbb{F}_2 + u\mathbb{F}_2$ and $\mathbb{F}_4 + u\mathbb{F}_4$ will appear. In order to avoid writing long vectors with elements that can be confused with other elements, we will be describing the elements of this ring in a shorthand way, which will make the tables more compact.

For the elements of $\mathbb{F}_2 + u\mathbb{F}_2$ we will use $0 \rightarrow 0, 1 \rightarrow 1, u \rightarrow u$ and $1 + u \rightarrow 3$.

For the elements of $\mathbb{F}_4 + u\mathbb{F}_4$, we use the ordered basis $\{\omega, \bar{\omega}, u, 1\}$ to express the elements of $\mathbb{F}_4 + u\mathbb{F}_4$ as binary strings of length 4. Then we will use the hexadecimal number system to describe each element:

$0 \leftrightarrow 0000, 1 \leftrightarrow 0001, 2 \leftrightarrow 0010, 3 \leftrightarrow 0011, 4 \leftrightarrow 0100, 5 \leftrightarrow 0101, 6 \leftrightarrow 0110, 7 \leftrightarrow 0111, 8 \leftrightarrow 1000, 9 \leftrightarrow 1001, A \leftrightarrow 1010, B \leftrightarrow 1011, C \leftrightarrow 1100, D \leftrightarrow 1101, E \leftrightarrow 1110, F \leftrightarrow 1111.$

For example $1 + u\omega$ corresponds to 1001, which is represented by the hexadecimal 9, while $\omega + u\omega$ corresponds to 1100, which is represented by C.

We are going to use the following extension method for computational results in the upcoming sections.

Theorem 2.3. ([7]) *Let R be a commutative ring of characteristic 2 with identity. Let C be a self-dual code over R of length n and $G = (r_i)$ be a $k \times n$ generator matrix for C , where r_i is the i -th row of G , $1 \leq i \leq k$. Let c be a unit in R such that $c^2 = 1$ and X be a vector in R^n with $\langle X, X \rangle = 1$. Let $y_i = \langle r_i, X \rangle$ for $1 \leq i \leq k$.*

Then the following matrix

$$\left[\begin{array}{cc|c} 1 & 0 & X \\ \hline y_1 & cy_1 & r_1 \\ \vdots & \vdots & \vdots \\ y_k & cy_k & r_k \end{array} \right],$$

generates a self-dual code D over R of length $n + 2$.

2.1. Special Matrices. Circulant matrices play an important role in many applications. In this section we briefly recall circulant matrices and its variations in the form of reverse-circulant and λ -circulant matrices. For more detailed information on circulant matrices we refer the reader to [18], [17] and the references therein.

With R a commutative ring with identity, let σ be the permutation on R^n that corresponds to the right shift, i.e.

$$(2.1) \quad \sigma(a_1, a_2, \dots, a_n) = (a_n, a_1, \dots, a_{n-1}).$$

A circulant matrix is a square matrix where each row is a right-circular shift of the previous row. In other words, if \bar{r} is the first row, a typical circulant matrix is of the form

$$(2.2) \quad \left[\begin{array}{c} \bar{r} \\ \hline \sigma(\bar{r}) \\ \hline \sigma^2(\bar{r}) \\ \hline \vdots \\ \hline \sigma^{n-1}(\bar{r}) \end{array} \right].$$

It is clear that, with T denoting the permutation matrix corresponding to the n -cycle $(123\dots n)$, a circulant matrix with first row (a_1, a_2, \dots, a_n) can be expressed as a polynomial in T as:

$$a_1 I_n + a_2 T + a_3 T^2 + \dots + a_n T^{n-1},$$

with $T^n = I_n$. This shows that circulant matrices commute.

A reverse-circulant matrix is a square matrix where each row is a left-circular shift of the previous row. It is clear to see that if \bar{r} is the first row, a reverse-circulant matrix is of the form

$$(2.3) \quad \left[\begin{array}{c} \bar{r} \\ \hline \sigma^{-1}(\bar{r}) \\ \hline \sigma^{-2}(\bar{r}) \\ \hline \vdots \\ \hline \sigma^{-(n-1)}(\bar{r}) \end{array} \right].$$

An $n \times n$ square matrix A is called λ -circulant if every row is a λ -cyclic shift of the previous one, in other words A is in the following form;

$$\begin{pmatrix} a_1 & a_2 & a_3 & \cdots & a_n \\ \lambda a_n & a_1 & a_2 & \cdots & a_{n-1} \\ \lambda a_{n-1} & \lambda a_n & a_1 & \cdots & a_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \lambda a_2 & \lambda a_3 & \lambda a_4 & \cdots & a_1 \end{pmatrix}.$$

λ -circulant matrices are an immediate generalization of circulant matrices and like circulant matrices, two λ -circulant matrices also commute.

λ -reverse-circulant matrices can also be defined in exactly the same way as an extension of reverse circulant matrices.

The following lemma gives us an important result that we will be using in the upcoming sections.

Lemma 2.4. [13] *Let A and C be λ -circulant matrices. Then $C' = CR$ is a λ -reverse-circulant matrix and it is symmetric. Here R is the back-diagonal matrix. Moreover, $AC' - C'A^T = 0$. Equivalently, $ARC^T - CRA^T = 0$.*

A special case of Lemma 2.4 is as follows;

Lemma 2.5. *Symmetric circulant matrices commute with reverse circulant matrices.*

2.2. On four circulant construction. The four-circulant construction, which was inspired by orthogonal designs, was introduced in [2]:

Theorem 2.6. [2] *Let A and B be $n \times n$ circulant matrices over \mathbb{F}_p such that $AA^T + BB^T = -I_n$ then the matrix*

$$G = \left(I_{2n} \left| \begin{array}{cc} A & B \\ -B^T & A^T \end{array} \right. \right)$$

generates a self-dual code over \mathbb{F}_p .

Recently, the four circulant construction was applied on $\mathbb{F}_2 + u\mathbb{F}_2$ in [11], which resulted in a new binary self-dual code of length 64.

The following is a variation of the four circulant construction, which was used in [13] to obtain new extremal binary self-dual codes.

Theorem 2.7. [13] *Let λ be a unit of the commutative Frobenius ring \mathcal{R} , A be a λ -circulant matrix and B be a λ -reverse-circulant matrix with $AA^T + BB^T = -I_n$ then the matrix*

$$G = \left(I_{2n} \left| \begin{array}{cc} A & B \\ -B & A \end{array} \right. \right)$$

generates a self-dual code \mathcal{C} over \mathcal{R} .

3. A GENERALIZATION OF THE FOUR CIRCULANT CONSTRUCTION

In this section, we give a generalization of the four circulant construction. We also propose two specific variations of the construction.

Theorem 3.1. *Let \mathcal{R} be a commutative Frobenius ring of characteristics 2, A and B be circulant matrices and C be a reverse circulant matrix. Then the code generated by*

$$G := \left(I_{2n} \left| \begin{array}{cc} A & B + C \\ B^T + C & A^T \end{array} \right. \right)$$

is self-dual when $AA^T + BB^T + C^2 = I_n$ and $AC = CA$.

Proof. Let $M := \begin{pmatrix} A & B+C \\ B^T+C & A^T \end{pmatrix}$. We are to show that $MM^T = I_{2n}$ under the given conditions. Indeed

$$\begin{aligned} MM^T &= \begin{pmatrix} AA^T + BB^T + BC + CB^T + C^2 & AB + AC + BA + CA \\ B^T A^T + CA^T + A^T B^T + A^T C & B^T B + B^T C + CB + C^2 + A^T A \end{pmatrix} \\ &= \begin{pmatrix} AA^T + BB^T + C^2 & AC + CA \\ CA^T + A^T C & B^T B + C^2 + A^T A \end{pmatrix} = \begin{pmatrix} I_n & 0_n \\ 0_n & I_n \end{pmatrix}. \end{aligned}$$

The above equality holds because we have $AB = BA$ and $A^T B^T = B^T A^T$ since circulant matrices commute and also because by Lemma 2.4 $BC + CB^T = 0$ and $B^T C + CB = 0$. \square

We obtain the following corollary when A is a symmetric circulant matrix:

Corollary 3.2. *Let \mathcal{R} be a commutative Frobenius ring of characteristic 2, A be a symmetric circulant matrix, B be a circulant matrix and C be a reverse circulant matrix. Then the code generated by*

$$G := \left(I_{2n} \mid \begin{array}{cc} A & B+C \\ B^T+C & A \end{array} \right)$$

is a self-dual code over \mathcal{R} whenever $A^2 + BB^T + C^2 = I_n$.

Proof. It follows by Theorem 3.1 and Lemma 2.5. \square

We may also propose another special case of Theorem 3.1.

Corollary 3.3. *Let \mathcal{C} be a self-dual four circulant code of length $4n$ over \mathcal{R} (of characteristic 2) generated by*

$$G := \left(I_{2n} \mid \begin{array}{cc} A & B \\ B^T & A^T \end{array} \right).$$

Then for any reverse circulant matrix C , which commutes with A and satisfies $C^2 = 0$, the matrix

$$\left(I_{2n} \mid \begin{array}{cc} A & B+C \\ B^T+C & A^T \end{array} \right)$$

generates a self-dual code \mathcal{C}' .

Corollary 3.3 allows us to reduce the size of the search field for that specific variation. We may consider a four circulant code and search for reverse circulant matrices C under the restrictions.

Example 3.4. Let $n = 7$ and \mathcal{C} be the four circulant code where $A = I_7$ and $B = 0_7$, i.e. $r_A = (1, 0, 0, 0, 0, 0, 0)$ and $r_B = (0, 0, 0, 0, 0, 0, 0)$. The code \mathcal{C} is binary self-dual with parameters $[28, 14, 2]$. Let C be the reverse circulant matrix with first row $r_C = (1110100)$ which satisfies $C^2 = 0_7$ and obviously it commutes with A . Then the code \mathcal{C}' obtained by Corollary 3.3 is an extremal binary self-dual $[28, 14, 6]$ code with an automorphism group of order $2^6 \times 3 \times 7$.

4. COMPUTATIONAL RESULTS

In this section, we provide examples to demonstrate the effectiveness of the methods introduced in Section 3. We also compare the methods with the well known four circulant construction for various lengths over the alphabets \mathbb{F}_2 , $\mathbb{F}_2 + u\mathbb{F}_2$ and $\mathbb{F}_4 + u\mathbb{F}_4$.

4.1. Comparison of the methods over \mathbb{F}_2 for length 40. We construct Type I self-dual codes of length 40 by the four circulant construction and also by the methods given in Section 3. The weight enumerator of a singly even binary self-dual code of parameters $[40, 20, 8]$ is in the following form:

$$W_{40} = 1 + (125 + 16\beta)y^8 + (1664 - 64\beta)y^{10} + \dots, 0 \leq \beta \leq 10$$

The existence of a code with $\beta = 9$ is still an open problem. There are codes for the other values. In Table 1, we list four circulant self-dual binary codes of length 40.

TABLE 1. $[40, 20, 8]$ four circulant codes

$\mathcal{C}_{40,i}$	r_A	r_B	$ Aut(\mathcal{C}_{40,i}) $	β in W_{40}
$\mathcal{C}_{40,1}$	(0100001110)	(0100110011)	$2^2 \times 5$	0
$\mathcal{C}_{40,2}$	(0000110011)	(0010111001)	$2^3 \times 5$	0
$\mathcal{C}_{40,3}$	(0101100111)	(1111001011)	$2^3 \times 3 \times 5$	0
$\mathcal{C}_{40,4}$	(1001000100)	(0101101101)	$2^{14} \times 3 \times 5$	10
$\mathcal{C}_{40,5}$	(1000000010)	(1101011101)	$2^{16} \times 3^3 \times 5^2$	10

The binary self-dual codes of length 40 constructed from the construction given in Corollary 3.2 are given in Table 2. Since A is symmetric circulant, we only list the necessary entries of the first row r_A .

TABLE 2. $[40, 20, 8]$ codes by Corollary 3.2

$\mathcal{D}_{40,i}$	r_A	r_B	r_C	$ Aut(\mathcal{D}_{40,i}) $	β in W_{40}
$\mathcal{D}_{40,1}$	(110111)	(0010001100)	(1110100101)	2^3	0
$\mathcal{D}_{40,2}$	(011010)	(0010111000)	(1110010010)	2^4	0
$\mathcal{D}_{40,3}$	(011010)	(0010011100)	(1010000100)	$2^3 \times 3$	0
$\mathcal{D}_{40,4}$	(100111)	(1010110011)	(0000100110)	2^6	0
$\mathcal{D}_{40,5}$	(111001)	(1001011010)	(0010011001)	2^7	0
$\mathcal{D}_{40,6}$	(010111)	(1001111001)	(0001011111)	2^2	1
$\mathcal{D}_{40,7}$	(011001)	(0111000011)	(0111100001)	2^3	1
$\mathcal{D}_{40,8}$	(011010)	(0010000110)	(1100000000)	2^4	1
$\mathcal{D}_{40,9}$	(110001)	(1011000011)	(0100110100)	2^3	2
$\mathcal{D}_{40,10}$	(000110)	(0100010011)	(1100001000)	2^6	2
$\mathcal{D}_{40,11}$	(111001)	(1100101001)	(0100111110)	$2^9 \times 3^2$	2
$\mathcal{D}_{40,12}$	(110100)	(1101111011)	(0011111001)	2^{13}	2
$\mathcal{D}_{40,13}$	(001010)	(1010000111)	(1111110011)	2^5	4
$\mathcal{D}_{40,14}$	(011000)	(0101010110)	(0000001100)	2^6	4
$\mathcal{D}_{40,15}$	(100111)	(0000110011)	(1010011001)	2^7	4
$\mathcal{D}_{40,16}$	(000010)	(0011111000)	(0100101010)	2^8	6
$\mathcal{D}_{40,17}$	(000010)	(1011101000)	(0100101010)	$2^8 \times 3$	6
$\mathcal{D}_{40,18}$	(010100)	(1000111111)	(0100100101)	2^{15}	10
$\mathcal{D}_{40,19}$	(110100)	(0101010101)	(1001110010)	$2^{14} \times 3 \times 5$	10
$\mathcal{D}_{40,20}$	(101010)	(1000100010)	(0101111101)	$2^{16} \times 3^3 \times 5^2$	10

We apply Corollary 3.3 to $\mathcal{C}_{40,4}$ from Table 1. In other words, we fix the circulant matrices A and B and search for reverse circulant matrices which satisfy the given conditions. The results are tabulated in Table 3. The results have shown the method to be quite effective.

TABLE 3. $[40, 20, 8]$ codes by Corollary 3.3 for $\mathcal{C}_{40,4}$

$\mathcal{E}_{40,i}$	r_C	$ Aut(\mathcal{E}_{40,i}) $	β in W_{40}
$\mathcal{E}_{40,1}$	(1101011010)	2^3	0
$\mathcal{E}_{40,2}$	(1100011000)	2^2	2
$\mathcal{E}_{40,3}$	(0111001110)	2^3	2
$\mathcal{E}_{40,3}$	(0100001000)	2^8	2
$\mathcal{E}_{40,4}$	(0011100111)	2^{13}	2
$\mathcal{E}_{40,5}$	(0111101111)	2^{11}	4
$\mathcal{E}_{40,6}$	(1010010100)	2^8	6
$\mathcal{E}_{40,7}$	(1101011010)	$2^8 \times 3$	6
$\mathcal{E}_{40,8}$	(1000110001)	2^{15}	10
$\mathcal{E}_{40,9}$	(1111011110)	2^{16}	10

4.2. Comparison of the methods over \mathbb{F}_2 for length 64. There are two possibilities for the weight enumerators of extremal Type I self-dual codes of length 64 (hence of parameters $[64, 32, 12]$) ([4]):

$$\begin{aligned} W_{64,1} &= 1 + (1312 + 16\beta) y^{12} + (22016 - 64\beta) y^{14} + \dots, \\ W_{64,2} &= 1 + (1312 + 16\beta) y^{12} + (23040 - 64\beta) y^{14} + \dots \end{aligned}$$

where β and γ are parameters.

Self-dual four circulant $[64, 32, 12]_2$ type I codes exist for weight enumerators $\beta = 0, 8, 16, 24, 32, 40, 48, 56, 64$ and 72 in $W_{64,2}$. We provide codes that we obtained from Corollary 3.2 in Table 4. The results show that the limited version of the generalized four circulant construction gives some codes which do not have four circulant representation (The ones with $\beta = 4, 10, 12, 13, 17, 18, 20, 28, 34$.)

TABLE 4. Type I extremal self-dual codes of length 64 by Corollary 3.2

\mathcal{C}_i	r_A	r_B	r_C	$ \text{Aut}(\mathcal{C}_i) $	β in $W_{64,2}$
\mathcal{C}_1	(101001111)	(0001111001100011)	(1101001110011001)	2^3	4
\mathcal{C}_2	(001101011)	(0101001000001001)	(0101011110111101)	2^4	8
\mathcal{C}_3	(111101110)	(0011110000111101)	(0101011110111101)	2^5	8
\mathcal{C}_4	(111101011)	(0000100001000001)	(0100000011001101)	2^3	10
\mathcal{C}_5	(101101111)	(1000101000101000)	(0010111110100010)	2^4	12
\mathcal{C}_6	(100110100)	(0001001001111111)	(0101011110111101)	2^3	13
\mathcal{C}_7	(100011101)	(1010110001101011)	(1100111001100100)	2^4	16
\mathcal{C}_8	(110001001)	(0010111110100010)	(1100011011111000)	2^3	17
\mathcal{C}_9	(000011110)	(0000011101110111)	(0100000011001101)	2^3	18
\mathcal{C}_{10}	(101101100)	(1111001000100111)	(1100101101001011)	2^4	20
\mathcal{C}_{11}	(000011100)	(0010001101001111)	(0101011110111101)	2^3	24
\mathcal{C}_{12}	(010111001)	(1010000110110000)	(0100010001011111)	2^4	24
\mathcal{C}_{13}	(010010011)	(0000000100000101)	(0101011110111101)	2^4	28
\mathcal{C}_{14}	(111011011)	(0100000101111111)	(0001010001000001)	2^4	32
\mathcal{C}_{15}	(011111000)	(1000101000100010)	(0001001101000011)	2^3	34

Remark 4.1. The first code with weight enumerator $\beta = 34$ in $W_{64,2}$ has been recently constructed in [1]. Here we give an alternative construction.

4.3. Comparison of the methods over $\mathbb{F}_2 + u\mathbb{F}_2$. In this section we compare the methods; four circulant construction and generalized four circulant construction over $\mathbb{F}_2 + u\mathbb{F}_2$ for length 32. A complete classification of four circulant codes of length 32 over $\mathbb{F}_2 + u\mathbb{F}_2$ is given by Karadeniz et al. in [11]. Four circulant type I codes of length 32 have binary images corresponding to weight enumerators with $\beta = 0, 16, 32, 48$ and 80 in $W_{64,2}$. In Table 5 we provide generalized four circulant codes. It is observed that the latter method is more efficient as it produces many more codes of length 64 with parameters that could not be obtained from the ordinary four circulant construction.

TABLE 5. $[64, 32, 12]$ codes via Theorem 3.1 over $\mathbb{F}_2 + u\mathbb{F}_2$ ($W_{64,2}$)

\mathcal{F}_i	r_A	r_B	r_C	$ Aut(\mathcal{F}_i) $	β
1	$(0, u, 0, 0, 1, u, 3, 0)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(0, u, 3, 0, 0, u, 3, 0)$	2^4	0
2	$(u, u, 0, u, 1, u, 1, u)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(0, 0, 3, 0, 0, 0, 3, 0)$	2^5	0
3	$(u, u, u, u, 1, u, 3, u)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(u, u, 1, u, u, u, 1, u)$	2^6	0
4	$(u, 0, u, 0, 0, 1, u, 3)$	$(u, u, u, u, 0, 1, 1, 1)$	$(1, u, 3, 0, 3, u, 1, 0)$	2^3	4
5	$(u, u, u, u, 1, 1, 3, 1)$	$(u, u, 0, 1, 0, 0, 1, 3)$	$(u, u, 0, u, u, u, 0, u)$	2^4	4
6	$(0, u, 0, u, 1, u, 3, u)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(u, 0, 3, 0, u, 0, 3, 0)$	2^5	4
7	$(0, u, 0, u, 1, 1, 0, 1)$	$(u, u, 0, u, 1, 1, 1, 3)$	$(u, 0, u, 0, u, 3, u, 3)$	2^3	8
8	$(u, 0, 0, u, 1, u, 1, 0)$	$(u, u, u, u, 0, 1, 1, 1)$	$(u, 0, 1, 0, u, 0, 1, 0)$	2^4	8
9	$(u, u, u, u, 1, 1, u, 1)$	$(u, u, u, u, u, 1, 0, 1)$	$(u, 0, u, u, 1, 3, 1, 1)$	2^5	8
10	$(u, u, 0, 0, 1, 1, 0, 3)$	$(u, u, 0, u, 1, 1, 1, 3)$	$(u, u, 0, u, u, 1, 0, 1)$	2^3	12
11	$(u, u, u, u, 0, 1, 0, 3)$	$(u, u, u, u, 0, 1, 1, 1)$	$(1, u, 3, u, 1, u, 3, u)$	2^4	12
12	$(0, u, 0, u, 0, 1, u, 3)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(1, 0, 3, 0, 1, 0, 3, 0)$	2^5	12
13	$(u, 0, 0, 0, u, 1, u, 3)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(1, u, 3, u, 1, u, 3, u)$	2^6	12
14	$(u, 0, u, 0, 1, 1, u, 1)$	$(u, u, u, u, u, 1, 0, 1)$	$(0, 0, 0, u, 1, 3, 1, 1)$	2^4	16
15	$(u, u, u, u, 0, 1, 0, 3)$	$(u, u, u, u, 0, 1, 1, 1)$	$(3, 0, 3, 0, 3, 0, 3, 0)$	2^5	16
16	$(u, u, 0, u, u, 1, u, 3)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(3, 0, 3, 0, 3, 0, 3, 0)$	2^6	16
17	$(0, 0, u, 0, 0, 1, 0, 3)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(3, 0, 3, 0, 3, 0, 3, 0)$	2^7	16
18	$(u, u, u, u, 1, 1, u, 1)$	$(u, u, 0, 0, u, 1, u, 3)$	$(u, u, u, 0, 3, 3, 3, 1)$	2^3	20
19	$(u, 0, 0, u, 1, u, 1, 0)$	$(u, u, u, u, 0, 1, 1, 1)$	$(0, 0, 1, 0, 0, 0, 1, 0)$	2^4	20
20	$(u, u, u, u, 1, 1, 0, 1)$	$(u, u, 0, u, 1, 3, 3, 1)$	$(u, u, u, u, u, 3, 0, 3)$	2^3	24
21	$(u, 0, 0, u, 0, 1, u, 1)$	$(u, u, u, u, 0, 1, 1, 1)$	$(3, u, 3, 0, 3, u, 3, 0)$	2^4	24
22	$(u, u, 0, u, 0, 1, 0, 1)$	$(u, u, u, 0, 0, u, 0, 1)$	$(1, 0, 1, 0, 1, 3, 1, 3)$	2^5	24
23	$(u, 0, 0, 0, u, 1, u, 3)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(1, u, 3, 0, 1, u, 3, 0)$	2^4	28
24	$(u, u, u, u, u, 1, 0, 3)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(1, u, 1, 0, 1, u, 1, 0)$	2^5	32
25	$(u, u, 0, 0, 1, 0, 1, u)$	$(u, u, u, u, 0, 1, 1, 1)$	$(u, 0, 3, 0, u, 0, 3, 0)$	2^4	36
26	$(u, u, u, 0, 1, u, 3, 0)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(u, u, 3, u, u, u, 3, u)$	2^5	36
27	$(0, u, u, 0, 0, 1, 0, 1)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(1, 0, 3, 0, 1, 0, 3, 0)$	2^4	44
28	$(0, u, u, 0, 0, 1, 0, 1)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(1, u, 1, u, 1, u, 1, u)$	2^5	48
29	$(u, 0, u, 0, u, 1, 0, 3)$	$(u, u, u, 0, 0, 1, 1, 3)$	$(3, u, 3, u, 3, u, 3, u)$	2^7	80

4.4. **Computational results over $\mathbb{F}_4 + u\mathbb{F}_4$.** Four circulant codes of length 16 over $\mathbb{F}_4 + u\mathbb{F}_4$ had been studied in [12]. The binary images of these codes are Type I extremal self-dual binary codes with weight enumerators $\beta = 0, 4, 8, 12, 24, 28, 32, 36, 40, 48$ and 52 in $W_{64,2}$. We apply Corollary 3.2 and observe that it provides many new parameters that could not be constructed from the four circulant construction. The results are tabulated in Table 6.

TABLE 6. Self-dual codes via Corollary 3.2 over $\mathbb{F}_4 + u\mathbb{F}_4$ of length 16 whose binary images are self-dual codes of length 64

\mathcal{E}_i	r_A	r_B	r_C	$ Aut(\mathcal{E}_i) $	β in $W_{64,2}$
\mathcal{E}_1	$(D, F, 5, F)$	$(E, C, 0, 1)$	$(7, B, 4, A)$	2^3	1
\mathcal{E}_2	$(5, B, D, B)$	(A, E, B, D)	$(7, F, 1, 8)$	2^3	5
\mathcal{E}_3	$(B, 9, D, 9)$	$(2, E, 9, 7)$	$(D, 7, 1, 2)$	2^3	9
\mathcal{E}_4	(D, F, F, F)	$(E, E, 9, 8)$	$(A, 6, 9, 7)$	2^3	13
\mathcal{E}_5	$(9, 7, 7, 7)$	$(0, F, C, 0)$	$(4, 1, F, A)$	2^3	17
\mathcal{E}_6	$(F, 9, 7, 9)$	$(2, 4, 3, F)$	$(F, 5, B, 8)$	2^3	21
\mathcal{E}_7	$(D, 0, F, 0)$	$(9, E, C, A)$	$(D, B, 4, 2)$	2^3	29
\mathcal{E}_8	$(5, 8, 6, 8)$	$(F, 5, E, 8)$	$(0, 8, 9, 7)$	2^5	40
\mathcal{E}_9	$(B, 4, 4, 4)$	$(7, E, 8, D)$	$(0, 6, 4, 2)$	2^5	52

Remark 4.2. The codes with weight enumerators for 1, 5, 13, 17, 21, 29 and 52 were first constructed in [10] as R_3 lifts of the extended binary Hamming code. These are reconstructed by a circulant construction in Table 6.

5. NEW EXTREMAL BINARY SELF-DUAL CODES OF LENGTH 68

The possible weight enumerator of an extremal binary self-dual code of length 68 (of parameters $[68, 34, 12]$) is in one of the following forms by [3, 9]:

$$\begin{aligned}
 W_{68,1} &= 1 + (442 + 4\beta)y^{12} + (10864 - 8\beta)y^{14} + \dots, 104 \leq \beta \leq 1358, \\
 W_{68,2} &= 1 + (442 + 4\beta)y^{12} + (14960 - 8\beta - 256\gamma)y^{14} + \dots
 \end{aligned}$$

where $0 \leq \gamma \leq 9$. Recently, Yankov et al. constructed the first examples of codes with a weight enumerator for $\gamma = 7$ in $W_{68,2}$. Together with these, the existence of codes in $W_{68,2}$ is known for many values. In order to save space we only give the lists for $\gamma = 5$ and $\gamma = 6$, which are updated in this work;

$$\gamma = 5 \text{ with } \beta \in \{113, 116, \dots, 181\} \quad \gamma = 6 \text{ with } \beta \in \{2m \mid m = 69, 77, 78, 79, 81, 88\}$$

We construct 36 new codes with the rare parameter $\gamma = 6$ and 7 codes with $\gamma = 5$ in $W_{68,2}$.

We first construct two new codes of length 68 by applying the extension method described in Theorem 2.3 over $\mathbb{F}_2 + u\mathbb{F}_2$ to \mathcal{F}_2 from Table 5.

TABLE 7. New codes of length 68 as extensions of codes in Table 5 by Theorem 2.3

$\mathcal{D}_{68,i}$	\mathcal{F}_i	c	X	γ	β
$\mathcal{D}_{68,1}$	2	1	$(31u011u30uu113u3333u11u010301101)$	5	101
$\mathcal{D}_{68,2}$	2	$1 + u$	$(130031u300013101313u31uu301u3103)$	5	105

Now, applying the neighboring construction to the codes obtained in Table 7, we get the following new codes of length 68:

TABLE 8. New codes of length 68 as neighbors of codes in Table 7

$\mathcal{N}_{68,i}$	$\mathcal{D}_{68,i}$	x	γ	β
$\mathcal{N}_{68,1}$	2	(1010111001011100010100000010000000)	5	109
$\mathcal{N}_{68,2}$	2	(0000110011011010111101110100011100)	5	111
$\mathcal{N}_{68,3}$	2	(0000111110011111111011010001100000)	5	112
$\mathcal{N}_{68,4}$	2	(1101110000001001011100101100010101)	5	114
$\mathcal{N}_{68,5}$	1	(1110110100000001001100000111001010)	5	115
$\mathcal{N}_{68,6}$	2	(1001010001010101110010111110111000)	6	133

Example 5.1. Let \mathcal{C}_{68} be the code obtained by extending $\varphi(\mathcal{E}_6)$ over $\mathbb{F}_2 + u\mathbb{F}_2$ where $X = (3, 0, 1, 1, u, 0, u, 3, 0, 1, 1, u, 1, 0, 0, u, 3, 0, 0, u, u, u, 1, 3, 3, 0, 1, 3, 1, u, 0, 3)$, then the binary image of \mathcal{C}_{68} is an extremal self-dual [68, 34, 12] code with weight enumerator for $\gamma = 6$ and $\beta = 157$ in $W_{68,2}$. As listed above only 6 codes with $\gamma = 6$ were known before. So this is the first example of a self-dual code with the corresponding weight enumerator.

Without loss of generality, we consider the standard form of the generator matrix of $\varphi(\mathcal{C}_{68})$. Let $x \in \mathbb{F}_2^{68} - \varphi(\mathcal{C}_{68})$ then $D = \langle \langle x \rangle^\perp \cap \varphi(\mathcal{C}_{68}), x \rangle$ is a neighbor of $\varphi(\mathcal{C}_{68})$. The first 34 entries of x are set to be 0, the rest of the vectors are listed in Table 9. As neighbors of $\varphi(\mathcal{C}_{68})$ we obtain 34 new codes with weight enumerators for $\gamma = 6$ in $W_{68,2}$, which are listed in Table 9. All the codes have an automorphism group of order 2.

TABLE 9. New extremal binary self-dual codes of length 68 with $\gamma = 6$ as neighbors of \mathcal{C}_{68}

$\mathcal{C}_{68,i}$	X	β	$\mathcal{C}_{68,i}$	X	β
$\mathcal{C}_{68,1}$	(1111111100111100001100001000000111)	137	$\mathcal{C}_{68,2}$	(0101001001001111111011100010111011)	139
$\mathcal{C}_{68,3}$	(1000001100000110110110000111100010)	140	$\mathcal{C}_{68,4}$	(0010011101110110011001110110110110)	141
$\mathcal{C}_{68,5}$	(1111111111000011111101100010011001)	142	$\mathcal{C}_{68,6}$	(100100000111111111010010000011110)	143
$\mathcal{C}_{68,7}$	(110010001000011100110011111110001)	144	$\mathcal{C}_{68,8}$	(0000110001110110011011011010000110)	145
$\mathcal{C}_{68,9}$	(100001010000101011010111011110101)	146	$\mathcal{C}_{68,10}$	(1100110100000010010000110010011110)	147
$\mathcal{C}_{68,11}$	(111010100001111010010111111101011)	148	$\mathcal{C}_{68,12}$	(0110011001001101000111010101011000)	149
$\mathcal{C}_{68,13}$	(111111110010110100000001011111000)	150	$\mathcal{C}_{68,14}$	(000010000110001011101001111111000)	151
$\mathcal{C}_{68,15}$	(1110000010100000001110110110000101)	152	$\mathcal{C}_{68,16}$	(1010100100110011111101001101001001)	153
$\mathcal{C}_{68,17}$	(1111010010000100100000101000011101)	155	$\mathcal{C}_{68,18}$	(100000101111011110010110000001000)	159
$\mathcal{C}_{68,19}$	(0001010001010101010010010001100010)	160	$\mathcal{C}_{68,20}$	(1100000100011110101111110001010101)	161
$\mathcal{C}_{68,21}$	(0101110011110010110000111111010011)	163	$\mathcal{C}_{68,22}$	(100000011111100000010111100010001)	164
$\mathcal{C}_{68,23}$	(0100000001010000001001110110010110)	165	$\mathcal{C}_{68,24}$	(0111001010010100000010010010101000)	166
$\mathcal{C}_{68,25}$	(1111010011000111000101101001011100)	167	$\mathcal{C}_{68,26}$	(0010110010110100000010001111000000)	168
$\mathcal{C}_{68,27}$	(0000010011010110001010010000101001)	169	$\mathcal{C}_{68,28}$	(1110101000110000011111010101010101)	170
$\mathcal{C}_{68,29}$	(1110100001100111100100000010010010)	171	$\mathcal{C}_{68,30}$	(1000001101101110010001101010111101)	172
$\mathcal{C}_{68,31}$	(1100100001110011101001010001100000)	173	$\mathcal{C}_{68,32}$	(0100011001000011000100010100101100)	174
$\mathcal{C}_{68,33}$	(0110000001110110000111101000101011)	177	$\mathcal{C}_{68,34}$	(101111100010000000101101000101010)	184

6. CONCLUSION

In this paper, we generalize the well known four circulant construction for constructing self-dual codes. We compare both methods to highlight the significance of the generalized construction. Additionally, we construct many new codes of length

68. For codes of length 68, we constructed the following codes with new weight enumerators in $W_{68,2}$:

$$\gamma = 5, \quad \beta = \{101, 105, 109, 111, 112, 114, 115\}.$$

$$\gamma = 6, \quad \beta = \{133, 137, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, \\ 153, 155, 157, 159, 160, 161, 163, 164, 165, 166, 167, 168, 169, 170, 171, \\ 172, 173, 174, 177, 184\}.$$

The binary generator matrices of the new codes we have constructed are available online at [14].

The results we have obtained have demonstrated the effectiveness of the new construction and the difference from the ordinary four-circulant construction. A possible direction for future research could be applying these constructions for different rings and lengths.

REFERENCES

- [1] D. Anev, M. Harada and N. Yankov, “New extremal singly even self-dual codes of lengths 64 and 66”, *J. Algebra Comb. Discrete Appl.*, Vol. 5, No. 3, pp. 143–151, 2017.
- [2] K. Betsumiya, S. Georgiou, T.A. Gulliver, M. Harada and C. Koukouvinos, “On self-dual codes over some prime fields”, *Discrete Math*, Vol. 262, pp. 37–58, 2003.
- [3] S. Buyuklieva and I. Boukliev, “Extremal self-dual codes with an automorphism of order 2”, *IEEE Trans. Inform. Theory*, Vol. 44, pp. 323–328, 1998.
- [4] J.H. Conway and N.J.A. Sloane, “A new upper bound on the minimal distance of self-dual codes”, *IEEE Trans. Inform. Theory*, Vol. 36, No. 6, pp. 1319–1333, 1990.
- [5] S.T. Dougherty, T.A. Gulliver and M. Harada, “Extremal binary self dual codes”, *IEEE Trans. Inform. Theory*, Vol. 43, pp. 2036–2047, 1997.
- [6] S.T. Dougherty, P. Gaborit, M. Harada and P. Sole, “Type II codes over $\mathbb{F}_2 + u\mathbb{F}_2$ ”, *IEEE Trans. Inform. Theory*, Vol. 45, pp. 32–45, 1999.
- [7] S.T. Dougherty, J.-L. Kim, H. Kulosman and H. Liu, “Self-dual codes over Commutative Frobenius rings”, *Finite Fields and Applications*, Vol. 16, No. 1, pp. 14–26, 2010.
- [8] P. Gaborit, V. Pless, P. Sole and P. Atkin, “Type II codes over \mathbb{F}_4 ”, *Finite Fields Appl.*, Vol. 8, No. 2, pp. 171–183, 2002.
- [9] M. Harada and A. Munemasa, “Some restrictions on weight enumerators of singly even self-dual codes”, *IEEE Trans. Inform. Theory*, Vol. 52, pp. 1266–1269, 2006.
- [10] S. Karadeniz and B. Yildiz, “New extremal binary self-dual codes of length 64 from R_3 lifts of the extended binary Hamming code”, *Des. Codes Cryptogr.*, Vol. 74, No. 3, pp. 673–680, 2015.
- [11] S. Karadeniz, B. Yildiz and N. Aydin, “Extremal binary self-dual codes of lengths 64 and 66 from four-circulant constructions over codes $\mathbb{F}_2 + u\mathbb{F}_2$ ”, *FILOMAT*, Vol. 28, No. 5, pp. 937–945, 2014.
- [12] A. Kaya and B. Yildiz “Various constructions for self-dual codes over rings and new binary self-dual codes”, *Discrete Math*, Vol. 339, No. 2 pp. 460–469, 2016.
- [13] A. Kaya, B. Yildiz and A. Pasa “New extremal binary self-dual codes from a modified four circulant construction”, *Discrete Math*, Vol. 339, No. 3 pp. 1086–1094, 2016.
- [14] J. Gildea, A. Kaya and B. Yildiz “Binary generator matrices for extremal binary self-dual codes of length 68”, available online at <http://abidinkaya.wixsite.com/math/research6>.
- [15] S. Ling and P. Sole, “Type II codes over $\mathbb{F}_4 + u\mathbb{F}_4$ ”, *Europ. J. Combinatorics*, Vol. 22, pp. 983–997, 2001.
- [16] E.M. Rains, “Shadow Bounds for Self Dual Codes”, *IEEE Trans. Inform. Theory*, Vol. 44, pp. 134–139, 1998.
- [17] S. Solak and M. Bahsi, “Some Properties of Circulant Matrices with Ducci Sequences”, *Linear Algebra and Appl.*, vol. 542, pp. 557–568, 2018.
- [18] P. Zellini, “On Some Properties of Circulant Matrices”, *Linear Algebra and Appl.*, vol. 26, pp. 31–43, 1979.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHESTER, UK

E-mail address: `j.gildea@chester.ac.uk`

DEPARTMENT OF MATHEMATICS EDUCATION, SAMPOERNA UNIVERSITY, 12780, JAKARTA, INDONESIA

E-mail address: `abidin.kaya@sampoernauniversity.ac.id`

DEPARTMENT OF MATHEMATICS & STATISTICS, NORTHERN ARIZONA UNIVERSITY, FLAGSTAFF 86011, AZ

E-mail address: `bahattin.yildiz@nau.edu`