

Research

Long term analysis of social structure: evidence of age-based consistent associations in male Alpine ibex

Alice Brambilla, Achaz von Hardenberg, Claudia Canedoli, Francesca Brivio, Cédric Sueur and Christina R. Stanley

A. Brambilla (<https://orcid.org/0000-0003-3061-9544>) ✉ (alicebrambilla1@gmail.com), Dept of Evolutionary Biology and Environmental Studies, Univ. of Zurich, Zurich, Switzerland and Alpine Wildlife Research Center, Gran Paradiso National Park, Torino, Italy. – A. von Hardenberg and C. R. Stanley, Conservation Biology Research Group, Dept of Biological Sciences, Univ. of Chester, Chester, UK. – C. Canedoli, Dept of Earth and Environmental Sciences, Univ. of Milano Bicocca, Milano, Italy. – F. Brivio (<https://orcid.org/0000-0002-1449-8335>), Dept of Veterinary Medicine, Univ. of Sassari, Sassari, Italy. – C. Sueur (<https://orcid.org/0000-0001-8206-2739>), Univ. de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France and Inst. Universitaire de France, Saint-Michel 103, Paris, France.

Oikos

2022: e09511

doi: 10.1111/oik.09511

Subject Editor:

Marjo Saastamoinen

Editor-in-Chief: Dries Bonte

Accepted 14 May 2022



Despite its recognized importance for understanding the evolution of animal sociality as well as for conservation, long term analysis of social networks of animal populations is still relatively uncommon. We investigated social network dynamics in males of a gregarious mountain ungulate (Alpine ibex, *Capra ibex*) over ten years focusing on groups, sub-groups and individuals, exploring the dynamics of sociality over different scales. Despite the social structure changing between seasons, the Alpine ibex population was highly cohesive: fission–fusion dynamics lead almost every male in the population to associate with each other male at least once. Nevertheless, we found that male Alpine ibex showed preferential associations that were maintained across seasons and years. Age seemed to be the most important factor driving preferential associations while other characteristics, such as social status, appeared less crucial. We also found that centrality measures were influenced by age and were also related to individual physical condition. The multi-scale and long-term frame of our study helped us show that ecological constraints, such as resource availability, may play a role in shaping associations in a gregarious species, but they cannot solely explain sociality and preferential association that are likely also to be driven by life-history linked physiological and social needs. Our results highlight the importance of long-term studies based on individually recognizable subjects to help us build on our understanding of the evolution of animal sociality.

Keywords: age-based preference, Alpine ibex, *Capra ibex*, fission–fusion dynamic, long term research, social network analysis

Introduction

Understanding the drivers of spatial and temporal interactions between animals is of great importance to some of the most pressing questions in biology, such as understanding the evolution of animal sociality (Couzin and Laidre 2009), determining



www.oikosjournal.org

© 2022 The Authors. Oikos published by John Wiley & Sons Ltd on behalf of Nordic Society Oikos. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

how genetic and cultural information spread within and among inter-connected populations (van de Waal and Bshary 2011), tracking disease transmission (MacIntosh et al. 2012, Marchand et al. 2017, Silk et al. 2019), and predicting the invasion dynamics of introduced species (Fogarty et al. 2011).

In many social species, the process of group formation is highly dynamic, with frequent changes in group size and composition (Grueter et al. 2020). Groups can merge (fusion) or split (fission) over time and space, making group composition a dynamic property (Sueur et al. 2011). Social systems characterized by fission–fusion dynamics are widespread among vertebrates as for example fish (*Poecilia reticulata*, Wilson et al. 2014), birds (Silk et al. 2014, Papageorgiou and Farine 2020) and several mammals including primates (Smuts et al. 1987), Bechstein's bats *Myotis bechsteinii* (Kerth and König 1999), bottlenose dolphins *Tursiops* spp. (Connor and Wells 2000), spotted hyenas *Crocuta crocuta* (Holekamp et al. 1997), African elephants *Loxodonta africana* (Wittemyer et al. 2005), giraffes *Giraffa camelopardalis* (Bond et al. 2019), feral goats *Capra hircus* (Stanley and Dunbar 2013) and sheep *Ovis aries* (Jewell et al. 1974). The timescale over which fission–fusion dynamics occur may differ as environmental conditions or individual requirements change (Aureli et al. 2008, Sueur et al. 2011). Indeed, there is pronounced variation in the degree of fission–fusion dynamics both across and within species which can vary from a highly cohesive society with stable group membership to a highly fluid society with either relatively stable or flexible subgroup membership (Aureli et al. 2008). The flexibility of fission–fusion dynamics is likely to have evolved because it allows individuals to optimize the costs and benefits of group-living (Moscovice et al. 2020). Fission–fusion dynamics are predicted to be most frequent in environments with spatial variability and to increase with temporal uncertainty and unpredictability of the environment (Sueur et al. 2011). Other factors known to influence the structure and cohesion of groups are demographic processes such as deaths, births, dispersal and immigration (İlany and Akçay 2016) that may influence social structure through the loss of some social connections and the formation of new ones (Shizuka and Johnson 2020) as well as by driving changes in patterns of association between remaining individuals (Flack et al. 2006).

Studies on social systems often focus on the association between individuals, as members of a group can show a preference to associate with specific individuals and maintain certain social links despite the frequent splitting and merging of the overall group. In many social mammals there is a tendency towards homophily, i.e. for individuals to associate with others that share similar characteristics such as age or sex (Le Pendu et al. 1995). A possible explanation for these preferences is that the cost of living in groups with members of the same age class or sex is lower due to similar physiological needs and a reduction in harmful inter or intra-sexual interactions (Conradt and Roper 2000, Ruckstuhl and Neuhaus 2000). Likewise, social relationships between individuals may be affected by kinship (Cassinello and Calabuig 2008, Wittemyer et al. 2009, Podgórski et al. 2014) as well as

by social rank as observed in horses *Equus caballus* (Kimura 1998) and red deer *Cervus elaphus* (Appleby 1983), where individuals spent more time close to other of similar rank.

Associations can be stable or can change during the life on an individual, varying from long-term or even life-long associations (as shown, e.g. in birds, Teitelbaum et al. 2017; wild boars *Sus scrofa*, Podgórski et al. 2014; feral horses and ponies, Cameron et al. 2009, Stanley et al. 2018) to transitory or short-term associations (e.g. in spotted hyenas, Smith et al. 2011). Also the levels of gregariousness can differ between individuals of the same species or population and can be influenced by various factors such as age (Carter et al. 2013, Machanda and Rosati 2020) or reproductive status (Fischhoff et al. 2009, Vander Wal et al. 2015, Machanda and Rosati 2020). This can, however, also be a consistent individual characteristic that is maintained throughout life as recent studies on personality (i.e. the presence of behavioural differences that are repeatable over time and across situations) have shown (Krause et al. 2010, Firth et al. 2018). Sociability is indeed one of the five personality traits described in literature and it can affect reproductive success and survival and hence fitness (Réale et al. 2007).

Social network analysis (SNA) is a powerful tool that can be applied to the analysis of several aspects of social behaviour (Sueur et al. 2011, Sosa et al. 2021b). One of the most attractive features of SNA is that it allows to study the social organisation of animals at all levels (individual, dyad, group, population) and for all types of interaction (e.g. aggressive, cooperative, sexual, Krause et al. 2009), allowing a plethora of novel insights into the evolution and maintenance of sociality to be elucidated (Wey et al. 2008, Pinter-Wollman et al. 2014, Krause et al. 2015, Webber and Vander Wal 2019, Sosa et al. 2021b). As social structure can also affect population growth rates, dispersal and gene flow, network analysis also has the potential to be an important tool in the management of wild populations (Tarlow and Blumstein 2007, Schakner et al. 2017, Snijders et al. 2017, Welch et al. 2020).

To date, few studies have investigated dynamic changes in social network structures of wild populations in the long term, especially where the individuals composing the network change due to demographic processes (e.g. death, emigration or immigration of individuals, Cantor et al. 2012, Borgeaud et al. 2017). Shizuka and Johnson (2020) called for the need to integrate demographic processes and consequent social processes into social network analyses. However, the scarcity of such studies is probably due to the intensity of consistent data collection required in order to incorporate environmental and demographic changes or stochastic events, in addition to limitations in the available analytical tools (Pinter-Wollman et al. 2014). Instead, many studies making use of SNA in free ranging populations rely on relatively short-term datasets and, therefore, are only able to describe snapshots of the social systems. Since long term monitoring of wild species has provided important contributions to the study of ecology and evolution as well as to conservation (Festa-Bianchet et al. 2017, Clutton-Brock 2021), long term analysis of social network dynamics could bring rich

rewards in terms of a better understanding of population-level processes (Pinter-Wollman et al. 2014).

Ungulate species are widespread in all continents except Antarctica and show high diversity both in terms of size, habitat as well as behaviour (Wilson and Mittermeier 2011). Their social organization has been extensively studied and provided crucial elements for understanding the evolution of vertebrates' social systems (Jarman 1974, Krause et al. 2002). However, many of the studies on social networks of ungulates have focused on female associations (Vander Wal et al. 2016, Ramos et al. 2019) meaning knowledge on male sociality in ungulates remains scarce. To fill those gaps, we performed a long-term analysis of male social structure in a gregarious ungulate: the Alpine ibex *Capra ibex*. We took advantage of a detailed long-term dataset resulting from ten years of behavioural observations of individually identifiable male Alpine ibex to explore the male social network of a gregarious ungulate and to investigate population-level changes in social structure over time.

The Alpine ibex, a mountain ungulate of conservation concern (Brambilla et al. 2015, 2018, 2020), is a gregarious species that lives in open-membership groups and shows fission–fusion dynamics. Alpine ibex are highly sexually dimorphic and exhibit strong sexual segregation all year round except for during the mating season, that occurs between December and January (Villaret and Bon 1995, Ruckstuhl and Neuhaus 2001). As the degree of sexual segregation increases with male age (Ruckstuhl and Neuhaus 2001), outside the rut, adult males join groups mainly composed of males, while yearlings stay in groups mainly composed of females, and young males (≤ 2 –3 years old) may move between male and female groups.

This study was conducted during spring and summer months, when sexual segregation is at its peak, and focused on the social network of male Alpine ibex. Specifically, we aimed to determine: 1) whether males showed consistent associations and which factors influenced the choice of preferred companions; 2) which individual characteristics (age, life stage) influenced network metrics at the individual level and if these metrics were consistent over time; 3) whether social structure was stable across seasons (i.e. across periods with different ecological conditions); 4) whether any changes in the social network structure over time could be explained by demographic factors.

Similarly to observations in other bovids (Vander Wal et al. 2015, Ramos et al. 2019), we expected to find a highly connected and cohesive social network, with all the individuals connected either directly or indirectly. Despite the aforementioned studies having analysed social structure in females, based on field observations we expected similar results also for males. Age was expected to predict certain network node-based measures (Turner et al. 2018, Sosa et al. 2021b). Particularly, we expected adult males to occupy the more central positions in the network due to their reproductive status and generally higher rank. In addition, we expected males to become less central once they had passed their reproductive peak due to their decreased

competitive ability. Age often underpins social preferences (Wey and Blumstein 2010, Welch et al. 2020); we therefore expected stronger associations between males of the same age class due to their similarity in body size, social motivation and behaviours. At the same time, and as rank-based homophily was observed in other species (Sosa 2016), we also expected stronger associations between males with similar social rank.

At the population level, we wanted to verify if the structure of the social network remained stable between seasons and between years over a period of ten years. Environmental conditions during the study period were reasonably stable within seasons so we did not expect variations in global network measures. However, as more experienced, older individuals are expected to help maintain group cohesion (Allen et al. 2020), we expected to find a correlation between global centrality measures and the proportion of old males in the population which could change due to stochastic events (e.g. harsh winters).

Methods

Study area and population

The study was conducted on a free-ranging population of Alpine ibex in the Levionaz basin (Valsavarenche valley, AO), within the Gran Paradiso National Park (north-western Italian Alps – 45°26'N, 7°08'E).

In the study site, ibex are captured and individually marked with coloured plastic ear tags in the framework of a long-term study on the life-history and conservation of the species (Bassano et al. 2003, Bergeron et al. 2010, Brambilla and Brivio 2018). The capture and marking protocol used in this study has been authorised by the Italian Ministry of Environment (authorisation no. 25114 of 21/09/2004) after the positive review by the Italian National Institute for Environmental Protection and Research (ISPRA) and was developed trying to minimize the effects on the welfare of the animals (Brambilla et al. 2013, Brivio et al. 2015).

The mean number \pm SD of individuals counted in the Levionaz population during the 10 years of this study was 180 ± 41 of which 63.9 ± 10.8 were males. Censuses were conducted every year in July block-counts by the personnel of Gran Paradiso National Park. The proportion of marked males varied from a minimum of 50% in 2017 to a maximum of 88% in 2011 and 2012 for a total of 111 marked individuals observed during the study.

Data collection

Data were collected for male Alpine ibex during a period of ten years from 2008 to 2017 in spring (May–June) and summer (July–Aug–Sept). As Alpine ibex perform yearly altitudinal migrations following the green up of the vegetation (Parrini et al. 2003), exact dates of the seasonal ranges varied between years and were selected according to the altitudinal movement of the animals on the study site.

Individual attributes

Age

The age of the marked individuals was determined at capture by counting the number of annual horn segments (Brambilla and Canedoli 2014). The age class of the unmarked individuals was estimated by using binoculars to count the number of annual horn segments. Unmarked individuals were divided in age classes, defined as follows based on clearly visible body mass differences (Couturier 1962): 2–5 years old (young), 6–8 years old (sub-adults); 9–11 years old (fully grown adults); > 11 years old (old individuals).

Season preceding death

Male Alpine ibex adult survival is very high and most of them reach senescence (Toigo et al. 2007). However, some individuals die earlier and, particularly in case of chronic diseases, their behaviour could change in the months preceding death. We therefore recorded all deaths of identified individuals to obtain a binary variable ‘season preceding death’; this indicated whether the animal died within that year of data collection or if it survived until the following year. This was possible as carcasses of dead individuals were often found in the field within a few days of death. When possible, the cause of death was also ascertained. Some individuals’ carcasses, however, were not found. This happened especially when animals died during winter or in an inaccessible part of the study area. If an animal was not observed in the study site nor in the surrounding areas (monitored daily by park rangers) for a whole year, we considered it as dead. We could not exclude that some of those individuals migrated; however, as all the animals included in our study were observed several times every year and their movements were rather predictable, we were confident in considering them as dead if they were not observed again. Indeed, during the time of our study, we had no cases of animals that were observed again in our or in other study areas after one whole year when they were not observed.

Male associations

Association patterns between individuals were defined via the ‘gambit of the group’ method (Whitehead 2008), i.e. assuming that each animal in a group is associating with every other individual in that group (Croft et al. 2008, Franks et al. 2010). Association data were collected daily by means of surveys conducted when the animals were active/feeding. When possible, associations were recorded twice per day, after dawn and before dusk, considering each survey as independent due to the strongly bimodal daily activity pattern typical of the species, but time and number of daily observations varied based on seasonal and weather conditions (details on the number of surveys per season are provided in the Supporting information). Data were collected by an observer scanning the study area while walking along transects consisting of the GPNP paths used during block counts. Transect routes changed during the season according to the altitudinal movement of the animals in the study site (Parrini et al. 2003). Each session of data collection lasted around one to three hours depending on the location of the animals. Every year,

data were collected by two to four observers. New observers were trained and tested by the same person that was present for the whole duration of the study. Before new observers could start collecting data, blind contemporary data collection with the expert observer was conducted for up to one week, until consensus in group identification was reached. For each group, the total number of individuals, identity of the marked individuals and age class of the unmarked individuals were recorded. Two groups were considered as distinct if their closest members were more than 50 m apart. This threshold was set after field observations considered this distance sufficient to avoid social interactions during random movements while foraging. Any animal observed alone was considered to be a separate group. Data on association were used to build association matrices for each season of each year.

Male dominance

All observed agonistic interactions when an individual was clearly dominant over another, as defined by Bergeron et al. (2010), were recorded using all occurrence sampling (Altmann 1974). Agonistic interactions were collected opportunistically during the group composition data collection as well as during focal samples conducted for other studies. The identity of the winner and loser in each interaction was recorded, allowing dominance matrices to be built based on the frequency of dominance interactions per dyad. If the identity of one or both of the individuals involved in the interaction was unknown, this interaction was not included in the matrix.

The outcome of agonistic interactions was also used to calculate hierarchical rank via the Elo-rating method (Elo 1978, Albers and De Vries 2001, Neumann et al. 2011). This method is based on the sequence in which interactions occur and continuously updates ratings by looking at interactions sequentially (Neumann et al. 2011). As previous analysis showed that hierarchical ranks are established early in spring and remain rather stable during summer, we calculated the Elo score (R package EloRating, Neumann et al. 2011) for each individual as of the 30 June of each year in which the individual was observed.

Network analysis

Association networks

The association network was built based on the half weight index (HWI; Whitehead 2008). HWI is defined as:

$$\text{HWI} = \frac{X}{X + 0.5(Y_a + Y_b) + Y_{ab}} \quad (1)$$

where X is the number of sampling periods during which individuals a and b were observed together, Y_a is the number of sampling periods when a was observed without b , Y_b is the number of sampling periods in which b was observed without a , and Y_{ab} is the number of sampling periods when a and b were both observed separately (Whitehead 2008). The HWI is considered to be less biased when not all associating individuals can be identified (Whitehead 2008) and it was

chosen as association index to account for the unequal detection probability typical of gregarious mountain ungulates (Vander Wal et al. 2015) and for potential uneven sampling between individuals. As ecological conditions (e.g. resource quality and distribution, temperatures, precipitations) change during the year and Alpine ibex perform seasonal altitudinal migrations (Parrini et al. 2003), the HWI for each pair of individuals was calculated separately for spring and summer of each year (2008–2017) resulting in 20 time-aggregated association matrices (for details on the methods and tools used to build the networks, see the data analysis section).

Dominance networks

Dominance networks were built using the absolute frequency of dominance interactions between each pair of individuals within each season and year (as we assume that our method allowed random sampling of dominance interactions), resulting in 20 dominance networks. Dominance networks were non-symmetrical as the directionality of the dominance resulting from the interaction was preserved.

Network metrics

To analyse the network structure of male Alpine ibex, we used the metrics described below.

Individual-level metrics

Strength centrality and eigenvector centrality were calculated for each individual in each network and averaged among individuals for each network. Strength centrality is calculated as the sum of the weights of the edges in a weighted network (Wasserman and Faust 1994). This measure represents the sociality of an individual as it estimates the frequency of its interactions (Sosa et al. 2021a) and it was also used to compute gregariousness at the group level. Eigenvector centrality is defined as the first non-negative eigenvector value obtained by transforming an adjacency matrix linearly (Wasserman and Faust 1994). It measures the centrality of a node by examining its connections as well as that of its alters (with alters being the other individuals connected to the node). Eigenvector centrality can therefore be interpreted as the social resources available to an individual (Brent et al. 2011).

Global metrics

As measures to describe the global structure of the network we calculated network density, gregariousness and typical group size (TGS). Network density calculates the ratio between existing links and all potential links of a network, and it assesses the connection of the network as a whole (Sosa et al. 2021a). Gregariousness represents the tendency of individuals to associate with few or many individuals (Godde et al. 2013). At the individual level, it is represented by strength centrality and it is calculated as the sum of the values of the association indices involving that individual. The overall gregariousness of a population is calculated as the average gregariousness of all individuals in the population. TGS quantifies group size as experienced by an average individual of the population and it emphasises the extent to which members of the population tend to associate

(Jarman 1974). It is an animal-centred measure defined as the sum of the squares of the number of individuals in each group, divided by the total number of animals sampled (Jarman 1974).

Data analysis

Association matrices were built in SOCPROG ver. 2.9 (Whitehead 2009). Network analysis was performed in R (<www.r-project.org>) using the packages *tnet* (Opsahl 2009), *network* and *sna* (Butts 2020), included in the *statnet* suite (Statnet Development Team 2003–2020, Handcock et al. 2008) and with the package *ANTs* (Sosa et al. 2020). Generalised linear mixed models (GLMMs) were conducted using the R packages *ANTs* and *lme4* (Bates et al. 2015).

Data and code are accessible on the Dryad Digital Repository <<https://doi.org/10.5061/dryad.w0vt4b8st>>.

Association patterns

To investigate association patterns, we used a quadratic assignment procedure (QAP) approach to compare pairs of association networks. The QAP test is a specialised version of the Mantel test that uses random permutations of node labelling to determine whether a correlation between two matrices is significantly higher than expected (Krackhardt 1988, Krause et al. 2015). QAP tests were carried out with 10 000 permutations using the *sna* package.

To test for consistency in associations between seasons, a QAP test was carried out between the two seasonal association networks within each year. For this analysis, the networks were built excluding individuals that were not observed across both seasons within a year.

Furthermore, to test for consistency in associations between years, we compared the networks built for the summer seasons of the years 2012–2016. The network used for this analysis only included a subset of individuals that were observed in all those five years (hereafter referred to as subset networks). This time period was selected as being optimal in terms of having the largest number of individuals present across the entire period, hence allowing the greatest power for testing for consistency in associations; the summer seasonal network was selected as spring associations may partly be driven by resource availability. Moreover, spring and summer networks within each year were correlated. QAP tests were carried out between each subset network and the subset network of the following year.

To test whether the age difference between pairs of individuals had an impact on whether they were likely to associate or to be dominant over each other, we built matrices of difference in age (in years) within each dyad for each year. We built both a symmetrical matrix with absolute age difference as well as an asymmetrical matrix with the exact age difference between dyads. We then used QAP tests to determine whether there was significant consistency in structure between the summer association matrix and the absolute age difference matrix, then between the dominance matrix and the age difference matrix. Finally, to test whether dominance rank differences between individuals predicted their likelihood of association, we built

matrices of absolute difference in their Elo scores (used as a proxy for hierarchical rank) and performed a QAP test to test for structural consistency between the summer association matrix and the absolute Elo score difference matrix. As this resulted in multiple hypothesis testing, for each of the sets of QAP tests described before, we applied a sequential Bonferroni correction (Holm 1979) for the assessment of significance levels. A combined p-value for all years was calculated using Fisher's method with the package `poolr` (Cinar and Viechtbauer 2021).

Factors affecting node measures and seasonal network structure

To determine whether association network structure changed between seasons (i.e. between periods with different ecological conditions) and which individual characteristics predicted node-based measures, we carried out generalized linear mixed models (GLMMs) on permuted association matrices with the package `ANTs`: time-aggregated networks for each season of each year were built through data stream permutations resulting in a list of 20 symmetrical association matrices (each with 10 000 permutations) used to calculate the centrality measures of interest (strength and eigenvector centrality) and to run permuted GLMMs.

The fixed and random structures of the permuted GLMMs were selected using non-permuted GLMMs (`lme4` package): each GLMM included as the dependent variable either strength centrality (modelled with a gaussian distribution) or eigenvector centrality (modelled with a binomial distribution) and, as fixed effects, the season and individual characteristics: age as a quadratic term (Bergeron et al. 2008) and the season preceding death (as a binary variable that indicated whether the animal died within that year of data collection or if it survived until the following year). The models also included year and individual identity (ID) as random effects. The choice of the fixed effects as well as the assessment of the importance of individual ID as a random effect was made based on the Akaike information criterion (AIC, Akaike 1973, Burnham et al. 2011), with $\Delta AIC > 2$ used as a threshold for the selection of the best fitting models. Graphical analysis of the residuals and the coefficient of determination R^2 were used to check model fit. Both marginal (R^2_m , that describes the proportion of variance explained by the fixed factors alone) and conditional coefficient of determination (R^2_c , which describes the proportion of variance explained by both the fixed and random factors) were calculated with the R package `MuMIn` (Barton 2009).

Seasonal and annual changes in the social structure

To investigate seasonal differences in the social structure, we run linear models to compare the spring and summer values of the global measures (mean gregariousness and TGS, calculated in R with own-built functions). To account for the sampling effort, the number of surveys for each season was added as an explanatory variable to the model.

To detect possible changes in the network structure during the ten years of the study, we calculated the global measures of density (calculated within the network R package), gregariousness and TGS within each year (across seasons).

As we hypothesised that the age structure of the population could affect the cohesion of the social structure and hence the above-mentioned global network measures, we calculated the proportion of adult individuals (i.e. of individuals of nine years and more) and the total number of males in the population (counting the maximum total number of males, including both marked and unmarked individuals, observed during the daily observations). We then ran two separate generalised linear models (GLM) with binomial distribution to test the effect of the proportion of old individuals and of the total number of males on density. In addition, we ran three GLMs to test the effect of the proportion of old individuals and of the total number of males on TGS and to test the effect of the total number of males on gregariousness.

Results

The association networks of Alpine ibex showed a high level of connectivity (Fig. 1) with the network density being close to one in all the years of the study.

Association patterns

QAP tests performed to test for consistent associations across seasons within each year showed significant correlations between association matrices within all the ten years of the study (Table 1) indicating that male Alpine ibex maintain consistent bonds throughout the year, despite changes in environment and location. Consistent associations between adjacent years were also observed over a five year period as shown by the results of the pairwise QAP tests performed on the subset networks between summer association networks in adjacent years (Table 2).

We also found that dominance matrices mostly correlated positively with age difference matrices within the summer season for each year (Table 3a), indicating that dominance interactions were more likely to occur between individuals the greater their difference in age, with the older individual of the dyad being more likely to be dominant over the younger. Association matrices, instead, mostly correlated negatively with their associated absolute age difference matrices, showing that individuals were more likely to associate with those closest in age (Table 3b). Finally, the tests between association matrices and absolute difference in Elo score matrices showed no evidence of a correlation between association choice and dominance rank differences or similarities (Table 3c).

Factors affecting node measures and seasonal network structure

The results of the GLMMs, performed to test whether the association network structure changed between seasons and which individual characteristics predicted node-level centrality measures, are presented in Table 4 and 5. Model selection results are presented in the Supporting information.

Strength centrality was significantly lower in the summer season compared to spring and was correlated with age

Table 3. Results of QAP tests for correlations between (a) dominance matrices and their associated exact age difference matrices (combined p-value < 0.001); (b) association matrices and their associated and absolute age difference matrices (combined p-value < 0.001); (c) association matrices and their associated absolute Elo score difference matrices. QAP tests were performed only on the summer season matrices. Significant correlations after a Bonferroni sequential correction are indicated by bold p-values.

Year	(a) Dominance versus age-diff.		(b) Association versus abs. age-diff.		(c) Association versus abs. Elo score-diff.	
	Correlation coefficient	p-value	Correlation coefficient	p-value	Correlation coefficient	p-value
2008	0.078	0.039	-0.049	0.304	0.148	0.131
2009	0.233	0.033	-0.239	0.009	0.106	0.166
2010	0.063	0.119	-0.274	< 0.001	0.158	0.038
2011	0.068	0.028	-0.285	< 0.001	-0.239	0.010
2012	0.108	0.009	-0.320	< 0.001	0.004	0.544
2013	0.160	< 0.001	-0.302	< 0.001	-0.119	0.053
2014	0.138	0.004	-0.070	0.230	0.043	0.342
2015	0.247	< 0.001	-0.261	< 0.001	-0.036	0.334
2016	0.188	< 0.001	-0.158	0.007	-0.103	0.114
2017	0.108	0.007	-0.295	< 0.001	0.059	0.287

coefficients of determination of the selected model were: $R^2_m = 0.12$, $R^2_c = 0.55$.

Seasonal and annual changes in the social structure

Annual and seasonal measures calculated to describe the global structure of the network are presented in Table 6. The density of the network was close to one in all the years of the study with little variation (min.=0.82, max.=0.99). TGS and gregariousness showed more variation over the years and between spring and summer seasons with both TGS ($\beta = -6.66$, $SE = -2.05$, $p = 0.005$) and gregariousness ($\beta = -5.09$, $SE = 1.43$, $p = 0.002$) being significantly lower during summer than during spring seasons (with no effect of sampling effort in either model). Gregariousness also showed differences in the coefficient of variation between years (min=0.18, max=0.40). Neither the proportion of old individuals nor the total number of males in the population had a significant effect on density or gregariousness. The proportion of old individuals also had no effect on TGS.

Discussion

We analysed the social network structure of male Alpine ibex using association and behavioural data collected over ten years. Male and female Alpine ibex are strongly segregated all year round, except for during the rutting season that occurs

Table 4. Output of the permuted GLMM performed to explain the variance of strength centrality. Perm p-value represents the permuted p-values obtained after 10 000 permutations in ANTs. Model specification: strength ~ age+age²+season+season preceding death+(1|year)+(1|ID), family=gaussian. Significant explanatory variables are provided in bold.

	Estimate	SE	t-value	Perm p-value
(Intercept)	10.69	2.42	4.41	< 0.001
age	2.93	0.41	7.08	< 0.001
age²	-0.18	0.02	-7.47	< 0.001
season	-9.63	0.36	-26.87	0.022
season prec. death	-2.72	0.65	-4.18	0.001

in December–January (Ruckstuhl and Neuhaus 2001). As they do not associate during spring and summer, our study did not consider interactions between males and females and focused on the social network structure of males alone.

Association patterns

The association network for male Alpine ibex formed one discrete component, with a high density of connections. Fission–fusion dynamics lead almost every male in the population to associate with each other male at least once by being members of the same sub-group; field observations showed frequent splitting and merging events and that all males occasionally joined together in a single group. For example, during bad weather (i.e. mainly thunderstorms), male ibex tend to group at specific places used as shelters, usually at a lower altitude (Brambilla unpubl.). The Alpine ibex has a relatively small home range (Parrini et al. (2003) reported an average yearly home range of around 450–500 ha for males in the same study area) and lives in open high-altitude alpine habitat. During summer, males feed in large alpine pasturelands where resources are not concentrated in specific sites making resource attraction unlikely to explain the high network density observed in this species. The high density of connections observed in male Alpine ibex was therefore probably due to the general gregariousness of the species (Villaret and Bon 1998) that is also observed in other mountain ungulates with a similar social structure and mating system (e.g. bighorn

Table 5. Output of the permuted GLMM performed to explain the variance of eigenvector centrality. Perm p-value represents the permuted p values obtained after 10 000 permutations in ANTs. Model specification: eigenvector ~ age+age²+season+season preceding death+(1|year)+(1|ID), family=binomial. Significant explanatory variables are provided in bold.

	Estimate	SE	z-value	Perm p-value
(Intercept)	-0.97	0.71	-1.37	0.137
age	0.76	0.17	4.51	0.060
age²	-0.05	0.01	-4.89	0.009
season	-0.54	0.18	-2.93	0.003
season prec. death	-0.83	0.25	-3.29	0.030




Figure 2. Partial regression plots representing the relationship between (a) strength and (b) eigenvector centrality respectively with age. Plots were built with the package visreg (Breheny and Burchett 2017). The y-axis is scaled around the mean value of the response variable.

sheep). The Alpine ibex is a polygynous species with males actively tending oestrus females and preventing subordinate males from approaching them (Willisch and Neuhaus 2010, Willisch et al. 2012, Apollonio et al. 2013). As the rutting season of most mountain ungulates occurs in winter, when environmental conditions can be harsh and moving in the snow is energetically demanding (Signer et al. 2011), the establishment of the hierarchies often begins earlier. Indeed, agonistic interactions in Alpine ibex occur all year round except immediately after the rutting season (Brivio et al. 2010) and are more frequent during spring and summer months. Being part of a highly connected network may allow males of polygynous species living in highly seasonal environments to interact with many other males reducing the need of agonistic contests during winter, when it is more energetically costly. This advantage may compensate for the costs of group-living that, in the case of gregarious ungulates, are probably mostly related to infection risk (Brambilla et al. 2013, Marchand et al. 2017) as highly connected networks allows for potentially fast spreading of diseases (Wey et al. 2008, Marchand et al. 2017).

We found significant correlations between association matrices for spring and summer seasons within all the ten

years of the study and also between adjacent years. The correlation coefficients were rather low, requiring caution in the interpretation of these results. However, the fact that all correlations between spring and summer association matrices of the same year and between summer association matrices of adjacent years were significant, seems to indicate that there are consistent associations within the population, both across seasons and years. These stable associations must convey some advantage as they are maintained despite a negligible risk of predation in this population and despite seasonal changes to the global association network structure. Our analyses carried out across a ten year period revealed that the factors leading to assortativity in Alpine ibex appeared constant throughout the study period. Males tended to associate with individuals of a similar age; dominance rank did not seem to predict associations, but difference in age did predict the likelihood of engaging in dominance interactions. The finding that age predicts association confirmed our expectation and supports the results of previous studies (*Capra ibex*, Villaret and Bon 1995; *Capra pyrenaica*, Alados 1986). However, we were not able to disentangle whether preferential associations were also present within individuals in the same age class and which factors drove these associations (e.g. kinship as observed by

Table 6. Global measures calculated for the 2008–2017 networks. Density was calculated for the summer networks. Gregariousness (with coefficient of variation, CV, calculated as the ratio of the standard deviation to the mean) and TGS were calculated for both spring and summer networks.

Year	n tot	Proportion ind > 9 years	Density	Gregariousness spring mean (CV)	Gregariousness summer mean (CV)	TGS spring	TGS summer
2008	59	0.32	0.89	6.01 (0.20)	3.09 (0.40)	12.03	8.17
2009	55	0.31	0.82	5.68 (0.34)	2.74 (0.32)	14.72	11.65
2010	60	0.29	0.95	9.36 (0.26)	6.85 (0.25)	15.65	13.16
2011	57	0.26	0.92	7.70 (0.28)	5.58 (0.36)	11.90	10.08
2012	58	0.33	0.95	12.66 (0.22)	7.59 (0.28)	18.67	12.93
2013	61	0.43	0.89	12.26 (0.21)	6.55 (0.31)	20.50	9.90
2014	58	0.45	0.99	9.20 (0.20)	6.76 (0.23)	17.38	13.42
2015	63	0.40	0.98	16.51 (0.19)	6.19 (0.27)	27.07	11.27
2016	84	0.25	0.96	18.07 (0.18)	9.20 (0.27)	27.83	18.84
2017	84	0.19	0.99	11.56 (0.26)	6.84 (0.29)	22.65	16.71

Cassinello and Calabuig (2008) in aoudades). A possible explanation of the fact that age predicts association is that individuals with similar characteristics (e.g. age and hence body size, Bergeron et al. 2010) also have similar energetic requirements and this leads them to associate and synchronize their activity so as to maintain group cohesion (Conradt and Roper 2000, Ruckstuhl and Neuhaus 2000). The large differences in body size of male Alpine ibex of different ages (Bergeron et al. 2010) further strengthen this explanation and in addition suggests that life history as well can contribute to the evolution of social structures. Furthermore, individuals may associate with others of similar age or size because they share similar social motivations and behaviours that enhance social cohesion and increase association (Cransac et al. 1998, Bon et al. 2001). In a polygynous species with male–male competition that lives in a strongly seasonal environment such as the Alps, adult males may share the need to interact to establish hierarchies before the reproductive season (Willisch and Neuhaus 2009, Brivio et al. 2010). At the same time, young males may share the need to interact with other males through sparring to increase fighting skills, as observed for example in African elephants (Chiyo et al. 2011).

In this Alpine ibex population, groups of males of similar age appear to merge and split with other groups regardless of their age. An implication of this could be that there is not a constant transfer of information between old and young males. In this species, the transfer of information between old and young individuals, if present, is likely to be more important for females that have to learn where to find safe and suitable places for parturition and the weaning of calves. In the case of males, since resources are homogeneously distributed during the summer season and predation risk is negligible, young individuals may avoid the costs of being constantly associated with mature males (with different time activity rhythm and physiological needs) but yet join them on some occasions.

As expected, and not surprisingly, age difference was also positively correlated with an increased likelihood of dominance interactions. Adult individuals were more frequently dominant, as has also been observed in other studies on the same species and on other polygynous mountain ungulates (Pelletier and Festa-Bianchet 2006, Bergeron et al. 2010, Willisch et al. 2012, Apollonio et al. 2013). The Alpine ibex is a capital breeder with a slow life history strategy and reaches social sexual maturity at around 9–10 years of age (Willisch et al. 2012). Young individuals are therefore often in the lower positions of the hierarchy (Bergeron et al. 2010, Willisch et al. 2012). However, we did not find a correlation between likelihood of association and dominance rank differences, indicating that dominance rank is not a factor that seems to predict social preferences. Only a few of the agonistic interactions among males escalate into physical contact. When an individual's dominance is clear, most of the interactions consists of threat displays. Therefore, associating with individuals of higher rank may not be costly, in terms of a risk of being injured, for low-rank males of species that evolved threat displays as a mean to establish hierarchies.

Factors affecting node measures and seasonal network structure

Analysing the network at the node level, we also found that age predicted the position of the individuals in the network; as we were expecting, both strength centrality and eigenvector centrality showed a significant quadratic relationship with age. We found lower centrality values for young and old individuals and maximum centrality values for adult individuals at the peak of fitness (9–10 years old). Young individuals may join groups of adults but then leave and form separate groups, thus explaining the lower centrality values observed in this age class. Moreover, in accordance with the known life history of the species, old and senescent individuals also showed lower strength centrality and eigenvector centrality values. Because of their conservative life history strategy, adult survival is very high in Alpine ibex (Toigo et al. 2007). Many males therefore reach senescence with progressively deteriorating body condition (Bergeron et al. 2008, Brambilla et al. 2015) and changes in activity rhythm and spatial behaviour, meaning they are less likely to remain central in the network. Our results suggests that social behaviour changes in Alpine ibex may occur as a result of senescence (Siracusa et al. 2022). Similar patterns were recently observed in red deer where Albery et al. (2021a, preprint) found that older animals changed their spatial behaviour and became less socially connected. In accordance with our interpretation, we also found that strength and eigenvector centrality were lower if an animal died within subsequent months. Further evidence for this explanation came from field observations as well as from the structure of the network graphs (Fig. 1), where it appears that some individuals occupied a more peripheral position in the months preceding their death (Fig. 1, Supporting information). In particular, this seemed to happen for individuals that died from medium to long term diseases (e.g. pneumonia). In the months preceding death, those individuals were often observed alone, away from the other males. This likely happened because the progressive deterioration of their physical condition did not allow them to follow the daily movements and rhythm of the other individuals. It has been observed that Alpine ibex significantly decrease investment in horn growth in the years preceding death (von Hardenberg et al. 2004) suggesting that energetic investment in non-vital traits is reduced when the animal is in poor physical condition. Our observations suggest that the same is also true for behavioral traits. The observation of changes in individual behavior due to physical conditions and health status may have important implications for understanding the spreading dynamic of infectious diseases (Craft 2015, Silk et al. 2019). However, we were not able to establish the cause of death for all the individuals and therefore our interpretation of this result remains speculative; a more detailed collection of data on the presence of infectious disease could help shed light on this important issue.

We also found preliminary evidence that inter-individual differences other than age are likely to play a role in determining social network positions in male ibex.

Interestingly, an individual's identity concurred to explain their association network position in terms of their strength centrality, suggesting that sociability varies consistently between individuals independent of their age. The presence of behavioural differences that are repeatable over time and across situations (i.e. personality or temperament) in animals has become widely recognized in the last few decades (Réale et al. 2007) and, since then, it has been observed in several species, including mountain ungulates (e.g. in bighorn sheep, Réale et al. 2000, Réale and Festa-Bianchet 2003, Poissant et al. 2013). Sociability is one of the five temperament traits described in literature (Réale et al. 2007) and our ten-year dataset provided a unique opportunity to follow the associations of several individuals for many years, highlighting the potential impact of personality on network position. However, we have only found very preliminary evidence of repeatable individual differences in sociability (i.e. retention of the individual identity as a random effect in the selected model) while other measurements of sociability as well as its heritability should be assessed in order to reliably quantify personality in this species (Réale et al. 2010).

Seasonal and annual changes in the social structure

Analysing seasonal differences in the social structure, we found that mean strength and mean eigenvector centrality in the association network, gregariousness and typical group size (TGS) were all significantly lower in the summer compared with the spring. The overall higher values of sociality observed during spring compared to summer can partly be explained by seasonal differences in resource availability (Peignier et al. 2019). During spring, resources are more concentrated as there are only a few pastures available at low altitude, where snow melts earlier. Hence, all male Alpine ibex group together there following the winter. In summer, as snow melts and more pastures become available, male Alpine ibex migrate to a higher altitude (Parrini et al. 2003) and therefore have fewer constraints on where to feed and with whom to associate. Accordingly, overall gregariousness and TGS decrease in addition to the mean strength centrality. Similar results have been found in other ungulates, for example in reindeer *Rangifer tarandus*: Peignier et al. (2019) observed low home range overlap and social associations in summer, when resources were distributed homogeneously, but higher home range overlap and social associations in winter, when resources were distributed heterogeneously. Our results hence strengthen the hypothesis that ecological constraints may influence social behaviour in ungulates. On the other side, in horses, strength centrality was found to be lower outside of the mating season; this highlighted a similar increased freedom in the choice of social partners and feeding locations when constraints were reduced, but in this species, the constraints appeared to be social rather than ecological (Stanley et al. 2018).

Finally, we aimed to verify if the overall structure of the social network remained consistent over a period of ten years and whether possible changes could be explained by

demographic characteristics of the population. Due to their experience and knowledge, older individuals are expected to help maintain group cohesion, e.g. leading group movement (McComb et al. 2001). Despite most of the study on this topic being conducted on females in matrilineal societies (McComb et al. 2001, Brent et al. 2015), it has recently been shown that the same may apply also to males (Allen et al. 2020). We therefore expected to find the highest values for global centrality measures in years with a higher number of mature individuals. However, the proportion of males over nine years of age seemed not to affect any of the global measures considered (TGS, gregariousness, density). Also the total number of individuals in the population did not affect either gregariousness or density. The absence of a relationship between global network measures and the demographic structure of the population could be due to the seasonal movement patterns apparent in this species. Unlike animals such as African elephants, the home range of the Alpine ibex in our study area is rather small, around 450–500 ha (Parrini et al. 2003). The seasonal movement of most male Alpine ibex occurs within the same valley, and they show a high site fidelity, possibly because they live in a rather predictable environment (Morrison et al. 2021). This likely makes the role of old, experienced individuals less crucial. Similar conditions may apply to other mountain ungulates with low dispersal rate and high site fidelity (Festa-Bianchet 1986). If this hypothesis is confirmed, it could mean that the social cohesion of male mountain ungulates group is stable and is not dependent on the presence of old and experienced individuals, i.e. it shows resilience to changes in group membership. However, we have to acknowledge that, during the years of the study, the demographic structure of the population was not subject to drastic changes as the environmental conditions remained relatively stable and some old males were alive in the population across all the years of the study, even after harsh winters. This could mean that the presence of a few older individuals may be enough to maintain social cohesion; however, it could also signify that we did not capture enough variation to draw biologically meaningful conclusions. As this aspect could have important conservation implications, further data should be collected to determine the effect of demographic structure on the social structure of mountain ungulates. Particularly, building networks on a finer scale (e.g. daily networks) coupled with more detailed data collection on associations (e.g. using the nearest neighbour method or conducting focal observations), as well as accounting for spatial behaviour (Albery et al. 2021b), could help in better understanding the factors shaping associations and the global network structure.

In conclusion, this study investigated social network dynamics of male of a mountain ungulate over a relatively long time period focusing at the level of group and individual, allowing us to explore ungulate sociality over different scales. As most previous studies on ungulate social networks have focused on female associations, this is one of the few to quantify male social structure in a gregarious ungulate with high levels of sexual segregation. We found that the male

Alpine ibex social network is highly cohesive with all individuals being connected; this has important implications for the management of disease outbreaks. Social structure changed during seasons, suggesting that ecological constraints such as resource availability may play a role in shaping associations in gregarious ungulates. However, we also demonstrated that male Alpine ibex did not associate randomly but showed stable associations, and that the strength of these associations varies across individuals, with age similarity being a factor driving these associations. This, in turn, suggests that ecological constraints are not enough to explain gregariousness and preferential associations in ungulates; these are likely to also be driven by physiological and social needs, ultimately varying according to individual characteristics and species life history. Finally, our results highlight the importance of long-term studies based on individually recognizable subjects (Festa-Bianchet et al. 2017).

Acknowledgements – We thank Brenda McCowan and Sandra Smith Aguilar for constructive reviews of the manuscript and Gabriel Ramos-Fernández for the recommendation and the evaluation process. We deeply thank Bruno Bassano from the Gran Paradiso National Park biodiversity and scientific research service and the Park Rangers of the surveillance service for captures and marking of Alpine ibex and for providing logistic support during field work. We are grateful to all the students and researchers that helped with data collection in the ten years of the study and to Edward Monovich for the artwork. We also thank David Laniado and Cristian Pasquaretta for helpful discussion and suggestion on data analysis. Open access funding enabled and organized by Projekt DEAL.

Funding – The authors conducted this research as part of their academic activity. This research did not receive specific grant from funding agencies.

Conflict of interest – The authors declare they have no conflict of interest relating to the content of this article. Cédric Sueur is a recommender for PCI Network Science.

Ethics statement/Permits

– The capture and marking protocol used in this study has been authorised by the Italian Ministry of Environment (authorisation no. 25114 of 21/09/2004) after the positive review by the Italian National Institute for Environmental Protection and Research (ISPRA) and was developed in order to minimize the effects on the welfare of the animals.

Author contributions

Alice Brambilla: Conceptualization (lead); Data curation (lead); Formal analysis (equal); Investigation (lead); Methodology (equal); Visualization (lead); Writing – original draft (lead); Writing – review and editing (lead). **Achaz von Hardenberg:** Conceptualization (equal); Formal analysis (supporting); Methodology (supporting); Supervision (supporting); Validation (supporting); Writing – original draft (supporting). **Claudia Canedoli:** Data curation (equal); Investigation (equal); Methodology (supporting); Writing – original draft (supporting). **Francesca Brivio:** Data curation (equal); Investigation (equal); Writing – review

and editing (supporting). **Cédric Sueur:** Formal analysis (supporting); Methodology (equal); Supervision (equal); Validation (equal); Writing – original draft (supporting); Writing – review and editing (supporting). **Christina R. Stanley:** Conceptualization (equal); Formal analysis (equal); Methodology (equal); Validation (equal); Writing – original draft (lead); Writing – review and editing (supporting).

Version 3 of the preprint of this manuscript has been peer-reviewed and recommended by Gabriel Ramos-Fernandez (2022) for Peer Community In Network Science.

Data availability availability

The data used to produce this manuscript are accessible on the Dryad Digital Repository: <<https://doi.org/10.5061/dryad.w0vt4b8st>>. The code used for the data analysis contained in this manuscript is accessible at Zenodo: <<https://doi.org/10.5281/zenodo.6499633>>.

Supporting information

The Supporting information associated with this article is available with the online version.

References

- Akaike, H. 1973. Information theory and the maximum likelihood principle. – In: Petrov, B. N. and Csáki, F. (eds), 2nd international symposium on information theory. Akademiai Kiado.
- Alados, C. L. 1986. Spatial structure in groups of Spanish ibex *Capra pyrenaica*. – Biol. Behav. 11: 176–185.
- Albers, P. C. and Vries, H. D. 2001. Elo-rating as a tool in the sequential estimation of dominance strengths. – Anim. Behav. 61: 489–495.
- Albery, G. F. et al. 2021a. Ageing red deer alter their spatial behaviour and become less social. – bioRxiv, <<https://doi.org/10.1101/2021.06.11.448092>>.
- Albery, G. F. et al. 2021b. Multiple spatial behaviours govern social network positions in a wild ungulate. – Ecol. Lett. 24: 676–686.
- Allen, C. R. et al. 2020. Importance of old bulls: leaders and followers in collective movements of all-male groups in African savannah elephants *Loxodonta africana*. – Sci. Rep. 10: 13996.
- Altmann, J. 1974. Observational study of behavior: sampling methods. – Behaviour 49: 227–266.
- Apollonio, M. et al. 2013. Consequences of snowy winters on male mating strategies and reproduction in a mountain ungulate. – Behav. Process. 98: 44–50.
- Appleby, M. C. 1983. Competition in a red deer stag social group: rank, age and relatedness of opponents. – Anim. Behav. 31: 913–918.
- Aureli, F. et al. 2008. Fission–fusion dynamics: new research frameworks. – Curr. Anthropol. 49: 627–654.
- Barton, K. 2009 Mu-MIn: multi-model inference. – R package ver. 1.43.17. <<https://CRAN.R-project.org/package=MuMIn>>.
- Bassano, B. et al. 2003. A method to weigh free-ranging ungulates without handling. – Wildl. Soc. Bull. 31: 1205–1209.

- Bastian, M. et al. 2009. Gephi: an open source software for exploring and manipulating networks. – In: Proc. Int. AAAI Conf. on web and social media, vol. 3, pp. 361–362, <<https://ojs.aaai.org/index.php/ICWSM/article/view/13937>>.
- Bates, D. et al. 2015. Fitting linear mixed-effects models using lme4. – J. Stat. Softw. 67: 1–48.
- Bergeron, P. et al. 2008. Heterogeneity in male horn growth and longevity in a highly sexually dimorphic ungulate. – Oikos 117: 77–82.
- Bergeron, P. et al. 2010. Secondary sexual characters signal fighting ability and determine social rank in Alpine ibex *Capra ibex*. – Behav. Ecol. Sociobiol. 64: 1299–1307.
- Bon, R. et al. 2001. Segregation is not only a matter of sex in Alpine ibex, *Capra ibex ibex*. – Anim. Behav. 62: 495–504.
- Bond, M. L. et al. 2019. Fission–fusion dynamics of a megaherbivore are driven by ecological, anthropogenic, temporal and social factors. – Oecologia 191: 335–347.
- Borgeaud, C. et al. 2017. The influence of demographic variation on social network stability in wild vervet monkeys. – Anim. Behav. 134: 155–165.
- Brambilla, A. and Brivio, F. 2018. Assessing the effects of helicopter disturbance in a mountain ungulate on different time scales. – Mammal. Biol. 90: 30–37.
- Brambilla, A. and Canedoli, C. 2014. How to continue measuring horn growth after capture in Alpine ibex. – J. Mountain Ecol. 9: 35–46.
- Brambilla, A. et al. 2013. Don't spit in the soup: faecal avoidance in foraging wild Alpine ibex, *Capra ibex*. – Anim. Behav. 86: 153–158.
- Brambilla, A. et al. 2015. Direct and indirect causal effects of heterozygosity on fitness-related traits in Alpine ibex. – Proc. R. Soc. B 282: 20141873.
- Brambilla, A. et al. 2018. Heterozygosity–fitness correlation at the major histocompatibility complex despite low variation in Alpine ibex *Capra ibex*. – Evol. Appl. 11: 631–644.
- Brambilla, A. et al. 2020. Distribution, status and recent population dynamics of Alpine ibex *Capra ibex* in Europe. – Mammal Rev. 50: 267–277.
- Breheny, P. and Burchett, W. 2017. Visualization of regression models using visreg. – R J. 9: 56–71.
- Brent, L. J. et al. 2011. Social network analysis in the study of nonhuman primates: a historical perspective. – Am. J. Primatol. 73: 720–730.
- Brent, L. J. et al. 2015. Ecological knowledge, leadership and the evolution of menopause in killer whales. – Curr. Biol. 25: 746–750.
- Brivio, F. et al. 2010. To feed or not to feed? Testing different hypotheses on rut-induced hypophagia in a mountain ungulate. – Ethology 116: 406–415.
- Brivio, F. et al. 2015. Assessing the impact of capture on wild animals: the case study of chemical immobilisation on alpine ibex. – PLoS One 10: e0130957.
- Burnham, K. P. et al. 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations and comparisons. – Behav. Ecol. Sociobiol. 65: 23–35.
- Butts, C. T. 2020. sna: tools for social network analysis. – R package ver. 2.6, <<https://CRAN.R-project.org/package=sna>>.
- Cameron, E. Z. et al. 2009. Social bonds between unrelated females increase reproductive success in feral horses. – Proc. Natl Acad. Sci. 106: 13850–13853.
- Cantor, M. et al. 2012. Disentangling social networks from spatiotemporal dynamics: the temporal structure of a dolphin society. – Anim. Behav. 84: 641–651.
- Carter, K. D. et al. 2013. Social networks, long-term associations and age-related sociability of wild giraffes. – Anim. Behav. 86: 901–910.
- Cassinello, J. and Calabuig, G. 2008. Spatial association in a highly inbred ungulate population: evidence of fine-scale kin recognition. – Ethology 114: 124–132.
- Chiyo, P. I. et al. 2011. Association patterns of African elephants in all-male groups: the role of age and genetic relatedness. – Anim. Behav. 81: 1093–1099.
- Cinar, O. and Viechtbauer, W. 2021. poolr: methods for pooling p-values from (dependent) tests. – R package ver. 1.0-0, <<https://CRAN.R-project.org/package=poolr>>.
- Clutton-Brock, T. 2021. Social evolution in mammals. – Science 373: eabc9699.
- Connor, R. C. and Wells, R. S. 2000. The bottlenose dolphin: social relationships in a fission–fusion society. – In: Mann, J. et al. (eds), Cetacean societies: field studies of dolphins and whales. Univ of Chicago Press, pp. 91–126.
- Conradt, L. and Roper, T. J. 2000. Activity synchrony and social cohesion: a fission–fusion model. – Proc. R. Soc. B 267: 2213–2218.
- Couturier, M. A. J. 1962. Le Bouquetin des Alps: *Capra aegagrus ibex ibex*. L. – Arthaud.
- Couzin, I. D. and Laidre, M. E. 2009. Fission–fusion populations. – Curr. Biol. 19: R633–R635.
- Craft, M. E. 2015. Infectious disease transmission and contact networks in wildlife and livestock. – Phil. Trans. R. Soc. B 370: 20140107.
- Cransac, N. et al. 1998. An example of segregation between age and sex classes only weakly related to habitat use in mouflon sheep *Ovis gmelini*. – J. Zool. 244: 371–378.
- Croft, D. P. et al. 2008. Exploring animal social networks. – Princeton Univ. Press.
- Elo, A. E. 1978. The rating of chess players, past and present. – Arco.
- Festa-Bianchet, M. 1986. Site fidelity and seasonal range use by bighorn rams. – Can. J. Zool. 64: 2126–2132.
- Festa-Bianchet, M. et al. 2017. Successes and challenges of long-term field studies of marked ungulates. – J. Mammal. 98: 612–620.
- Firth, J. A. et al. 2018. Personality shapes pair bonding in a wild bird social system. – Nat. Ecol. Evol. 2: 1696–1699.
- Fischhoff, I. R. et al. 2009. Reproductive status influences group size and persistence of bonds in male plains zebra *Equus burchelli*. – Behav. Ecol. Sociobiol. 63: 1035–1043.
- Flack, J. C. et al. 2006. Policing stabilizes construction of social niches in primates. – Nature 439: 426–429.
- Fogarty, S. et al. 2011. Social personality polymorphism and the spread of invasive species: a model. – Am. Nat. 177: 273–287.
- Franks, D. W. et al. 2010. Sampling animal association networks with the gambit of the group. – Behav. Ecol. Sociobiol. 64: 493–503.
- Godde, S. et al. 2013. Correcting for the impact of gregariousness in social network analyses. – Anim. Behav. 85: 553–558.
- Grueter, C. C. et al. 2020. Multilevel organisation of animal sociality. – Trends Ecol. Evol. 35: 834–847.
- Handcock, M. S. et al. 2008. statnet: software tools for the representation, visualization, analysis and simulation of network data. – J. Stat. Softw. 24: 1548.
- Holekamp, K. E. et al. 1997. Patterns of association among female spotted hyenas *Crocuta crocuta*. – J. Mammal. 78: 55–64.
- Holm, S. 1979. A simple sequentially rejective multiple test procedure. – Scand. J. Stat. 6: 65–70.

- Ilany, A. and Akçay, E. 2016. Social inheritance can explain the structure of animal social networks. – *Nat. Commun.* 7: 12084.
- Jarman, P. 1974. The social organisation of antelope in relation to their ecology. – *Behaviour* 48: 215–267.
- Jewell, P. A. et al. 1974. Island survivors. The ecology of the Soay sheep of St. Kilda. – The Athlone Press, Univ. of London.
- Kerth, G. and König, B. 1999. Fission, fusion and nonrandom associations in female Bechstein's bats *Myotis bechsteinii*. – *Behaviour* 136: 1187–1202.
- Kimura, R. 1998. Mutual grooming and preferred associate relationships in a band of free-ranging horses. – *Appl. Anim. Behav. Sci.* 59: 265–276.
- Krackhardt, D. 1988. Predicting with networks: nonparametric multiple regression analysis of dyadic data. – *Soc. Netw.* 10: 359–381.
- Krause, J. et al. 2002. Living in groups. – Oxford Univ. Press.
- Krause, J. et al. 2009. Animal social networks: an introduction. – *Behav. Ecol. Sociobiol.* 63: 967–973.
- Krause, J. et al. 2010. Personality in the context of social networks. – *Phil. Trans. R. Soc. B* 365: 4099–4106.
- Krause, J. et al. 2015. Animal social networks. – Oxford Univ. Press.
- Le Pendu, Y. et al. 1995. Inter-individual associations and social structure of a mouflon population *Ovis orientalis musimon*. – *Behav. Processes.* 34: 67–80.
- Machanda, Z. P. and Rosati, A. G. 2020. Shifting sociality during primate ageing. – *Phil. Trans. R. Soc. B* 375: 20190620.
- MacIntosh, A. J. et al. 2012. Monkeys in the middle: parasite transmission through the social network of a wild primate. – *PLoS One* 7: e51144.
- Marchand, P. et al. 2017. Sociospatial structure explains marked variation in brucellosis seroprevalence in an Alpine ibex population. – *Sci. Rep.* 7: 1–12.
- McComb, K. et al. 2001. Matriarchs as repositories of social knowledge in African elephants. – *Science* 292: 491–494.
- Morrison, T. A. et al. 2021. Drivers of site fidelity in ungulates. – *J. Anim. Ecol.* 90: 955–966.
- Moscovice, L. R. et al. 2020. How socio-ecological factors influence the differentiation of social relationships: an integrated conceptual framework. – *Biol. Lett.* 16: 20200384.
- Neumann, C. et al. 2011. Assessing dominance hierarchies: validation and advantages of progressive evaluation with Elo-rating. – *Anim. Behav.* 82: 911–921.
- Opsahl, T. 2009. Structure and evolution of weighted networks. – Univ. of London (Queen Mary College), pp. 104–122. <<http://toreopsahl.com/tnet/>>.
- Papageorgiou, D. and Farine, D. R. 2020. Multilevel societies in birds. – *Trends Ecol. Evol.* 36: P15–P17.
- Parrini, F. et al. 2003. Spatial behaviour of adult male Alpine ibex *Capra ibex ibex* in the Gran Paradiso National Park, Italy. – *Acta Theriol.* 48: 411–423.
- Peignier, M. et al. 2019. Space use and social association in a gregarious ungulate: testing the conspecific attraction and resource dispersion hypotheses. – *Ecol. Evol.* 9: 5133–5145.
- Pelletier, F. and Festa-Bianchet, M. 2006. Sexual selection and social rank in bighorn rams. – *Anim. Behav.* 71: 649–655.
- Pinter-Wollman, N. et al. 2014. The dynamics of animal social networks: analytical, conceptual and theoretical advances. – *Behav. Ecol.* 25: 242–255.
- Podgórski, T. et al. 2014. Long-lasting, kin-directed female interactions in a spatially structured wild boar social network. – *PLoS One* 9: e99875.
- Poissant, J. et al. 2013. A quantitative trait locus analysis of personality in wild bighorn sheep. – *Ecol. Evol.* 3: 474–481.
- Ramos, A. et al. 2019. The social network structure of a semi-free roaming European bison herd *Bison bonasus*. – *Behav. Processes.* 158: 97–105.
- Ramos-Fernández, G. 2022. A social network of bucks. – *Peer Community in Network Science*, p. 100004.
- Réale, D. and Festa-Bianchet, M. 2003. Predator-induced natural selection on temperament in bighorn ewes. – *Anim. Behav.* 65: 463–470.
- Réale, D. et al. 2000. Consistency of temperament in bighorn ewes and correlates with behaviour and life history. – *Anim. Behav.* 60: 589–597.
- Réale, D. et al. 2007. Integrating animal temperament within ecology and evolution. – *Biol. Rev.* 82: 291–318.
- Réale, D. et al. 2010. Evolutionary and ecological approaches to the study of personality. – *Phil. Trans. R. Soc. B* 365: 3937–3946.
- Ruckstuhl, K. and Neuhaus, P. 2000. Sexual segregation in ungulates: a new approach. – *Behaviour* 137: 361–377.
- Ruckstuhl, K. and Neuhaus, P. 2001. Behavioral synchrony in ibex groups: effects of age, sex and habitat. – *Behaviour* 138: 1033–1046.
- Schakner, Z. A. et al. 2017. Social associations between California sea lions influence the use of a novel foraging ground. – *R. Soc. Open Sci.* 4: 160820.
- Shizuka, D. and Johnson, A. E. 2020. How demographic processes shape animal social networks. – *Behav. Ecol.* 31: 1–11.
- Signer, C. et al. 2011. Hypometabolism and basking: the strategies of Alpine ibex to endure harsh over-wintering conditions. – *Funct. Ecol.* 25: 537–547.
- Silk, M. J. et al. 2014. The importance of fission–fusion social group dynamics in birds. – *Ibis* 156: 701–715.
- Silk, M. J. et al. 2019. Integrating social behaviour, demography and disease dynamics in network models: applications to disease management in declining wildlife populations. – *Phil. Trans. R. Soc. B* 374: 20180211.
- Siracusa, E. R. et al. 2022. Social ageing: exploring the drivers of late-life changes in social behaviour in mammals. – *Biol. Lett.* 18: 20210643.
- Smith, J. E. et al. 2011. Greetings promote cooperation and reinforce social bonds among spotted hyaenas. – *Anim. Behav.* 81: 401–415.
- Smuts, B. B. et al. 1987. Primate societies. – Univ. of Chicago Press.
- Snijders, L. et al. 2017. Animal social network theory can help wildlife conservation. – *Trends Ecol. Evol.* 32: 567–577.
- Sosa, S. 2016. The influence of gender, age, matriline and hierarchical rank on individual social position, role and interactional patterns in *Macaca sylvanus* at 'La Forêt des singes': a multilevel social network approach. – *Front. Psychol.* 7: 529.
- Sosa, S. et al. 2020. A multilevel statistical toolkit to study animal social networks: the animal network toolkit software (ants) R package. – *Sci. Rep.* 10: 1–8.
- Sosa, S. et al. 2021a. Network measures in animal social network analysis: their strengths, limits, interpretations and uses. – *Methods Ecol. Evol.* 12: 10–21.
- Sosa, S. et al. 2021b. Animal social networks: towards an integrative framework embedding social interactions, space and time. – *Methods Ecol. Evol.* 12: 4–9.
- Stanley, C. R. and Dunbar, R. I. M. 2013. Consistent social structure and optimal clique size revealed by social network analysis of feral goats, *Capra hircus*. – *Anim. Behav.* 85: 771–779.
- Stanley, C. R. et al. 2018. Social stability in semiferal ponies: networks show interannual stability alongside seasonal flexibility. – *Anim. Behav.* 136: 175–184.

- Statnet Development Team 2003–2020. Statnet: software tools for the statistical modeling of network data. – <<http://statnet.org>>.
- Sueur, C. et al. 2011. Collective decision-making and fission–fusion dynamics: a conceptual framework. – *Oikos* 120: 1608–1617.
- Tarlow, E. M. and Blumstein, D. T. 2007. Evaluating methods to quantify anthropogenic stressors on wild animals. – *Appl. Anim. Behav. Sci.* 102: 429–451.
- Teitelbaum, C. S. et al. 2017. Birds choose long-term partners years before breeding. – *Anim. Behav.* 134: 147–154.
- Toïgo, C. et al. 2007. Sex- and age-specific survival of the highly dimorphic Alpine ibex: evidence for a conservative life-history tactic. – *J. Anim. Ecol.* 76: 679–686.
- Turner, J. W. et al. 2018. Ontogenetic change in determinants of social network position in the spotted hyena. – *Behav. Ecol. Sociobiol.* 72: 10.
- van de Waal, E. and Bshary, R. 2011. Social-learning abilities of wild vervet monkeys in a two-step task artificial fruit experiment. – *Anim. Behav.* 81: 433–438.
- Vander Wal, E. et al. 2015. Sex-based differences in the adaptive value of social behavior contrasted against morphology and environment. – *Ecology* 96: 631–641.
- Vander Wal, E. et al. 2016. Dyadic associations and individual sociality in bighorn ewes. – *Behav. Ecol.* 27: 560–566.
- Villaret, J. C. and Bon, R. 1998. Sociality and relationships in Alpine ibex *Capra ibex*. – *Rev. d'Écol.* 53: 153–170.
- Villaret, J. C. and Bon, R. 1995. Social and spatial segregation in Alpine ibex *Capra ibex* in Bargy, French Alps. – *Ethology* 101: 291–300.
- von Hardenberg, A. et al. 2004. Horn growth but not asymmetry heralds the onset of senescence in male Alpine ibex *Capra ibex*. – *J. Zool.* 263: 425–432.
- Wasserman, S. and Faust, K. 1994. Social network analysis: methods and applications, vol. 8. – Cambridge Univ. Press.
- Webber, Q. M. and Vander Wal, E. 2019. Trends and perspectives on the use of animal social network analysis in behavioural ecology: a bibliometric approach. – *Anim. Behav.* 149: 77–87.
- Welch, M. J. et al. 2020. Social experience of captive Livingstone's fruit bats *Pteropus livingstonii*. – *Animals* 10: 1321.
- Wey, T. et al. 2008. Social network analysis of animal behaviour: a promising tool for the study of sociality. – *Anim. Behav.* 75: 333–344.
- Wey, T. W. and Blumstein, D. T. 2010. Social cohesion in yellow-bellied marmots is established through age and kin structuring. – *Anim. Behav.* 79: 1343–1352.
- Whitehead, H. 2008. Analysing animal societies: quantitative methods for vertebrate social analysis. – Univ. of Chicago Press.
- Whitehead, H. 2009. SOCPROG programs: analysing animal social structures. – *Behav. Ecol. Sociobiol.* 63: 765–778.
- Willisch, C. S. and Neuhaus, P. 2009. Alternative mating tactics and their impact on survival in adult male Alpine ibex *Capra ibex ibex*. – *J. Mammal.* 90: 1421–1430.
- Willisch, C. S. and Neuhaus, P. 2010. Social dominance and conflict reduction in rutting male Alpine ibex, *Capra ibex*. – *Behav. Ecol.* 21: 372–380.
- Willisch, C. S. et al. 2012. Male reproductive pattern in a polygynous ungulate with a slow life-history: the role of age, social status and alternative mating tactics. – *Evol. Ecol.* 26: 187–206.
- Wilson, A. D. et al. 2014. Dynamic social networks in guppies *Poecilia reticulata*. – *Behav. Ecol. Sociobiol.* 68: 915–925.
- Wilson, D. E. and Mittermeier, R. A. 2011. Handbook of the mammals of the world. Vol. 2: Hoofed mammals. – Lynx Editions.
- Wittemyer, G. et al. 2005. The socioecology of elephants: analysis of the processes creating multitiered social structures. – *Anim. Behav.* 69: 1357–1371.
- Wittemyer, G. et al. 2009. Where sociality and relatedness diverge: the genetic basis for hierarchical social organization in African elephants. – *Proc. R. Soc. B* 276: 3513–3521.