

Regulation and roles of Ca²⁺ stores in human sperm

Item Type	Article
Authors	Correia, Joao;Michelangeli, Francesco;Publicover, Stephen
Citation	Correia, J., Michelangeli, F., & Publicover, S. (2015). Regulation and roles of Ca ²⁺ stores in human sperm. <i>Reproduction</i> , 150(2), R65-R76. https://doi.org/10.1530/REP-15-0102
DOI	10.1530/REP-15-0102
Publisher	Bioscientifica
Journal	Reproduction
Rights	An error occurred on the license name.
Download date	2026-06-13 19:16:15
Item License	An error occurred getting the license - uri.
Link to Item	http://hdl.handle.net/10034/606590

1

2

3

4

5 Regulation and roles of Ca²⁺ stores in human sperm.

6

7 Joao Correia, Francesco Michelangeli, Stephen Publicover*

8 School of Biosciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

9 * Author for correspondence (s.j.publicover@bham.ac.uk)

10

11 Abstract.

12 $[Ca^{2+}]_i$ signalling is a key regulatory mechanism in sperm function. In mammalian sperm the Ca^{2+} -
13 permeable, plasma membrane ion channel CatSper is central to $[Ca^{2+}]_i$ signalling but there is good
14 evidence that Ca^{2+} stored in intracellular organelles is also functionally important. Here we briefly
15 review current understanding of the diversity of Ca^{2+} stores and the mechanisms for the regulation
16 of their activity. We then consider the evidence for the involvement of these stores in $[Ca^{2+}]_i$
17 signalling in mammalian (primarily human) sperm, the agonists that may activate these stores and
18 their role in control of sperm function. Finally we consider the evidence that membrane Ca^{2+}
19 channels and stored Ca^{2+} may play discrete roles in the regulation of sperm activities and propose a
20 mechanism by which these different components of the sperm Ca^{2+} -signalling apparatus may
21 interact to generate complex and spatially diverse $[Ca^{2+}]_i$ signals.

22

23 1. Ca²⁺ signalling in sperm

24 Cellular activity is constantly regulated by environmental cues and signals from other cells. Long
25 term regulation of cell function is normally achieved by control of gene expression, changing the
26 complement and levels of proteins in the cell, but rapid or short-term changes are achieved by 'post-
27 translational' protein modification, such as phosphorylation, sumoylation and nitrosylation, which
28 alter the function/activity of proteins already present. Ca²⁺-signalling is a key regulator of such post-
29 translational modifications, with changes in cytoplasmic Ca²⁺ concentration ([Ca²⁺]_i) controlling the
30 activities of key enzymes and proteins. Large changes in [Ca²⁺]_i can be achieved 'instantaneously' by
31 flux of Ca²⁺ into the cytoplasm from the extracellular fluid or from storage organelles (primarily the
32 endoplasmic reticulum) within the cell (fig 1a). The rapidity with which [Ca²⁺]_i-signals can be
33 generated is crucial for 'instantaneous' cellular responses such as activation of muscle contraction
34 and secretion of neurotransmitter, that are achieved by rapid post-translational-modification of
35 protein function.

36 The highly-condensed nucleus of sperm is transcriptionally silent (Miller and Ostermeier 2006,
37 Miller, et al. 2005) and translational activity is also negligible [though evidence has been presented
38 for translation occurring at mitochondrial ribosomes (Chandrashekan, et al. 2014a,
39 Chandrashekan, et al. 2014b, Gur and Breitbart 2007, Zhao, et al. 2009)]. Regulation of sperm
40 function is therefore dependent primarily on post-translational processes. [Ca²⁺]_i signalling is pivotal
41 to this regulation and in mammalian sperm it plays a central role in controlling the cell's behaviour
42 (motility type and potentially chemotaxis), the induction of acrosome reaction and the process of
43 capacitation (Darszon, et al. 2011, Darszon, et al. 2007, Publicover, et al. 2007). The importance for
44 sperm function of membrane Ca²⁺-channels and Ca²⁺-influx is well-established (Darszon, et al. 2011)
45 but there is also good evidence for the existence and functional importance of intracellular Ca²⁺-
46 storage organelles in sperm (Darszon, et al. 2007, Publicover, et al. 2007). Previously we reviewed
47 the identities and functions of Ca²⁺ stores in sperm, focussing on the evidence for the existence of

48 such stores, their components (pumps and channels) and their possible roles in the regulation of
49 function in the mature sperm cell (Costello, et al. 2009). Since then considerable progress has been
50 made in understanding the central role of Ca^{2+} signalling in the regulation of mammalian and non-
51 mammalian sperm function and the mechanisms by which sperm $[\text{Ca}^{2+}]_i$ signals are generated. In
52 particular successful application of whole cell patch clamp technique, in human as well as mouse
53 sperm, has revealed the central importance of Ca^{2+} influx through CatSper, a sperm specific, Ca^{2+} -
54 permeable channel in the membrane of the flagellar principal piece. Male mice null for CatSper are
55 infertile (Ren, et al. 2001) and their sperm show defective motility (Carlson, et al. 2003). Here we
56 review recent progress in understanding the diversity of mechanisms for the regulation of Ca^{2+} store
57 activity and the evidence for their involvement in controlling sperm function.

58 2. Ca^{2+} stores and their regulation

59 The importance of Ca^{2+} stores in generating complex Ca^{2+} signals in somatic cells has long been
60 recognized. Until relatively recently the endoplasmic reticulum Ca^{2+} store has been the major focus
61 for research as this was the first organelle to show controllable mobilization of Ca^{2+} through second
62 messengers acting upon intracellular Ca^{2+} channels, as well as being able to be refilled via Ca^{2+}
63 pumps. Additionally, these Ca^{2+} signals could also be re-modelled through the regulation of these
64 Ca^{2+} transporters to generate complex spatial and temporal Ca^{2+} transients (Berridge, et al. 2003). It
65 has now become clear that many other organelles such as mitochondria, endosomes, lysosomes and
66 Golgi complexes also contribute to the generation and propagation of these complex Ca^{2+} signals
67 within cells (Michelangeli, et al. 2005). Furthermore, novel Ca^{2+} transporters have also been
68 identified within these other organelles and several have recently been identified in sperm (Costello,
69 et al. 2009).

70 (i) Intracellular Ca^{2+} Channels

71 The major intracellular Ca^{2+} channels that have been identified and appear to be almost ubiquitously
72 distributed within mammalian cells, especially on the endoplasmic reticulum, include the inositol-
73 1,4,5-trisphosphate-(IP_3)-sensitive Ca^{2+} channel (or IP_3 receptor; IP_3R) and the ryanodine receptor
74 (RyR) (Michelangeli, et al. 2005) (fig 1a). The IP_3 receptor, as the name implies, is activated by the
75 second messenger IP_3 that is generated through the hydrolysis of phosphatidylinositol-4,5-
76 bisphosphate. This channel has a specific IP_3 binding site that is located towards the N-terminus of
77 the protein (Seo, et al. 2012) and also has a requirement for Ca^{2+} which acts as a co-agonist in order
78 for the channel to open (Bezprozvanny, et al. 1991). The activation of RyR is likely to be through a
79 mechanism involving Ca^{2+} induced Ca^{2+} release (CICR) and by the action of the putative second
80 messenger cyclic-adenosine diphospho-ribose (cADPR) (Ogunbayo, et al. 2011) (fig 1a). cADPR is
81 made from nicotinamide-adenine-dinucleotide (NAD) by the action of an ADP-ribosyl cyclase enzyme
82 such as CD38 (Cosker, et al. 2010), although other as yet unidentified enzymes may also be involved
83 in catalysing this reaction (Guse 2014). It is as yet unclear whether, unlike the IP_3R , cADPR binds
84 directly to RyR or whether it binds to accessory proteins such as calmodulin or FK506-binding
85 protein, that then interact with the RyR (Guse 2014).

86 Another metabolite of NAD which is believed to have Ca^{2+} mobilizing ability is nicotinic acid adenine
87 dinucleotide phosphate (NAADP) (Genazzani, et al. 1997). NAADP is made from NADP through the
88 action of either CD38 acting as a base-exchanger, swapping the nicotinamide group for nicotinic acid
89 or via an unidentified NADP-deaminase (Guse 2014). NAADP is believed specifically to mobilize Ca^{2+}
90 from acidic stores such as lysosomes (Churchill, et al. 2002, Menteyne, et al. 2006), which can then
91 induce CICR at RyRs and IP_3Rs in mammalian cells (Cancela et 1999) (fig 1a). Results initially
92 presented by Calcraft and colleagues (Calcraft, et al. 2009), indicated that NAADP specifically
93 activates Ca^{2+} specific 'two-pore' channels (TPC) within the acidic organelles, these channels being
94 first described in plants (Peiter, et al. 2005). However, in kinetic studies there is a prominent lag
95 between addition of NAADP and Ca^{2+} mobilization (Genazzani, et al. 1997). Combined with the
96 observation that photo-affinity labelling with azido-NAADP (Lin-Moshier, et al. 2012) showed

97 labelling of only low molecular weight proteins, not consistent with TPCs, this suggests that NAADP
98 might function by binding to accessory proteins rather than directly to the channel. More recently
99 two studies (Cang, et al. 2013, Wang, et al. 2012) have raised considerable controversy as whether
100 the NAADP-sensitive Ca^{2+} channel is a TPC. Both studies suggest that TPCs are in fact Na^{2+} -specific
101 channels with very low Ca^{2+} selectivity that are activated by phosphoinositide lipids and modulated
102 by mTOR, but not by NAADP. More work is currently being undertaken to clarify this and a number
103 of possible theories as to what role TPCs play in NAADP-induced Ca^{2+} mobilization are being explored
104 (see Morgan and Galione (2014)).

105 Numerous kinases have been shown to modulate the activity of both the IP_3Rs and RyRs , these
106 include several ubiquitous ser/thr kinases such as PKA, PKG and CaMKII (Camors and Valdivia 2014,
107 Yule, et al. 2010). Indeed some of these kinases such as PKA appear to have both stimulatory and
108 inhibitory effects on the IP_3R dependent upon isoform subtype and the presence of multiple kinase-
109 dependent phosphorylation sites on the same receptor (Dyer, et al. 2003). Less ubiquitous ser / thr
110 kinases such as Akt and polo kinases as well as tyrosine kinases such as fyn kinase have also been
111 shown to affect these channels (Camors and Valdivia 2014, Yule, et al. 2010).

112 Both the RyRs and the IP_3Rs are modulated by changes in their oxidation states caused by reactive
113 oxygen species (ROS) and reactive nitrogen species (RNS) and this occurs mainly through
114 modification of specific cysteine (cys) amino acid residues. Oxidation of these cys residues in RyRs
115 occurs both by s-glutathionylation as well as s-nitrosylation by the second messenger nitric oxide
116 (NO) (Csordas and Hajnoczky 2009) and promotes the activity of the channel by enhancing RyR
117 subunit interactions and also by reducing the efficacy of inhibitory modulators (Hamilton and Reid
118 2000). In IP_3Rs the effects of oxidative stress are complex: low levels of cys oxidation caused by low
119 concentrations of thimerosal (a cys-modifying mercuric compound) and naturally generated ROS
120 cause sensitization of this channel, while higher concentrations of thimerosal inhibit channel activity

121 (Missiaen, et al. 1991, Sayers, et al. 1993). Currently, however, there is little evidence that NO can
122 affect the activity of the IP₃Rs.

123 (ii) Intracellular Ca²⁺ pumps

124 The major transporter involved in refilling Ca²⁺ stores within the endoplasmic reticulum is the
125 sarcoplasmic/endoplasmic reticulum Ca²⁺ ATPase (SERCA) (fig 1a) and these pumps occur abundantly
126 in all somatic cells. Their role is to pump Ca²⁺ back into the storage organelles to help terminate Ca²⁺
127 signals (Michelangeli and East 2011, Michelangeli, et al. 2005). There are 3 isoforms of this
128 Ca²⁺ATPase, each encoded by a different gene and each isoform can exist in a variety of spliced
129 variants that differ in size and regulatory properties (Michelangeli and East 2011). SERCA1 is mainly
130 confined to skeletal muscle, while SERCA2 is widely distributed in most other tissues and organs and
131 type 3 has a limited expression. Another related Ca²⁺ATPase that is also found ubiquitously within
132 somatic cells is the secretory pathway Ca²⁺ ATPase (SPCA) which is localized to the Golgi apparatus
133 (Wootton, et al. 2004). SPCA exists in 2 isoforms with the expression of type 1 being far more
134 widespread than type 2, which appears to be mainly located within glandular tissues (Vanoevelen, et
135 al. 2005). Recently there has been evidence to suggest the SPCA2 can interact with and regulate the
136 plasma membrane located Orai Ca²⁺ channels that are implicated in store-operated Ca²⁺ entry (Feng,
137 et al. 2010), which may indicate a dual function for this Ca²⁺ ATPase in cells that express it.

138 There is currently some debate as to which type of intracellular Ca²⁺ ATPase is expressed in mature
139 sperm. We have highlighted that SPCA1 is present in human sperm, where it appears to be mainly
140 localized to the neck region of the cell where the redundant nuclear envelope (RNE) and calreticulin-
141 containing vesicles are situated (Harper, et al. 2005). This study also found no evidence for
142 expression of SERCA in human sperm as no cross-reactivity was observed with a pan-isoform SERCA
143 antibody and no effects on [Ca²⁺]_i were observed with specific but saturating concentrations of the
144 SERCA-inhibitor thapsigargin. However, a more recent study (Lawson, et al. 2007) detected SERCA2,
145 mainly localized to the acrosome and mid-piece, using a SERCA2-specific antibody.

146 Unlike the intracellular Ca^{2+} channels, there is no strong evidence to suggest that either SERCA or
147 SPCA can be directly phosphorylated and regulated by protein kinases, although some Ca^{2+} ATPase
148 modulatory proteins like phospholamban (that is found almost exclusively in heart) are regulated
149 through phosphorylation by PKA, PKG and CamKII (Colyer 1998). There is considerable evidence
150 indicating that oxidative stress can modulate SERCA activity (although no studies have yet been
151 undertaken on SPCA). Again a number of critical cys residues such as cys674 can be s-
152 glutathionylated to cause an increase in SERCA pump activity (Adachi, et al. 2004). Modifications of
153 other cys residues on the Ca^{2+} ATPase, however, can have inhibitory effects (Csordas and Hajnoczky
154 2009, Sayers, et al. 1993, Sharov, et al. 2006).

155

156 3. Ca^{2+} stores, mechanisms for store mobilisation and store-operated Ca^{2+} channels in sperm

157 During the later stages of their development spermatozoa shed much of their cytoplasm including
158 intracellular organelles. Thus mammalian sperm contain no organised endoplasmic reticulum.
159 However, studies on the expression of Ca^{2+} store components and on the generation $[\text{Ca}^{2+}]_i$ signals
160 suggest that the remaining intracellular organelles function as Ca^{2+} -stores and play a significant role
161 in the regulation of cellular function (Costello, et al. 2009). In particular, the acrosomal vesicle at the
162 apex of the head and the collection of vesicular membranous structures that occur at the sperm
163 neck and anterior midpiece (including the cytoplasmic droplet of human sperm) appear to be
164 functionally important Ca^{2+} -stores (fig 1b; shown in green). At both these locations IP_3Rs have been
165 detected in human and in bovine sperm by immuno-staining (Dragileva, et al. 1999, Ho and Suarez
166 2001, 2003, Kuroda, et al. 1999, Naaby-Hansen, et al. 2001). Ryanodine receptors (RyRs) have also
167 been detected in human and rodent sperm (Lefievre, et al. 2007, Trevino, et al. 1998). Staining of
168 human sperm with anti-RyR1, anti-RyR2, pan-RyR and BODIPY-FLX ryanodine is localised primarily to
169 the neck region, though some acrosomal staining was also observed (Harper, et al. 2004, Lefievre, et
170 al. 2007, Park, et al. 2011). In contrast, other authors (Ho and Suarez 2001) have reported no
171 staining of bovine sperm with BODIPY-FLX ryanodine (see (Costello, et al. 2009) for further

172 discussion). Thus mobilisation of stored Ca^{2+} in mammalian sperm may occur in response to
173 generation of IP_3 by activity of phospholipase C and by Ca^{2+} -induced Ca^{2+} release (CICR) at IP_3 Rs or
174 RyRs. These processes can be sensitised by effects such as oxidative stress and S-nitrosylation (see
175 section 2). For instance, exposure of human sperm to $\text{NO}\cdot$ at levels equivalent to those produced by
176 explants of reproductive tract lining mobilises store Ca^{2+} and modifies flagellar activity (Lefievre, et
177 al. 2007, Machado-Oliveira, et al. 2008).

178 In addition to generation of IP_3 in sperm, there is evidence that other Ca^{2+} mobilising messengers
179 (NAADP and cADPR) are synthesised in sperm and/or produced in response to stimulation. Sea
180 urchin sperm contain significant levels of both cADPR and NAADP, which may contribute to oocyte
181 activation (Billington, et al. 2002, Chini, et al. 1997). Human sperm have been shown to contain
182 cADPR at micromolar concentrations but NAADP was not detected (Billington, et al. 2006).
183 Interestingly, this study also demonstrated synthesis of cADPR by human sperm but the ecto-
184 enzyme CD38 (an enzyme present on mammalian cells that synthesises both cADPR and NAADP; see
185 section 2) could not be detected by western blotting. In contrast, Park and colleagues (Park, et al.
186 2011), reported detection of CD38 in human sperm after co-incubation with prostasomes (prostate-
187 derived membrane vesicles; see below). Furthermore, the presence of a novel NAADP synthase,
188 which lacks the cyclase activity of CD38, has been described both in sea urchin (Vasudevan, et al.
189 2008) and human sperm (Sánchez-Tusie et al, 2014). In sea urchin sperm this enzyme is strongly
190 Ca^{2+} -regulated and most active at acid pH whereas the human enzyme shows only weak Ca^{2+} -
191 regulation and activity is maximal at pH 7 to 8 (Sanchez-Tusie, et al. 2014, Vasudevan, et al. 2008).

192 Recent findings have supported the idea that NAADP is functional in human sperm. Sanchez-Tusie et
193 al. (2014) investigated the effects of cell-permeant (AM-ester) derivatives of NAADP and cADPR. No
194 effects were observed with cADPR, consistent with previous pharmacological investigation by
195 Billington et al. (2006), but NAADP caused elevation of $[\text{Ca}^{2+}]_i$ both in cells incubated under standard
196 conditions and also when $[\text{Ca}^{2+}]_o$ was buffered to 100 nM, conditions under which Ca^{2+} influx is

197 negligible and $[Ca^{2+}]_i$ signalling depends solely on mobilisation of stored Ca^{2+} . Staining of NAADP
198 receptors using the fluorescent NAADP receptor ligand Ned-19 and identification of acidic organelles
199 using lysotracker highlighted both an anterior store (potentially the acrosome) and a store at the
200 sperm neck (fig 1b). Consistent with these findings, Arndt et al. (2014), studying acrosome reaction
201 (see below), provided evidence for involvement in this process of NAADP and two-pore channels
202 (TPCs), which have been proposed to be the NAADP receptor/ Ca^{2+} channel of acidic Ca^{2+} storage
203 organelles (Calcraft, et al. 2009) (Fig 1a; see section 2).

204 (Park et al. (2011) investigated the incorporation into human sperm of proteins from prostasomes
205 (prostate-derived vesicles which are normally added to sperm during ejaculation) and their effects
206 on $[Ca^{2+}]_i$ signalling. They concluded that CatSper channel proteins were present in the differentiated
207 sperm, but other Ca^{2+} signalling 'tools' including RyRs and CD38 were added to the freshly-ejaculated
208 sperm upon mixing with prostasomes, by fusion with the membrane of the midpiece. They also
209 examined the effects of stimulation with progesterone on $[Ca^{2+}]_i$ and motility of sperm exposed to
210 prostasomes and sperm that had been rapidly removed from semen to minimise mixing with
211 prostasomes. Their data suggest that the generation of sustained $[Ca^{2+}]_i$ signals (such as the second
212 component of the biphasic progesterone-induced $[Ca^{2+}]_i$ signal) and consequent effects on motility
213 may depend, at least partly, upon generation of cADPR by prostasome-derived enzymes.

214 Interestingly, CD38-null mice proved to be fertile, but analysis showed that 20% of normal ADPR
215 cyclase activity remained in prostasomes from these animals, indicating the presence of a non-CD38
216 ADPR-cyclase, potentially that described by Sánchez-Tusie et al (2014). Thus both NAADP and cADPR
217 are potentially synthesised by sperm and involved in regulation of sperm Ca^{2+} store activity but their
218 roles are not yet clear.

219 In somatic cells mobilisation of stored Ca^{2+} induces secondary Ca^{2+} influx through channels at the cell
220 membrane (store-operated channels; SOCs) by the process of capacitative Ca^{2+} entry (CCE) (fig 1a).
221 CCE both prolongs Ca^{2+} signals that are induced by store mobilisation and provides Ca^{2+} for re-

222 charging of the storage organelles. Recently great progress has been made in elucidating the key
223 players and mechanisms in this process. STIM (stromal interaction molecule) has been identified as
224 the sensor molecule present in the membrane of the Ca^{2+} store. The intraluminal part of STIM
225 includes a Ca^{2+} -binding EF hand that detects depletion of stored Ca^{2+} . STIM then redistributes,
226 moving to a position adjacent to the plasma membrane where it activates channel proteins (Orai
227 and possibly members of the TRPC [transient receptor potential canonical] family; Cahalan 2009).
228 $[\text{Ca}^{2+}]_i$ signals in human and other mammalian sperm induced by agonists and by treatments
229 designed to mobilise stored- Ca^{2+} show characteristics consistent with the occurrence of CCE
230 (Blackmore 1993, Dragileva, et al. 1999, Lefievre, et al. 2012, O'Toole, et al. 2000, Park, et al. 2011).
231 STIM1, Orai and TRPC proteins have been detected in human sperm (Castellano, et al. 2003,
232 Darszon, et al. 2012, Lefievre, et al. 2012), STIM1 being localised primarily to the neck
233 region/midpiece and the acrosome where Ca^{2+} stores are present (Lefievre, et al. 2012). To date the
234 application of whole-cell patch clamp has not provided evidence for the occurrence of CCE in human
235 sperm (Lefievre, et al. 2012) so these findings must be interpreted cautiously, but $[\text{Ca}^{2+}]_i$ signals
236 generated by mobilisation of Ca^{2+} stores in sperm may be amplified by activation of CCE. Induction of
237 CCE in somatic cells can have a latency of 10s of seconds due to the need for STIM to migrate to the
238 peripheral portions of the endoplasmic reticulum where it can interact with SOC proteins (Luik, et al.
239 2006, Wu, et al. 2006), but in sperm the storage organelles are close to the plasma membrane and
240 STIM proteins are localised here, such that CCE could be near 'instantaneous'. Pre-treatment of
241 human sperm with low concentrations of 2-aminoethoxydiphenyl borate (2-APB), which potentiates
242 CCE by promoting the interaction of STIM with SOCs (Navarro-Borelly, et al. 2008, Wang, et al. 2009,
243 Yamashita, et al. 2011) significantly enhances the amplitude of the progesterone-induced Ca^{2+}
244 transient at the sperm neck (where secondary release of stored Ca^{2+} may occur; fig 1b; see section 6)
245 but does not affect the response in the flagellum, where progesterone activates CatSper channels
246 (fig 1b), or the kinetics of the signal at either location (Lefievre, et al. 2012). Conversely, when sperm
247 were pre-treated with a cell-penetrating peptide that mimics part of the key SOAR region of STIM1

248 (potentially preventing auto-inhibitory folding of STIM upon store-refilling) there was a marked
249 prolongation of the progesterone-induced $[Ca^{2+}]_i$ transient in a subset of cells (Morris, et al. 2015).

250

251

252 4. Mobilisation of sperm Ca^{2+} stores by agonists

253 In the majority of somatic cells mobilisation of stored Ca^{2+} is induced by agonist-induced synthesis of
254 Ca^{2+} mobilising intracellular messengers. Thus agonist-induced synthesis of inositol trisphosphate,
255 cADPR and NAADP can lead to rapid release of stored Ca^{2+} and generation of local, global and
256 complex spatio-temporal signals (fig 1a). Is there evidence that such processes occur and are
257 functionally significant in responses to agonist stimulation of sperm?

258 The best-characterised agonist-induced $[Ca^{2+}]_i$ signals in sperm are responses to solubilised zona
259 pellucida/zona proteins in mouse cells and progesterone in human. Application of patch clamp has
260 clearly shown that the primary action of progesterone in human sperm is to activate CatSper
261 channels, leading to Ca^{2+} -influx (Lishko, et al. 2011, Strunker, et al. 2011). Strunker and colleagues
262 (Strunker, et al. 2011) investigated the $[Ca^{2+}]_o$ dependence of progesterone-induced $[Ca^{2+}]_i$ signals in
263 rapid-mixing experiments on human sperm and reported that buffering of $[Ca^{2+}]_o$ to ≤ 100 nM
264 abolished the response (though see Espino et al. (2009)), suggesting that any mobilisation of stored
265 Ca^{2+} is a secondary response. Synthesis of IP_3 is reported to occur downstream of progesterone-
266 induced Ca^{2+} influx (Thomas and Meizel 1989), an important observation that should be pursued.
267 Stimulation of mouse sperm with zona proteins induces acrosome reaction, which requires elevation
268 of $[Ca^{2+}]_i$ in the sperm head (Florman, et al. 2008) and is dependent on mobilisation of Ca^{2+} from the
269 acrosomal store ((De Blas, et al. 2002); see below). The nature of the Ca^{2+} influx following
270 stimulation is not clear and several channels may be involved (Cohen, et al. 2014, Florman, et al.
271 2008, Xia and Ren 2009), but Ca^{2+} signals are sensitive to inhibition of G-protein signalling (using
272 pertussis toxin) and inhibition of PLC (Florman, et al. 2008, Ren and Xia 2010). Furthermore, in sperm
273 from mice null for PLC $\delta 4$ (in which males fertility is 'severely impaired') the $[Ca^{2+}]_i$ response is

274 reduced and zona-induced AR does not occur (Fukami, et al. 2001, Fukami, et al. 2003). Thus
275 conventional IP₃-induced mobilisation of stored Ca²⁺ is apparently central to this essential aspect of
276 mammalian sperm physiology.

277 Evidence for the existence of other store-mobilising agonists is largely preliminary, but there are a
278 number of candidates, of which the best-studied is vitamin D (Blomberg Jensen 2014). Human sperm
279 have been shown to express vitamin D receptor (VDR) (Aquila, et al. 2009, Blomberg Jensen, et al.
280 2011, Blomberg Jensen, et al. 2010), the enzymes CYP2R1 and CYP27B (which produce the active
281 compound (1,25(OH)₂D₃) cholcalciferol) and the inactivating enzyme CYP24A1 (Blomberg Jensen, et
282 al. 2011, Blomberg Jensen, et al. 2010). All are expressed in the neck region of the sperm and
283 staining of cells for VDR and CYP24A1 shows a strong association. In sub-fertile patients the
284 proportion of cells expressing CYP24A1 varies greatly and is significantly correlated with semen
285 quality (sperm count, concentration, morphology and motility) (Blomberg Jensen, et al. 2011,
286 Blomberg Jensen, et al. 2012). Stimulation of human sperm with 1,25(OH)₂D₃ (100 pM-1 uM)
287 induced a [Ca²⁺]_i response, including a transient and plateau, that was blocked by pre-treatment with
288 the non-genomic VDR antagonist 1b,25(OH)₂D₃ but was insensitive to blockade of the nuclear VDR
289 antagonist ZK159222 (Blomberg Jensen, et al. 2011). This effect was greatly reduced by pre-
290 treatment with the phospholipase C inhibitor U73122 (2 μM) but was inhibited by incubation in
291 EGTA-buffered medium for up to 20 min prior to stimulation. Both motility and AR were significantly
292 increased upon stimulation with 1,25(OH)₂D₃ (Blomberg Jensen, et al. 2011).

293 Kisspeptin, a peptide agonist of the G-protein coupled receptor GPR54/KISS1R has also been shown
294 to cause sustained, dose-dependent elevation of [Ca²⁺]_i in human and in mouse sperm (Hsu, et al.
295 2014, Pinto, et al. 2012). In neurons binding of kisspeptin to its receptor activates PLC and results in
296 generation of IP₃ and diacylglycerol, leading to mobilisation of stored Ca²⁺ and also depolarisation
297 (Beltramo, et al. 2014, Liu, et al. 2008, Pielecka-Fortuna, et al. 2008). In human sperm the effect of
298 kisspeptin on [Ca²⁺]_i did not occlude the response to stimulation with the CatSper agonist

299 progesterone and was not reduced when applied in the presence of progesterone (Pinto, et al.
300 2012). Both KISS1R and kisspeptin itself were detected in the head of human sperm, suggesting that
301 an autocrine action of the peptide may occur. Motility parameters of kisspeptin-treated cells were
302 significantly altered, including an increase in lateral movement of the head and a decrease in
303 linearity of the sperm path, characteristics of hyperactivated sperm (Pinto, et al. 2012). Ghrelin,
304 another peptide hormone which also acts through mobilisation of stored Ca^{2+} (Camina, et al. 2003),
305 has also been detected in human sperm (Moretti, et al. 2014). Micromolar concentrations of ghrelin
306 have been shown to increase $[\text{Ca}^{2+}]_i$ and motility in rat sperm (Lukaszyk, et al. 2012) but expression
307 of ghrelin receptors or effect of ghrelin on human sperm $[\text{Ca}^{2+}]_i$ have not been investigated.

308 5. Functional significance of Ca^{2+} -stores

309 The acrosome

310 Acrosome reaction (AR) is the fusion between the outer acrosomal membrane and the overlying
311 plasma membrane. Fusion occurs at multiple points, resulting in vesiculation and loss of the fused
312 outer acrosomal membrane/plasmalemma so that the acrosomal content is released and the inner
313 acrosomal membrane becomes the new cell surface. Membrane fusion proteins from the SNARE
314 family are present in the acrosomal region and may be integrated into microdomains that facilitate
315 Ca^{2+} -regulated membrane fusion in a manner that has been compared with events at presynaptic
316 terminals (De Blas, et al. 2005, Mayorga, et al. 2007, Zitanski, et al. 2010). Zona pellucida proteins
317 interact with sperm surface receptors to activate a signalling cascade leading to AR (Florman, et al.
318 2008) and release of acrosomal content at the surface of the zona may, in combination with
319 hyperactivated motility, facilitate zona penetration. However, observation of mouse IVF using sperm
320 with GFP-labelled acrosomes showed that, in addition to cells that undergo AR at the surface of the
321 zona, sperm which arrive having already lost their acrosome (probably within the cumulus) may go
322 on to penetrate the zona and fertilise (Jin, et al. 2011). Physiological inducers of AR that have been
323 studied (primarily mouse ZP3 and progesterone) induce Ca^{2+} influx across the plasma membrane and
324 a sustained rise in $[\text{Ca}^{2+}]_i$. O'Toole, et al (2000) provided pharmacological evidence that ZP3-induced

325 AR in mouse sperm involved activation of store operated Ca^{2+} influx downstream of Ca^{2+} store
326 mobilisation and De Blas, et al (2002) showed that in streptolysin-permeabilised human sperm,
327 mobilisation of the acrosomal Ca^{2+} store was a requirement for acrosome reaction even when it was
328 'directly' induced by introduction of Rab3A into the cytoplasm. Further studies using this
329 permeabilised sperm model have provided information about the mechanisms by which fusion of
330 the plasma and outer acrosomal membranes is regulated, resulting in a detailed model in which
331 mobilisation of the acrosomal store is a central and necessary event (Ruete, et al. 2014). Stimulation
332 of PLC, leading to generation of IP_3 and activation of IP_3Rs in the outer acrosomal membrane may be
333 key to this process (Fukami, et al. 2001, Fukami, et al. 2003), but there is also evidence that the
334 acrosomal membrane contains the NAADP-sensitive, Ca^{2+} -permeable two-pore channels (Calcraft, et
335 al. 2009) and that NAADP mobilises acrosomal Ca^{2+} in mouse sperm (Arndt, et al. 2014).
336 Interpretation of this finding is complex since the regulation and Ca^{2+} -permeability of TPCs have
337 recently been questioned (Cang, et al. 2013, Wang, et al. 2012) (see section 2).

338

339 The redundant nuclear envelope and calreticulin-containing vesicles

340 A second area where Ca^{2+} storage organelles have been identified in mammalian sperm is at the
341 sperm neck and midpiece (fig 1b). Mitochondria have mechanisms for accumulation and release of
342 Ca^{2+} (Drago, et al. 2011, Pizzo, et al. 2012) and therefore may contribute to Ca^{2+} buffering and
343 signalling in this part of the sperm. Inhibition of mitochondrial function in sea urchin sperm, using
344 respiratory inhibitors or uncouplers, causes a rise in $[\text{Ca}^{2+}]_i$ and leads to activation of Ca^{2+} influx that
345 has characteristics consistent with store-operated channels (Ardon, et al. 2009). Treatment with
346 mitochondrial uncouplers (2,4 dinitrophenol [2,4 DNP], carbonyl cyanide-4-
347 (trifluoromethoxy)phenylhydrazone [FCCP]) also increases $[\text{Ca}^{2+}]_i$ in human sperm (Morris and
348 Publicover, unpublished). Mitochondria may thus contribute to shaping of Ca^{2+} signals in sperm.
349 However, the primary stimulus-regulated Ca^{2+} storage in this part of the sperm is in the redundant
350 nuclear envelope and/or a second, apparently separate group of calreticulin-containing vesicular

351 structures, both of which are sited at the sperm neck region and cytoplasmic droplet (Ho and Suarez
352 2001, 2003, Naaby-Hansen, et al. 2001). Mobilisation of Ca^{2+} stored in these compartments regulates
353 flagellar activity and treatment of mouse sperm with thimerosal stimulates hyperactivated motility
354 by activating Ca^{2+} release from these organelles (Ho and Suarez 2001, Marquez, et al. 2007). This
355 effect occurs in the absence of extracellular Ca^{2+} and can be induced in sperm that are null for
356 CatSper (Marquez, et al. 2007). In mouse sperm the direction of the major, high-amplitude flagellar
357 bend of hyperactivated sperm can be clearly characterised by reference to the hooked acrosomal
358 cap (pro-hook or anti-hook). Sperm that became hyperactivated during capacitation *in vitro* (due to
359 activation of CatSper) show pro-hook bends whereas those activated by store mobilisation (using
360 thimerosal) show anti-hook bends (Chang and Suarez 2011). When sperm were observed interacting
361 with the lining of isolated mouse oviducts, most hyperactivated cells showed anti-hook bending of
362 the type which is elicited by store mobilisation (Chang and Suarez 2012).

363 In human sperm a similar effect of store mobilisation is seen. Thimerosal greatly increases the
364 proportion of cells showing hyperactivated motility and 4-aminopyridine, which both alkalinises the
365 cytoplasm (and will thus activate CatSper) and mobilises stored Ca^{2+} , is similarly potent (Alasmari, et
366 al. 2013a, Alasmari, et al. 2013b). In contrast, manipulations that should activate CatSper (elevation
367 of pH_i , stimulation with progesterone or prostaglandin E_1) elevate $[\text{Ca}^{2+}]_i$ but have only minor
368 stimulatory effects on the proportion of hyperactivated cells. Instead, these manipulations
369 significantly increase penetration into viscous media (Alasmari, et al. 2013a, Alasmari, et al. 2013b,
370 Luo, et al. 2014).

371

372 6. Model for interaction of CatSper channels and Ca^{2+} -stores

373 Patch clamp recordings have provided no evidence that conventional voltage-operated Ca^{2+} channels
374 contribute to Ca^{2+} influx in mature mammalian sperm. In mouse sperm null for CatSper1 and the K^+
375 channel Slo3 only a small leak current was recorded even at high intracellular pH and strong

376 depolarisation (Zeng, et al. 2013). CatSper channels in mouse and human sperm are pH- and
377 (weakly) voltage-sensitive but in human sperm the channel is also ligand-sensitive. Established Ca^{2+} -
378 mobilising agonists of human sperm such as progesterone and prostaglandin E_1 have been shown to
379 activate CatSper but also a range of other small molecules including environmental pollutants such
380 as 1,1,1-trichloro-2,2-bis(4-chlorophenyl)ethane, 1,1-bis(4-chlorophenyl)-2,2,2-trichloroethane
381 (4,4'-DDT), p,p'-dichlorodiphenyldichloroethylene (p,p'-DDE) and 4-methylbenzylidene camphor (4-
382 MBC) are potent agonists (Schiffer, et al. 2014, Tavares, et al. 2013). In addition, agents used to
383 demonstrate cyclic-nucleotide-activated Ca^{2+} influx (such as 8-Br-AMP) have been shown directly to
384 activate CatSper by binding at the extracellular surface (Brenker, et al. 2012). Thus it is possible that
385 a significant proportion of the pharmacological data that apparently support the existence of
386 multiple Ca^{2+} influx pathways in sperm are misleading and in fact reflect actions of the drugs on Ca^{2+}
387 flux through CatSper channels (Brenker, et al. 2012). Furthermore, experiments using CatSper null
388 mice provide strong evidence that $[\text{Ca}^{2+}]_i$ elevation induced by solubilised ZP is dependent on Ca^{2+}
389 influx through the CatSper channel in the flagellum, which then propagates to the head (Xia and Ren
390 2009) (though see (Cohen, et al. 2014)). Interestingly, the ability of solubilised zona to induce
391 acrosome reaction was not diminished in CatSper-null sperm. These findings not only suggest that
392 CatSper is the primary Ca^{2+} influx pathway in mammalian sperm, but also that in human sperm it
393 may act as a Ca^{2+} -signalling 'hub' or 'node', such that the effects of diverse agonists are
394 summated/integrated in the rate of Ca^{2+} influx into the flagellum (Brenker, et al. 2012). This is an
395 elegant and simple model for which there is already a significant body of data, but in its basic form it
396 does not address the question of how a sperm can generate and use diverse $[\text{Ca}^{2+}]_i$ signals to control
397 diverse Ca^{2+} -sensitive functions.

398 Mouse sperm null for CatSper are unable to hyperactivate (Carlson, et al. 2003) and evidence from
399 clinical cases suggests that CatSper is also required for normal levels of motility in human sperm
400 (Avenarius, et al. 2009, Smith, et al. 2013). Why then, is manipulation of Ca^{2+} stores more effective in
401 inducing hyperactivated motility than treatments targeted to CatSper (Alasmari, et al. 2013b)? We

402 have proposed that CatSper activation acts as a trigger and consequent elevation of flagellar $[Ca^{2+}]_i$
403 stimulates secondary release of stored Ca^{2+} at the sperm neck, either by stimulating synthesis of IP_3
404 or by CICR, leading to hyperactivation (Alasmari, et al. 2013b). Mathematical modelling of the Ca^{2+}
405 signals induced by CatSper activation in mouse sperm suggests that forward diffusion of Ca^{2+} from
406 the flagellum cannot explain the $[Ca^{2+}]_i$ that occurs at the sperm head upon activation of CatSper and
407 that such a secondary Ca^{2+} release at the neck region occurs (Li, et al. 2014, Olson, et al. 2011, Olson,
408 et al. 2010). Recently we have investigated the occurrence of such secondary responses in human
409 sperm by uncaging Ca^{2+} in the principal piece of the flagellum. Uncaging induces a clear $[Ca^{2+}]_i$
410 transient in the flagellum that decays within 5-10 s. At the neck region of the sperm the transient is
411 truncated and rises more slowly, consistent with diffusion of Ca^{2+} from the uncaged pool, but in a
412 small proportion of cells ($\approx 10\%$) we have observed a late $[Ca^{2+}]_i$ response at the neck region, often
413 including multiple peaks (fig 2). The low incidence of this secondary Ca^{2+} -mobilisation is consistent
414 with our observation that, though direct release of stored Ca^{2+} can induce hyperactivated motility in
415 the majority of human sperm, only a small proportion of cells hyperactivate upon activation of
416 CatSper (Alasmari, et al. 2013a, Alasmari, et al. 2013b).

417 Ca^{2+} -store-mediated $[Ca^{2+}]_i$ oscillations occur more readily in sperm incubated for a prolonged period
418 (>24 h) under capacitating conditions (Kirkman-Brown, et al. 2004). Capacitation involves generation
419 of reactive oxygen and reactive nitrogen species (Aitken and Nixon 2013, Herrero, et al. 1999, 2001)
420 and we have observed that store mobilisation is sensitised and induced by low concentrations of NO
421 donors, through a mechanism that involves protein S-nitrosylation (Machado-Oliveira, et al. 2008).
422 RyRs were detected in the human sperm nitrosoproteome (Lefievre, et al. 2007) and it is well-
423 established that IP_3 Rs and RyRs are sensitised by oxidative stress (Bansaghi, et al. 2014, Bootman, et
424 al. 1992, Meissner 2004, Sayers, et al. 1993, Stoyanovsky, et al. 1997) (see section 2). We propose
425 that CICR from the sperm neck Ca^{2+} -store is regulated during capacitation, perhaps through the
426 effects of oxidative stress on Ca^{2+} release channels (Alasmari, et al. 2013b) (fig 3).

427 7. Final remarks

428 The central role of $[Ca^{2+}]_i$ signalling in the physiology of mammalian sperm and the pivotal
429 importance of CatSper in this process are well established - mice null for CatSper are infertile (Ren,
430 et al. 2001) and in men CatSper lesions are associated with impaired sperm function (Avenarius, et
431 al. 2009, Avidan, et al. 2003, Smith, et al. 2013, Zhang, et al. 2009) . The available evidence suggests
432 that Ca^{2+} -stores also play important roles in both acrosome reaction and the regulation of motility.
433 Future studies should address the mechanisms by which store mobilisation is achieved (both by CICR
434 and by direct activation by agonist-induced generation of Ca^{2+} -mobilising 2nd messengers) and
435 regulated, particularly the significance of capacitation in Ca^{2+} -store filling and in sensitising Ca^{2+}
436 release mechanisms. Also, similarly to the important species differences in expression and function
437 of sperm ion channels between human and mouse sperm (Brenker, et al. 2014, Miller, et al. 2014),
438 there may also be differences in store-regulation and/or function between species. An intriguing
439 possibility is that, at least in human sperm, it may prove possible to bypass the effects on motility of
440 lesions in the expression, function or regulation of CatSper channels by pharmacological activation of
441 store Ca^{2+} release.

442 Declaration of interest

443

444 There is no conflict of interest that could be perceived as prejudicing the impartiality of the research
445 reported

446

447 Funding

448

449 JC was supported by the Wellcome Trust (Grant #086470)

450

451 References

- 452 **Adachi T, Weisbrod RM, Pimentel DR, Ying J, Sharov VS, Schoneich C & Cohen RA** 2004 S-
453 Glutathiolation by peroxynitrite activates SERCA during arterial relaxation by nitric oxide.
454 *Nat Med* **10** 1200-1207.
- 455 **Aitken RJ & Nixon B** 2013 Sperm capacitation: a distant landscape glimpsed but unexplored. *Mol*
456 *Hum Reprod* **19** 785-793.
- 457 **Alasmari W, Barratt C, Publicover S, Whalley K, Foster E, Kay V, Silva SMd & Oxenham S** 2013a The
458 incidence and clinical significance of defects in calcium signalling pathways mediating human
459 sperm hyperactivation in donors and sub fertile patients. *Human Reproduction* **28** 866-876.
- 460 **Alasmari W, Costello S, Correia J, Oxenham SK, Morris J, Fernandes L, Ramalho-Santos J, Kirkman-**
461 **Brown J, Michelangeli F, Publicover S & Barratt CLR** 2013b Ca²⁺ signals generated by
462 CatSper and Ca²⁺ regulate different behaviours in human sperm. *J Biol Chem* **288** 6248-6258.
- 463 **Aquila S, Guido C, Middea E, Perrotta I, Bruno R, Pellegrino M & Ando S** 2009 Human male gamete
464 endocrinology: 1alpha, 25-dihydroxyvitamin D3 (1,25(OH)2D3) regulates different aspects of
465 human sperm biology and metabolism. *Reprod Biol Endocrinol* **7** 140.
- 466 **Ardon F, Rodriguez-Miranda E, Beltran C, Hernandez-Cruz A & Darszon A** 2009 Mitochondrial
467 inhibitors activate influx of external Ca²⁺ in sea urchin sperm. *Biochim Biophys Acta* **1787** 15-
468 24.
- 469 **Arndt L, Castonguay J, Arlt E, Meyer D, Hassan S, Borth H, Zierler S, Wennemuth G, Breit A, Biel M,**
470 **Wahl-Schott C, Gudermann T, Klugbauer N & Boekhoff I** 2014 NAADP and the two-pore
471 channel protein 1 participate in the acrosome reaction in mammalian spermatozoa. *Mol Biol*
472 *Cell* **25** 948-964.
- 473 **Avenarius MR, Hildebrand MS, Zhang Y, Meyer NC, Smith LL, Kahrizi K, Najmabadi H & Smith RJ**
474 2009 Human male infertility caused by mutations in the CATSPER1 channel protein. *Am J*
475 *Hum Genet* **84** 505-510.
- 476 **Avidan N, Tamary H, Dgany O, Cattani D, Pariente A, Thulliez M, Borot N, Moati L, Barthelme A,**
477 **Shalmon L, Krasnov T, Ben-Asher E, Olender T, Khen M, Yaniv I, Zaizov R, Shalev H,**
478 **Delaunay J, Fellous M, Lancet D & Beckmann JS** 2003 CATSPER2, a human autosomal
479 nonsyndromic male infertility gene. *Eur J Hum Genet* **11** 497-502.
- 480 **Bansaghi S, Golinar T, Madesh M, Csordas G, RamachandraRao S, Sharma K, Yule DI, Joseph SK &**
481 **Hajnoczky G** 2014 Isoform- and species-specific control of inositol 1,4,5-trisphosphate (IP3)
482 receptors by reactive oxygen species. *J Biol Chem* **289** 8170-8181.
- 483 **Beltramo M, Dardente H, Cayla X & Caraty A** 2014 Cellular mechanisms and integrative timing of
484 neuroendocrine control of GnRH secretion by kisspeptin. *Mol Cell Endocrinol* **382** 387-399.
- 485 **Berridge MJ, Bootman MD & Roderick HL** 2003 Calcium signalling: dynamics, homeostasis and
486 remodelling. *Nat Rev Mol Cell Biol* **4** 517-529.
- 487 **Bezprozvanny I, Watras J & Ehrlich BE** 1991 Bell-shaped calcium-response curves of Ins(1,4,5)P3-
488 and calcium-gated channels from endoplasmic reticulum of cerebellum. *Nature* **351** 751-
489 754.
- 490 **Billington RA, Harper C, Bellomo EA, Publicover S, Barratt CL & Genazzani AA** 2006 Characterization
491 of cyclic adenine dinucleotide phosphate ribose levels in human spermatozoa. *Fertil Steril* **86**
492 891-898.
- 493 **Billington RA, Ho A & Genazzani AA** 2002 Nicotinic acid adenine dinucleotide phosphate (NAADP) is
494 present at micromolar concentrations in sea urchin spermatozoa. *J Physiol* **544** 107-112.
- 495 **Blackmore PF** 1993 Thapsigargin elevates and potentiates the ability of progesterone to increase
496 intracellular free calcium in human sperm: possible role of perinuclear calcium. *Cell Calcium*
497 **14** 53-60.
- 498 **Blomberg Jensen M** 2014 Vitamin D and male reproduction. *Nat Rev Endocrinol* **10** 175-186.
- 499 **Blomberg Jensen M, Bjerrum PJ, Jessen TE, Nielsen JE, Joensen UN, Olesen IA, Petersen JH, Juul A,**
500 **Dissing S & Jorgensen N** 2011 Vitamin D is positively associated with sperm motility and
501 increases intracellular calcium in human spermatozoa. *Hum Reprod* **26** 1307-1317.

502 **Blomberg Jensen M, Jorgensen A, Nielsen JE, Bjerrum PJ, Skalkam M, Petersen JH, Egeberg DL,**
503 **Bangsbo S, Andersen AN, Skakkebaek NE, Juul A, Rajpert-De Meyts E, Dissing S, Leffers H**
504 **& Jorgensen N** 2012 Expression of the vitamin D metabolizing enzyme CYP24A1 at the
505 annulus of human spermatozoa may serve as a novel marker of semen quality. *Int J Androl*
506 **35** 499-510.

507 **Blomberg Jensen M, Nielsen JE, Jorgensen A, Rajpert-De Meyts E, Kristensen DM, Jorgensen N,**
508 **Skakkebaek NE, Juul A & Leffers H** 2010 Vitamin D receptor and vitamin D metabolizing
509 enzymes are expressed in the human male reproductive tract. *Hum Reprod* **25** 1303-1311.

510 **Bootman MD, Taylor CW & Berridge MJ** 1992 The thiol reagent, thimerosal, evokes Ca²⁺ spikes in
511 HeLa cells by sensitizing the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **267** 25113-
512 25119.

513 **Brenker C, Goodwin N, Weyand I, Kashikar ND, Naruse M, Krahling M, Muller A, Kaupp UB &**
514 **Strunker T** 2012 The CatSper channel: a polymodal chemosensor in human sperm. *Embo J* **31**
515 1654-1665.

516 **Brenker C, Zhou Y, Muller A, Echeverry F, Trotschel C, Poetsch A, Xia X, Bonigk W, Lingle C, Kaupp**
517 **U & T S** 2014 Slo3 in human sperm - a K⁺ channel activated by Ca²⁺ *eLife* **3:e01438**.

518 **Cahalan MD** 2009 STIMulating store-operated Ca²⁺ entry. *Nat Cell Biol* **11** 669-677.

519 **Calcraft PJ, Ruas M, Pan Z, Cheng X, Arredouani A, Hao X, Tang J, Rietdorf K, Teboul L, Chuang KT,**
520 **Lin P, Xiao R, Wang C, Zhu Y, Lin Y, Wyatt CN, Parrington J, Ma J, Evans AM, Galione A &**
521 **Zhu MX** 2009 NAADP mobilizes calcium from acidic organelles through two-pore channels.
522 *Nature* **459** 596-600.

523 **Camina JP, Carreira MC, Micic D, Pombo M, Kelestimur F, Dieguez C & Casanueva FF** 2003
524 Regulation of ghrelin secretion and action. *Endocrine* **22** 5-12.

525 **Camors E & Valdivia HH** 2014 CaMKII regulation of cardiac ryanodine receptors and inositol
526 triphosphate receptors. *Front Pharmacol* **5** 101.

527 **Cancela JM, Churchill GC & Galione A** 1999 Coordination of agonist-induced Ca²⁺-signalling patterns
528 by NAADP in pancreatic acinar cells. *Nature* **399** 74-77.

529 **Cang C, Zhou Y, Navarro B, Seo YJ, Aranda K, Shi L, Battaglia-Hsu S, Nissim I, Clapham DE & Ren D**
530 2013 mTOR regulates lysosomal ATP-sensitive two-pore Na⁺ channels to adapt to metabolic
531 state. *Cell* **152** 778-790.

532 **Carlson AE, Westenbroek RE, Quill T, Ren D, Clapham DE, Hille B, Garbers DL & Babcock DF** 2003
533 CatSper1 required for evoked Ca²⁺ entry and control of flagellar function in sperm. *Proc Natl*
534 *Acad Sci U S A* **100** 14864-14868.

535 **Castellano LE, Trevino CL, Rodriguez D, Serrano CJ, Pacheco J, Tsutsumi V, Felix R & Darszon A** 2003
536 Transient receptor potential (TRPC) channels in human sperm: expression, cellular
537 localization and involvement in the regulation of flagellar motility. *FEBS Lett* **541** 69-74.

538 **Chandrashekrana A, Isa I, Dudhia J, Thrasher AJ, Dibb N, Casimir C, Readhead C & Winston R** 2014a
539 Lentiviral vector transduction of spermatozoa as a tool for the study of early development.
540 *FEBS Open Bio* **4** 266-275.

541 **Chandrashekrana A, Sarkar R, Thrasher A, Fraser SE, Dibb N, Casimir C, Winston R & Readhead C**
542 2014b Efficient generation of transgenic mice by lentivirus-mediated modification of
543 spermatozoa. *Faseb J* **28** 569-576.

544 **Chang H & Suarez SS** 2011 Two distinct Ca²⁺ signaling pathways modulate sperm flagellar beating
545 patterns in mice. *Biol Reprod* **85** 296-305.

546 **Chang H & Suarez SS** 2012 Unexpected flagellar movement patterns and epithelial binding behavior
547 of mouse sperm in the oviduct. *Biol Reprod* **86** 140, 141-148.

548 **Chini EN, Thompson MA, Chini CC & Dousa TP** 1997 Cyclic ADP-ribose signaling in sea urchin
549 gametes: metabolism in spermatozoa. *Am J Physiol* **272** C416-420.

550 **Churchill GC, Okada Y, Thomas JM, Genazzani AA, Patel S & Galione A** 2002 NAADP mobilizes Ca²⁺
551 from reserve granules, lysosome-related organelles, in sea urchin eggs. *Cell* **111** 703-708.

552 **Cohen R, Buttke DE, Asano A, Mukai C, Nelson JL, Ren D, Miller RJ, Cohen-Kutner M, Atlas D &**
553 **Travis AJ** 2014 Lipid Modulation of Calcium Flux through CaV2.3 Regulates Acrosome
554 Exocytosis and Fertilization. *Dev Cell* **28** 310-321.

555 **Colyer J** 1998 Phosphorylation states of phospholamban. *Ann N Y Acad Sci* **853** 79-91.

556 **Cosker F, Cheviron N, Yamasaki M, Menteyne A, Lund FE, Moutin MJ, Galione A & Cancela JM.**
557 2010. The ecto-enzyme CD38 is a mammalian NAADP synthase which couples receptor
558 activation to Ca²⁺ mobilization from lysosomes. *J Biol Chem* **285** 38251-38259.

559 **Costello S, Michelangeli F, Nash K, Lefievre L, Morris J, Machado-Oliveira G, Barratt C, Kirkman-**
560 **Brown J & Publicover S** 2009 Ca²⁺ stores in sperm: their identities and functions.
561 *Reproduction* **138** 425-437.

562 **Csordas G & Hajnoczky G** 2009 SR/ER-mitochondrial local communication: calcium and ROS. *Biochim*
563 *Biophys Acta* **1787** 1352-1362.

564 **Darszon A, Nishigaki T, Beltran C & Trevino CL** 2011 Calcium channels in the development,
565 maturation, and function of spermatozoa. *Physiol Rev* **91** 1305-1355.

566 **Darszon A, Sanchez-Cardenas C, Orta G, Sanchez-Tusie AA, Beltran C, Lopez-Gonzalez I, Granados-**
567 **Gonzalez G & Trevino CL** 2012 Are TRP channels involved in sperm development and
568 function? *Cell Tissue Res* **349** 749-764

569 **Darszon A, Trevino CL, Wood C, Galindo B, Rodriguez-Miranda E, Acevedo JJ, Hernandez-Gonzalez**
570 **EO, Beltran C, Martinez-Lopez P & Nishigaki T** 2007 Ion channels in sperm motility and
571 capacitation. *Soc Reprod Fertil Suppl* **65** 229-244.

572 **De Blas G, Michaut M, Trevino CL, Tomes CN, Yunes R, Darszon A & Mayorga LS** 2002 The
573 intraacrosomal calcium pool plays a direct role in acrosomal exocytosis. *J Biol Chem* **277**
574 49326-49331.

575 **De Blas GA, Roggero CM, Tomes CN & Mayorga LS** 2005 Dynamics of SNARE assembly and
576 disassembly during sperm acrosomal exocytosis. *PLoS Biol* **3** e323.

577 **Dragileva E, Rubinstein S & Breitbart H** 1999 Intracellular Ca²⁺-Mg²⁺-ATPase regulates calcium influx
578 and acrosomal exocytosis in bull and ram spermatozoa. *Biol Reprod* **61** 1226-1234.

579 **Drago I, Pizzo P & Pozzan T** 2011 After half a century mitochondrial calcium in- and efflux
580 machineries reveal themselves. *Embo J* **30** 4119-4125.

581 **Dyer JL, Mobasher H, Lea EJ, Dawson AP & Michelangeli F** 2003 Differential effect of PKA on the
582 Ca²⁺ release kinetics of the type I and III InsP3 receptors. *Biochem Biophys Res Commun* **302**
583 121-126.

584 **Espino J, Mediero M, Lozano GM, Bejarano I, Ortiz A, Garcia JF, Pariente JA & Rodriguez AB** 2009
585 Reduced levels of intracellular calcium releasing in spermatozoa from asthenozoospermic
586 patients. *Reprod Biol Endocrinol* **7** 11.

587 **Feng M, Grice DM, Faddy HM, Nguyen N, Leitch S, Wang Y, Muend S, Kenny PA, Sukumar S,**
588 **Roberts-Thomson SJ, Monteith GR & Rao R** 2010 Store-independent activation of Orai1 by
589 SPCA2 in mammary tumors. *Cell* **143** 84-98.

590 **Florman HM, Jungnickel MK & Sutton KA** 2008 Regulating the acrosome reaction. *Int J Dev Biol* **52**
591 503-510.

592 **Fukami K, Nakao K, Inoue T, Kataoka Y, Kurokawa M, Fissore RA, Nakamura K, Katsuki M,**
593 **Mikoshiba K, Yoshida N & Takenawa T** 2001 Requirement of phospholipase Cdelta4 for the
594 zona pellucida-induced acrosome reaction. *Science* **292** 920-923.

595 **Fukami K, Yoshida M, Inoue T, Kurokawa M, Fissore RA, Yoshida N, Mikoshiba K & Takenawa T**
596 2003 Phospholipase Cdelta4 is required for Ca²⁺ mobilization essential for acrosome reaction
597 in sperm. *J Cell Biol* **161** 79-88.

598 **Genazzani AA, Mezna M, Summerhill RJ, Galione A & Michelangeli F** 1997 Kinetic properties of
599 nicotinic acid adenine dinucleotide phosphate-induced Ca²⁺ release. *J Biol Chem* **272** 7669-
600 7675.

601 **Gur Y & Breitbart H** 2007 Protein translation in mammalian sperm. *Soc Reprod Fertil Suppl* **65** 391-
602 397.

603 **Guse AH** 2014 Calcium mobilizing second messengers derived from NAD. *Biochim Biophys Acta*.
604 **Hamilton SL & Reid MB** 2000 RyR1 modulation by oxidation and calmodulin. *Antioxid Redox Signal* **2**
605 41-45.
606 **Harper C, Wootton L, Michelangeli F, Lefievre L, Barratt C & Publicover S** 2005 Secretory pathway
607 Ca^{2+} -ATPase (SPCA1) Ca^{2+} pumps, not SERCAs, regulate complex $[\text{Ca}^{2+}]_i$ signals in human
608 spermatozoa. *J Cell Sci* **118** 1673-1685.
609 **Harper CV, Barratt CL & Publicover SJ** 2004 Stimulation of human spermatozoa with progesterone
610 gradients to simulate approach to the oocyte. Induction of $[\text{Ca}^{2+}]_i$ oscillations and cyclical
611 transitions in flagellar beating. *J Biol Chem* **279** 46315-46325.
612 **Herrero MB, de Lamirande E & Gagnon C** 1999 Nitric oxide regulates human sperm capacitation and
613 protein-tyrosine phosphorylation in vitro. *Biol Reprod* **61** 575-581.
614 **Herrero MB, de Lamirande E & Gagnon C** 2001 Tyrosine nitration in human spermatozoa: a
615 physiological function of peroxynitrite, the reaction product of nitric oxide and superoxide.
616 *Mol Hum Reprod* **7** 913-921.
617 **Ho HC & Suarez SS** 2001 An inositol 1,4,5-trisphosphate receptor-gated intracellular Ca^{2+} store is
618 involved in regulating sperm hyperactivated motility. *Biol Reprod* **65** 1606-1615.
619 **Ho HC & Suarez SS** 2003 Characterization of the intracellular calcium store at the base of the sperm
620 flagellum that regulates hyperactivated motility. *Biol Reprod* **68** 1590-1596.
621 **Hsu MC, Wang JY, Lee YJ, Jong DS, Tsui KH & Chiu CH** 2014 Kisspeptin modulates fertilization
622 capacity of mouse spermatozoa. *Reproduction* **147** 835-845.
623 **Jin M, Fujiwara E, Kakiuchi Y, Okabe M, Satouh Y, Baba SA, Chiba K & Hirohashi N** 2011 Most
624 fertilizing mouse spermatozoa begin their acrosome reaction before contact with the zona
625 pellucida during in vitro fertilization. *Proc Natl Acad Sci U S A* **108** 4892-4896.
626 **Kirkman-Brown JC, Barratt CL & Publicover SJ** 2004 Slow calcium oscillations in human
627 spermatozoa. *Biochem J* **378** 827-832.
628 **Kuroda Y, Kaneko S, Yoshimura Y, Nozawa S & Mikoshiba K** 1999 Are there inositol 1,4,5-
629 triphosphate (IP3) receptors in human sperm? *Life Sci* **65** 135-143.
630 **Lawson C, Dorval V, Goupil S & Leclerc P** 2007 Identification and localisation of SERCA 2 isoforms in
631 mammalian sperm. *Mol Hum Reprod* **13** 307-316.
632 **Lefievre L, Chen Y, Conner SJ, Scott JL, Publicover SJ, Ford WC & Barratt CL** 2007 Human
633 spermatozoa contain multiple targets for protein S-nitrosylation: an alternative mechanism
634 of the modulation of sperm function by nitric oxide? *Proteomics* **7** 3066-3084.
635 **Lefievre L, Nash K, Mansell S, Costello S, Punt E, Correia J, Morris J, Kirkman-Brown J, Wilson SM,**
636 **Barratt CL & Publicover S** 2012 2-APB-potentiated channels amplify CatSper-induced Ca^{2+}
637 signals in human sperm. *Biochem J* **448** 189-200.
638 **Li LF, Xiang C, Zhu YB & Qin KR** 2014 Modeling of progesterone-induced intracellular calcium
639 signaling in human spermatozoa. *J Theor Biol* **351** 58-66.
640 **Lin-Moshier Y, Walseth TF, Churamani D, Davidson SM, Slama JT, Hooper R, Brailoiu E, Patel S &**
641 **Marchant JS** 2012 Photoaffinity labeling of nicotinic acid adenine dinucleotide phosphate
642 (NAADP) targets in mammalian cells. *J Biol Chem* **287** 2296-2307.
643 **Lishko PV, Botchkina IL & Kirichok Y** 2011 Progesterone activates the principal Ca^{2+} channel of
644 human sperm. *Nature* **471** 387-391.
645 **Liu X, Lee K & Herbison AE** 2008 Kisspeptin excites gonadotropin-releasing hormone neurons
646 through a phospholipase C/calcium-dependent pathway regulating multiple ion channels.
647 *Endocrinology* **149** 4605-4614.
648 **Luik RM, Wu MM, Buchanan J & Lewis RS** 2006 The elementary unit of store-operated Ca^{2+} entry:
649 local activation of CRAC channels by STIM1 at ER-plasma membrane junctions. *J Cell Biol* **174**
650 815-825.
651 **Lukaszyk A, Kotwicka M, Jankowska A, Kasprzak A, Rucinski M, Sterzynska K, Ziolkowska A,**
652 **Sawinski P & Ruchala M** 2012 Expression of ghrelin receptor (GHSR-1a) in rat epididymal
653 spermatozoa and the effects of its activation. *Reprod Biol* **12** 293-300.

654 **Luo T, Li N, He YQ, Weng SQ, Wang T, Zou QX & Zeng XH** 2014 Emodin inhibits human sperm
655 functions by reducing sperm [Ca] and tyrosine phosphorylation. *Reprod Toxicol* **51C** 14-21.
656 **Machado-Oliveira G, Lefievre L, Ford C, Herrero MB, Barratt C, Connolly TJ, Nash K, Morales-Garcia**
657 **A, Kirkman-Brown J & Publicover S** 2008 Mobilisation of Ca²⁺ stores and flagellar regulation
658 in human sperm by S-nitrosylation: a role for NO synthesised in the female reproductive
659 tract. *Development* **135** 3677-3686.
660 **Marquez B, Ignatz G & Suarez SS** 2007 Contributions of extracellular and intracellular Ca²⁺ to
661 regulation of sperm motility: Release of intracellular stores can hyperactivate CatSper1 and
662 CatSper2 null sperm. *Dev Biol* **303** 214-221.
663 **Mayorga LS, Tomes CN & Belmonte SA** 2007 Acrosomal exocytosis, a special type of regulated
664 secretion. *IUBMB Life* **59** 286-292.
665 **Meissner G** 2004 Molecular regulation of cardiac ryanodine receptor ion channel. *Cell Calcium* **35**
666 621-628.
667 **Menteyne A, Burdakov A, Charpentier G, Petersen OH & Cancela JM** 2006 Generation of specific
668 Ca²⁺ signals from Ca²⁺ stores and endocytosis by differential coupling to messengers. *Curr*
669 *Biol* **16** 1931-1937.
670 **Michelangeli F & East JM** 2011 A diversity of SERCA Ca²⁺ pump inhibitors. *Biochem Soc Trans* **39** 789-
671 797.
672 **Michelangeli F, Ogunbayo OA & Wootton LL** 2005 A plethora of interacting organellar Ca²⁺ stores.
673 *Curr Opin Cell Biol* **17** 135-140.
674 **Miller D & Ostermeier GC** 2006 Spermatozoal RNA: Why is it there and what does it do? *Gynecol*
675 *Obstet Fertil* **34** 840-846.
676 **Miller D, Ostermeier GC & Krawetz SA** 2005 The controversy, potential and roles of spermatozoal
677 RNA. *Trends Mol Med* **11** 156-163.
678 **Miller MR, Mansell SA, Meyers SA & Lishko PV** 2014 Flagellar ion channels of sperm: similarities and
679 differences between species. *Cell Calcium*.
680 **Missiaen L, Taylor CW & Berridge MJ** 1991 Spontaneous calcium release from inositol
681 trisphosphate-sensitive calcium stores. *Nature* **352** 241-244.
682 **Moretti E, Vindigni C, Tripodi SA, Mazzi L, Nuti R, Figura N & Collodel G** 2014 Immunolocalisation of
683 ghrelin and obestatin in human testis, seminal vesicles, prostate and spermatozoa.
684 *Andrologia* **46** 979-985.
685 **Morgan AJ & Galione A** 2014 Two-pore channels (TPCs): current controversies. *Bioessays* **36** 173-
686 183.
687 **Morris J, Jones S, Howl J, Lukanowska M, Lefievre L & Publicover S** 2015 Cell penetrating peptides,
688 targeting the regulation of store-operated channels, slow decay of the progesterone-
689 induced [Ca²⁺]_i signal in human sperm. *Mol Hum Reprod* pii: gav019. [Epub ahead of print]
690 **Naaby-Hansen S, Wolkowicz MJ, Klotz K, Bush LA, Westbrook VA, Shibahara H, Shetty J, Coonrod**
691 **SA, Reddi PP, Shannon J, Kinter M, Sherman NE, Fox J, Flickinger CJ & Herr JC** 2001 Co-
692 localization of the inositol 1,4,5-trisphosphate receptor and calreticulin in the equatorial
693 segment and in membrane bounded vesicles in the cytoplasmic droplet of human
694 spermatozoa. *Mol Hum Reprod* **7** 923-933.
695 **Navarro-Borelly L, Somasundaram A, Yamashita M, Ren D, Miller RJ & Prakriya M** 2008 STIM1-
696 Orai1 interactions and Orai1 conformational changes revealed by live-cell FRET microscopy. *J*
697 *Physiol* **586** 5383-5401.
698 **O'Toole CM, Arnoult C, Darszon A, Steinhardt RA & Florman HM** 2000 Ca²⁺ entry through store-
699 operated channels in mouse sperm is initiated by egg ZP3 and drives the acrosome reaction.
700 *Mol Biol Cell* **11** 1571-1584.
701 **Ogunbayo OA, Zhu Y, Rossi D, Sorrentino V, Ma J, Zhu MX & Evans AM** 2011 Cyclic adenosine
702 diphosphate ribose activates ryanodine receptors, whereas NAADP activates two-pore
703 domain channels. *J Biol Chem* **286** 9136-9140.

704 **Olson SD, Fauci LJ & Suarez SS** 2011 Mathematical modeling of calcium signaling during sperm
705 hyperactivation. *Mol Hum Reprod* **17** 500-510.

706 **Olson SD, Suarez SS & Fauci LJ** 2010 A model of CatSper channel mediated calcium dynamics in
707 mammalian spermatozoa. *Bull Math Biol* **72** 1925-1946.

708 **Park KH, Kim BJ, Kang J, Nam TS, Lim JM, Kim HT, Park JK, Kim YG, Chae SW & Kim UH** 2011 Ca²⁺
709 signaling tools acquired from prostasomes are required for progesterone-induced sperm
710 motility. *Sci Signal* **4** ra31.

711 **Peiter E, Maathuis FJ, Mills LN, Knight H, Pelloux J, Hetherington AM & Sanders D** 2005 The
712 vacuolar Ca²⁺-activated channel TPC1 regulates germination and stomatal movement.
713 *Nature* **434** 404-408.

714 **Pielecka-Fortuna J, Chu Z & Moenter SM** 2008 Kisspeptin acts directly and indirectly to increase
715 gonadotropin-releasing hormone neuron activity and its effects are modulated by estradiol.
716 *Endocrinology* **149** 1979-1986.

717 **Pinto FM, Cejudo-Roman A, Ravina CG, Fernandez-Sanchez M, Martin-Lozano D, Illanes M, Tena-
718 Sempere M & Cadenas ML** 2012 Characterization of the kisspeptin system in human
719 spermatozoa. *Int J Androl* **35** 63-73.

720 **Pizzo P, Drago I, Filadi R & Pozzan T** 2012 Mitochondrial Ca²⁺ homeostasis: mechanism, role, and
721 tissue specificities. *Pflugers Arch* **464** 3-17.

722 **Publicover S, Harper CV & Barratt C** 2007 [Ca²⁺]_i signalling in sperm--making the most of what
723 you've got. *Nat Cell Biol* **9** 235-242.

724 **Ren D, Navarro B, Perez G, Jackson AC, Hsu S, Shi Q, Tilly JL & Clapham DE** 2001 A sperm ion
725 channel required for sperm motility and male fertility. *Nature* **413** 603-609.

726 **Ren D & Xia J** 2010 Calcium signaling through CatSper channels in mammalian fertilization.
727 *Physiology (Bethesda)* **25** 165-175.

728 **Ruete MC, Lucchesi O, Bustos MA & Tomes CN** 2014 Epac, Rap and Rab3 act in concert to mobilize
729 calcium from sperm's acrosome during exocytosis. *Cell Commun Signal* **12** 43.

730 **Sanchez-Tusie AA, Vasudevan SR, Churchill GC, Nishigaki T & Trevino CL** 2014 Characterization of
731 NAADP-mediated calcium signaling in human spermatozoa. *Biochem Biophys Res Commun*
732 **443** 531-536.

733 **Sayers LG, Brown GR, Michell RH & Michelangeli F** 1993 The effects of thimerosal on calcium
734 uptake and inositol 1,4,5-trisphosphate-induced calcium release in cerebellar microsomes.
735 *Biochem J* **289 (Pt 3)** 883-887.

736 **Schiffer C, Muller A, Egeberg DL, Alvarez L, Brenker C, Rehfeld A, Frederiksen H, Waschle B, Kaupp
737 UB, Balbach M, Wachten D, Skakkebaek NE, Almstrup K & Strunker T** 2014 Direct action of
738 endocrine disrupting chemicals on human sperm. *EMBO Rep* **15** 758-765.

739 **Seo MD, Velamakanni S, Ishiyama N, Stathopoulos PB, Rossi AM, Khan SA, Dale P, Li C, Ames JB,
740 Ikura M & Taylor CW** 2012 Structural and functional conservation of key domains in InsP3
741 and ryanodine receptors. *Nature* **483** 108-112.

742 **Sharov VS, Dremina ES, Galeva NA, Williams TD & Schoneich C** 2006 Quantitative mapping of
743 oxidation-sensitive cysteine residues in SERCA in vivo and in vitro by HPLC-electrospray-
744 tandem MS: selective protein oxidation during biological aging. *Biochem J* **394** 605-615.

745 **Smith JF, Syritsyna O, Fellous M, Serres C, Mannowetz N, Kirichok Y & Lishko PV** 2013 Disruption of
746 the principal, progesterone-activated sperm Ca²⁺ channel in a CatSper2-deficient infertile
747 patient. *Proc Natl Acad Sci U S A* **110** 6323-6328.

748 **Stoyanovsky D, Murphy T, Anno PR, Kim YM & Salama G** 1997 Nitric oxide activates skeletal and
749 cardiac ryanodine receptors. *Cell Calcium* **21** 19-29.

750 **Strunker T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R & Kaupp UB** 2011 The CatSper
751 channel mediates progesterone-induced Ca²⁺ influx in human sperm. *Nature* **471** 382-386.

752 **Tavares RS, Mansell S, Barratt CL, Wilson SM, Publicover SJ & Ramalho-Santos J** 2013 p,p'-DDE
753 activates CatSper and compromises human sperm function at environmentally relevant
754 concentrations. *Hum Reprod* **28** 3167-3177.

- 755 **Thomas P & Meizel S** 1989 Phosphatidylinositol 4,5-bisphosphate hydrolysis in human sperm
756 stimulated with follicular fluid or progesterone is dependent upon Ca^{2+} influx. *Biochem J* **264**
757 539-546.
- 758 **Trevino CL, Santi CM, Beltran C, Hernandez-Cruz A, Darszon A & Lomeli H** 1998 Localisation of
759 inositol trisphosphate and ryanodine receptors during mouse spermatogenesis: possible
760 functional implications. *Zygote* **6** 159-172.
- 761 **Vanoevelen J, Dode L, Van Baelen K, Fairclough RJ, Missiaen L, Raeymaekers L & Wuytack F** 2005
762 The secretory pathway $\text{Ca}^{2+}/\text{Mn}^{2+}$ -ATPase 2 is a Golgi-localized pump with high affinity for
763 Ca^{2+} ions. *J Biol Chem* **280** 22800-22808.
- 764 **Vasudevan SR, Galione A & Churchill GC** 2008 Sperm express a Ca^{2+} -regulated NAADP synthase.
765 *Biochem J* **411** 63-70.
- 766 **Wang X, Zhang X, Dong XP, Samie M, Li X, Cheng X, Goschka A, Shen D, Zhou Y, Harlow J, Zhu MX,**
767 **Clapham DE, Ren D & Xu H** 2012 TPC proteins are phosphoinositide- activated sodium-
768 selective ion channels in endosomes and lysosomes. *Cell* **151** 372-383.
- 769 **Wang Y, Deng X, Zhou Y, Hendron E, Mancarella S, Ritchie MF, Tang XD, Baba Y, Kurosaki T, Mori Y,**
770 **Soboloff J & Gill DL** 2009 STIM protein coupling in the activation of Orai channels. *Proc Natl*
771 *Acad Sci U S A* **106** 7391-7396.
- 772 **Wootton LL, Argent CC, Wheatley M & Michelangeli F** 2004 The expression, activity and localisation
773 of the secretory pathway Ca^{2+} -ATPase (SPCA1) in different mammalian tissues. *Biochim*
774 *Biophys Acta* **1664** 189-197.
- 775 **Wu MM, Buchanan J, Luik RM & Lewis RS** 2006 Ca^{2+} store depletion causes STIM1 to accumulate in
776 ER regions closely associated with the plasma membrane. *J Cell Biol* **174** 803-813.
- 777 **Xia J & Ren D** 2009 Egg coat proteins activate calcium entry into mouse sperm via CATSPER
778 channels. *Biol Reprod* **80** 1092-1098.
- 779 **Yamashita M, Somasundaram A & Prakriya M** 2011 Competitive modulation of Ca^{2+} release-
780 activated Ca^{2+} channel gating by STIM1 and 2-aminoethyldiphenyl borate. *J Biol Chem* **286**
781 9429-9442.
- 782 **Yule DI, Betzenhauser MJ & Joseph SK** 2010 Linking structure to function: Recent lessons from
783 inositol 1,4,5-trisphosphate receptor mutagenesis. *Cell Calcium* **47** 469-479.
- 784 **Zeng XH, Navarro B, Xia XM, Clapham DE & Lingle CJ** 2013 Simultaneous knockout of Slo3 and
785 CatSper1 abolishes all alkalization- and voltage-activated current in mouse spermatozoa. *J*
786 *Gen Physiol* **142** 305-313.
- 787 **Zhang Y, Malekpour M, Al-Madani N, Kahrizi K, Zanganeh M, Mohseni M, Mojahedi F, Daneshi A,**
788 **Najmabadi H & Smith RJ** 2009 Sensorineural deafness and male infertility: a contiguous
789 gene deletion syndrome. *BMJ Case Rep* **2009**.
- 790 **Zhao C, Guo XJ, Shi ZH, Wang FQ, Huang XY, Huo R, Zhu H, Wang XR, Liu JY, Zhou ZM & Sha JH** 2009
791 Role of translation by mitochondrial-type ribosomes during sperm capacitation: an analysis
792 based on a proteomic approach. *Proteomics* **9** 1385-1399.
- 793 **Zitanski N, Borth H, Ackermann F, Meyer D, Vieweg L, Breit A, Gudermann T & Boekhoff I** 2010
794 The "acrosomal synapse": Subcellular organization by lipid rafts and scaffolding proteins
795 exhibits high similarities in neurons and mammalian spermatozoa. *Commun Integr Biol* **3**
796 513-521.

797

798

799

800

801 Figure legends

802 **Fig.1 a:** Simplified diagrammatic summary of $[Ca^{2+}]_i$ signalling toolkit in a somatic cell. Ion channels
803 are shown as rectangles with arrow indicating normal direction of Ca^{2+} flow (yellow=voltage gated;
804 green=ligand gated; purple=store-operated; light blue= IP_3 receptor; dark blue=ryanodine receptor;
805 red=NAADP-gated. Pumps are shown as circles with arrows indicating normal direction of Ca^{2+}
806 movement (red=PMCA'; blue= Na^+ - Ca^{2+} exchanger; green=SERCA; blue=SPCA). Activation of IP_3
807 receptors by membrane receptor activation and phospholipase C is shown in light blue. Generation
808 of cADPR and NAADP by CD38 and possibly other enzymes (leading to mobilisation of Ca^{2+} from
809 intracellular stores) is shown by yellow boxes. **b:** Structure of human sperm showing positions of
810 CatSper channels (yellow shading around anterior flagellum) and Ca^{2+} stores in the acrosome and at
811 the sperm (neck redundant nuclear envelope and calreticulin-containing vesicles) (shown in green).

812 **Fig. 2.** Ca^{2+} responses evoked in human sperm by uncaging of Ca^{2+} in the flagellum. Cells were
813 labelled with fluo-4 and loaded with 'caged' Ca^{2+} (NP-EGTA), then stimulated by an uncaging flash
814 (360 nm laser) at the central flagellum (shown by arrow) whilst collecting images at 33 Hz. Changes
815 in fluorescence, assessed at each of the positions shown by coloured circles in panel 'a', are plotted
816 (normalised to minimum and maximum) in panel 'b' using the same colour code. Green=neck;
817 yellow-midpiece; red=proximal flagellum; light blue=mid-distal flagellum; dark blue=distal flagellum.

818 **Fig. 3.** Model for triggering/regulation of CatSper-activated hyperactivation. CatSper channels in the
819 flagellum (yellow box; shown by yellow shading on sperm flagellum) are activated by diverse stimuli
820 including intracellular pH (pH_i), membrane potential (E_m), progesterone, prostaglandins and other
821 organic molecules. Ca^{2+} from the flagellum diffuses forward, raising $[Ca^{2+}]_i$ at the sperm neck and can
822 mobilise stored Ca^{2+} by Ca^{2+} -induced Ca^{2+} release (CICR). Susceptibility of the store to CICR is
823 potentially regulated/sensitised by processes occurring during capacitation including cAMP
824 signalling, oxidative stress, S-nitrosylation as well as Ca^{2+} store filling and effects of agonists on Ca^{2+} -
825 store release channels.

828

829




Figure 3