

cdr

Scenario-Led Habitat Modelling of Land Use Change Impacts on Key Species

Item Type	Article
Authors	Geary, Matthew;Fielding, Alan H.;McGowan, Philip J. K.;Marsden, Stuart J.
Citation	Geary, M., Fielding, A. H., McGowan, P. J. K., & Marsden, S. J. (2015). Scenario-led habitat modelling of land use change Impacts on key species. PLoS ONE 10(11): e0142477. doi:10.1371/journal.pone.0142477
Publisher	Public Library of Science
Journal	PLoS ONE
Download date	2026-05-12 07:11:56
Link to Item	http://hdl.handle.net/10034/592499

Scenario-led Habitat Modelling of Land Use Change Impacts on Key Species

Matthew Geary^{1#a*}, Alan H. Fielding¹, Philip J.K. McGowan^{2#b} & Stuart J. Marsden¹

¹Division of Biology & Conservation Ecology, School of Science & the Environment,
Manchester Metropolitan University, Manchester, UK

²Present address: School of Biology, Newcastle University, Newcastle upon Tyne UK

^{#a}Present address: Department of Biological Sciences, University of Chester, Chester, UK

^{#b}World Pheasant Association, Newcastle University Biology Field Station, Close House Estate,
Heddon-on-the-Wall, UK

* Corresponding author

M.geary@Chester.ac.uk (MG)

Abstract

Accurate predictions of the impacts of future land use change on species of conservation concern
5 can help to inform policy-makers and improve conservation measures. If predictions are spatially
explicit, predicted consequences of likely land use changes could be accessible to land managers
at a scale relevant to their working landscape. We introduce a method, based on open source
software, which integrates habitat suitability modelling with scenario-building, and illustrate its
use by investigating the effects of alternative land use change scenarios on landscape suitability
10 for black grouse *Tetrao tetrix*. Expert opinion was used to construct five near-future (twenty
years) scenarios for the 800 km² study site in upland Scotland. For each scenario, the cover of
different land use types was altered by 5-30% from 20 random starting locations and changes in
habitat suitability assessed by projecting a MaxEnt suitability model onto each simulated
landscape. A scenario converting grazed land to moorland and open forestry was the most
15 beneficial for black grouse, and ‘increased grazing’ (the opposite conversion) the most
detrimental. Positioning of new landscape blocks was shown to be important in some situations.
Increasing the area of open-canopy forestry caused a proportional decrease in suitability, but
suitability gains for the ‘reduced grazing’ scenario were nonlinear. ‘Scenario-led’ landscape
simulation models can be applied in assessments of the impacts of land use change both on
20 individual species and also on diversity and community measures, or ecosystem services. A next
step would be to include landscape configuration more explicitly in the simulation models, both
to make them more realistic, and to examine the effects of habitat placement more thoroughly. In
this example, the recommended policy would be incentives on grazing reduction to benefit black
grouse.

25

Keywords: Landscape simulation modelling, land use change scenarios, conservation planning, policy, Scotland, uplands, black grouse.

Introduction

30

Land use change has been shown to be an important driver of population change in a wide variety of taxa [1], with changes such as conversion of forest to agricultural land reducing habitat availability for species throughout the world [2]. Even subtle changes to the way the landscape is managed, such as the timing of ploughing cereal fields [3], can have a significant impact on the ecology of wildlife within an area [4]. Working landscapes, such as farmland or commercial forests, are often compromises between the interests of biodiversity preservation and economic benefit [5]. How much biodiversity persists within working landscapes, of course, varies widely, but, with careful planning, wildlife can thrive alongside human land use. For example, an area in Oregon, USA, retained around 97% of the biological value while still maintaining 92% of its economic value [6].

40

The Scottish uplands are a working landscape containing a mosaic of grouse moor, deciduous and coniferous woodland, and pasture [7]. The composition of the landscape is, of course, dynamic, with grant schemes such as the Scottish Rural Development Programme [8] influencing the amount and placement of various land use types. With these changes come changes in wildlife populations, some species being seriously affected [9,10]. One such species is the black grouse *Tetrao tetrix*, which experienced serious declines throughout the United Kingdom during the twentieth century [11] but which is expected to benefit from land use change brought about through both payments to land-owners and wind farm habitat management plans [8]. Black grouse is a bird of woodland edges and requires a habitat mosaic which can

50

consist of mixtures of moorland, forest and agricultural land [12,13], features which have been altered by land use changes in the UK uplands over the past decades[14].

Modelling species distributions in terms of their associations with habitats, land use or
55 environmental factors is a rapidly advancing area of ecological research [15]. Many applications
of these models have been static, assessing species distributions at a 'snapshot' in time [16].
Static species distribution modelling can inform conservationists about current [17] and potential
distributions [18] or population connectivity [19]. A logical extension is to project organism-
environment relationships into the future for example, to predict species distributions under
60 climate change scenarios [20]. Informed environmental policy can be of great benefit to
conservation [21,22] but the challenge for ecologists is to maximise the relevance and
accessibility of conservation research to policy-makers. To this end, it is important that the
projection of species distributions onto future landscape scenarios is ecologically realistic, and
therefore, restricts the projection to the shorter term and expand species distribution models to
65 include potential habitat-management or land use changes. Scenario-led models allow the
potential impacts of policy or conservation action to be quantified and compared [6,23].

Here we demonstrate the use of open source software to simulate land use change in the Scottish
uplands and its effect on black grouse. We developed five land use change scenarios: reduced
70 grazing, increased grazing, increased grouse moor (heather moorland actively managed to
provide high red grouse *Lagopus lagopus* densities for driven shooting), increased closed-canopy
forestry and increased open-canopy forestry to reflect potential land use changes in the study
area which may have a positive or negative impact on the species [14,24]. We 'grew' patches of

the new habitat, using cellular automata according to the land use change scenarios, in randomly
75 chosen positions to produce a range of estimates for the impact of the land use change on black
grouse over multiple runs. We then examined the effects of the landscape changes in more detail
by looking for nonlinearities in response as the land use change became more extensive. Finally,
we discuss the potential that scenario-led landscape simulation modelling has as a practical tool
for policy-makers interested in integrating conservation objectives and land use policy.

80

Materials and Methods

Study area and lek location data

The study area is 800 km² around Loch Tummel in Perthshire, Scotland (56°42'40"N 3°55'40"W).

85 Altitude ranges from 46 to 1078 m above sea level. Forestry, agriculture and moorland managed for shooting game (generally hunting of red deer *Cervus elaphus* and red grouse) are the main land uses in the area which is representative of black grouse habitat within working landscapes in much of Scotland. In order to demonstrate our modelling method, we required good quality data for both the location of black grouse display sites (*i.e.* leks; areas at which male black grouse display in order to obtain access
90 to females) along with a clear satellite image on which to base a habitat map. With this in mind, rather than demonstrating the modelling approach with a lower quality satellite image and the most recently available black grouse data we chose instead to base these simulations on 1994, a year in which the black grouse population was relatively large (663 displaying males compared with a mean number of 353 displaying males over the next 15 years), was thought to have been counted fully and for which a
95 clear Landsat image was available.

Lek locations used in this study were provided by Perthshire Black Grouse Study Group (PBGSG). The PBGSG is a group of volunteers, including core members from professional bodies including the Game and Wildlife Conservation Trust (GWCT), Royal Society for the Protection of Birds (RSPB) and
100 Forestry Commission. A core of a few volunteers surveyed large areas for several years (e.g. Forestry Commission land) while a number of volunteers surveyed smaller areas in just a few years. These data represent one of many years of a long-term study undertaken by PBGSG (1990-2008) and which used methods similar to those used in the two national black grouse surveys [11,25] covering as much of the landscape as possible rather than using transect methods. Volunteers walked within 0.5 km of all

105 suitable habitat [26] while listening for male calls to locate lek sites. These sites were subsequently
visited twice more, each up to two hours after dawn, to verify the presence of a lek and to count the
number of displaying males [25]. As the purpose of the survey was to count the number of displaying
males at existing leks, absences were not recorded systematically. Areas above 550 m (18% of the
study area) were considered unsuitable for black grouse leks by Hancock et al. [25] and so were not
110 searched by PBGS, and nor were dense forestry, built up areas and arable farmland. Surveys usually
focus attention on likely areas to search so as not to spend effort in areas known to be outside the range
for the target species. This was the case for the black grouse surveys which excluded areas above 550
m. However, below that altitude, there is still much local heterogeneity in habitat suitability for black
grouse [24] which is what the survey method attempted to capture [26].

115

Environmental data and habitat suitability modelling

National land cover maps are available for the UK but these have received criticism, particularly with
respect to their inability to accurately identify upland habitat types [27,28]. It therefore made sense to
120 develop our own habitat groupings based on unsupervised classification coupled with a good working
knowledge of the study area. Habitat data were taken from the USGS LANDSAT image at 30 m
resolution (path 206, row 20) from 1994; as the image for this region was clear, atmospheric correction
was not applied. The original image consisted of eight spectral bands of which six were combined
using principal components analysis (PCA; in ArcMap 9.2), chosen as the most straightforward method
125 in the available software, to remove redundant information. The thermal infra-red band and the
panchromatic band were omitted from the classification. The processed image, consisting of three
principal components, was classified into habitat types using an unsupervised classification in
Multispec [29]. As the satellite image was from 1994, contemporaneous land cover information was
unavailable; therefore unsupervised classification was preferred [30]. The k-means algorithm is a

130 simple and efficient algorithm which has two steps that are repeated (iterated) until an optimisation is reached [31]. Initially the data (pixels) are partitioned into a defined number of clusters, usually at random. Cluster centroids ('averages') are calculated and each case is moved to its most similar cluster centre. The cluster centres are recalculated and cases are again moved. These steps are repeated, leading to increasingly homogeneous clusters, until an endpoint is reached. Many habitats, particularly
135 woodland, agricultural grasslands and large anthropogenic features are clearly identifiable on a false composite colour image of the satellite scene, and these are correctly represented in the resulting automated classification. This classification resulted in 18 habitat types which were subsequently pooled into six broad categories using detailed knowledge of the study area, stakeholder interviews, field visits, and more recent (2005 & 2012) aerial photography as a reference. This photography was
140 used to confirm features which were likely to have remained fixed during this period (e.g. plantation forest edges, water bodies and field boundaries) and was used in conjunction with the other methods of verifying the landscape classification and tested with comparison to an existing land cover map from 1990 (S1 File). These broad categories were: human-dominated landscapes (buildings, urban areas, roads and roadside verges); grazed land (managed enclosed grasslands with relatively productive
145 grasses, as well as open grazed land dominated by rough, poor quality grassland such as *Nardus* and *Molinia*); moorland (open land usually dominated by heather *Calluna vulgaris*); open-canopy forestry; closed-canopy forestry; and water bodies. Proportions of each habitat type within a 2 km radius of each grid cell were calculated using the focal command in the 'raster' package [32] in R [33]. A radius of 2 km was considered to give a conservative estimate of the territory used by individuals throughout the
150 year [34,35]. This proof of concept study, as with many studies on black grouse, was based on lek location positions although it could be repeated or extended by using feeding locations, or locations of birds at particular life stages or in different seasons. A raster giving the altitude of each grid cell was

obtained from Ordnance Survey (Ordnance Survey data: OS Landform PROFILE; 50 m resolution), then cropped and converted to the same resolution (28.5 m) and extent as the habitat maps.

155

Habitat suitability modelling was performed using MaxEnt [36] within the 'dismo' package [37] in R using seven predictors (the proportion of six habitat types plus altitude). MaxEnt is a presence-only, machine learning process and has been shown to outperform other presence-only habitat suitability modelling methods [36,38]. MaxEnt produces values for the relative suitability of each pixel of a map (i.e. relative to the rest of the landscape used in the model rather than probability of presence [39]). We tested a range of values for the regularization parameter (β) within our models (values of 1, 2, 3, 5, 7, 9, 10, 11, 13, 15, 17, and 19) and selected the 'best' value for our model using the value with the lowest AIC score [40,41]. Default settings were used for all other variables within our model fitting 10 crossvalidated replicates [24]. Full details of the MaxEnt model used to assess the simulated landscapes in terms of habitat suitability for black grouse can be found in the supplementary information (S2 File).

165

To compare changes in landscape suitability after modelling with this base map, we converted the relative suitability predictions into a binary, presence/absence prediction for black grouse across the landscape using a habitat suitability threshold. The choice of threshold is extremely important [42] and in some cases can be a complex and case-specific choice. To keep these choices relatively simple for demonstration purposes, in this case, we tested three thresholds. These were low, medium and high suitability, corresponding to the first quartile, median and third quartile relative suitability scores, based on a habitat suitability model for the original habitat in 1994.

170

The most common metrics used to assess the predictive power of species distribution models require both presence and absence values for testing. Tests of predictive power were calculated from our presence-only dataset by producing a composite dataset consisting of our original presence points along

175

with 1000 background predictions from the binary presence/absence map similar to the map described above. This was created using a fourth threshold, the maximum sum of sensitivity and specificity. The MaxEnt model based on the original habitat was then tested using the area under the curve (AUC, ranges from 0 to 1) of the receiver operating characteristic plot (ROC [43]) as well as the true skill statistic (TSS, ranges from -1 to +1 [44]). In both cases, the closer the value is to 1, the more accurate the model.

185 **Scenario building and landscape change simulation**

The study area represents a highly dynamic working landscape. Analysis of satellite images from 1994 and 2008 show that while net gains or losses in each habitat type between 1994 and 2008 didn't exceed 3% for any major habitat type, the amount of land actually switching between habitat types was considerably larger (up to 17%; indicating both gains and losses in different parts of the landscape). During this period the black grouse population experienced a significant decline, followed by a recovery. Changes in habitat over this time and the impact on black grouse populations are explored in Geary *et al.* [45]. In order to choose appropriate future land use change scenarios for this proof of concept study, expert advice was sought. A questionnaire prompted respondents to choose their top five from ten potential scenarios, these coming from the authors' experience in upland research and the likelihood that they would take place during the next twenty years in Scotland. Surveys were received from ten professionals representing academic (30%), consultant (10%), conservation (30%), governmental (20%) and sporting interests (10%). The most likely future scenario was considered to be additional native forestry schemes resulting in more open-canopy forestry (Table 1). The second to fifth most likely scenarios were an increase in grouse moor, a decrease in grazing, an increase in grazing and an increase in closed-canopy forestry. Agreement among scenarios was generally good with each of the scenarios used in modelling chosen by over 50% of the experts (File S3).

Table 1 approximately here

205 For each scenario, land use change was simulated from 20 random starting locations (i.e. 20 random pixels from the 1994 classified image) using an iterative process which grows new pixels of habitat close to starting locations according to a probability of change (0.25; simply to determine the direction in which the patch grows) until they reached the new proportion of the habitat prescribed under the scenario. Starting locations refer to individual pixels around which simulated land use change was

210 centred and do not imply likely land use change at that point to be greater than at any other. The resulting habitat patches were non-uniform in shape and, due to the stochastic process, varied in size, but their combined area summed to the total area of new habitat. With the new habitat patches created, the proportion of each habitat type within 2 km of each grid cell in the study area was re-calculated. The 1994 MaxEnt model was then projected onto these novel landscapes and the percentage of the

215 study area predicted as suitable habitat for black grouse was calculated for each threshold suitability value. R scripts for the modelling functions as well as the scenarios are included in the supplementary information (File S4). Thirty new landscapes (i.e. 30 model runs) were generated for each scenario to allow average habitat suitability to be calculated. Results from these new scenarios could potentially have been influenced by changes in the spatial arrangement of patches rather than those produced by

220 the actual increase or decrease in the different habitat types. To explore this further we created a series of 'null' models for each scenario to compare situations with land use change against scenarios where the habitat is changed in the same way but with no net increase or decrease in the amount of each habitat type. Again, thirty new landscapes were generated for each of four null models (both increase in grazing and decrease in grazing have the same null model – no net change in grazing) and the 1994

225 MaxEnt model was then projected onto them, and the percentage of the study area predicted suitable for black grouse calculated.

The proportions of area predicted suitable under each of the five scenarios were compared using Kruskal-Wallis tests followed by pairwise Mann-Whitney U tests. In addition, each scenario was compared to its equivalent 'null' scenario using Mann-Whitney U tests. We examined in greater detail the impact of some closed-canopy forestry patches, a common land use change in the Scottish uplands, as they are thought to be detrimental for black grouse at a landscape scale [14] but changed land management around them could produce complex effects on habitat suitability. To do this, we present some examples of the effects of different placements of closed-canopy forest patches within the landscape. Supporting these examples, we present pre- and post-scenario landscape metrics such as number of closed-canopy forestry patches, mean patch sizes and total woodland edge computed using the SDMTools package [46]. Further to this, we investigated whether there was a linear (proportional) effect of adding increasing amounts of the given land use on habitat suitability. We varied the area affected by increased open-canopy forestry and reduced grazing, two land use changes which are thought to benefit black grouse [47], between 5% and 30% to identify any nonlinearities in benefit for black grouse.

While we use real data on lek presence for black grouse to illustrate our method, we do not include any demographic data in the study. As well as the suitability of habitat surrounding the lek we acknowledge that other considerations such as lek connectivity [48], the quality of habitat around surrounding leks [24], edge effects and source-sink dynamics are likely to also play a role in determining the overall quality of the landscape for this species. Although the model presented retains enough flexibility to incorporate these considerations, we have chosen to focus solely on habitat quality for the sake of simplicity while demonstrating the method.

250

Results

Comparisons across scenarios

The MaxEnt model, using a regularisation multiplier of 7 (Table 2), performed well in predicting black
255 grouse presence using the original environmental data (30 m pixels, AUC = 0.83, TSS = 0.64). All of
the scenarios produced outcomes that were significantly different from those of the ‘null’ scenarios (Fig
1) at each of the three thresholds (min $U = 0$, $n = 30$, max $P = 0.04$) except for the open canopy forestry
scenario which was not significantly different at the first and third quartile thresholds (min $U = 481$, n
 $= 30$, min $P = 0.06$) and the closed canopy forestry scenario which was not significantly different from
260 the null scenario at using the median threshold ($W = 390$, $n = 30$, $P = 0.38$). There were significant
differences between the amount of suitable habitat produced under the five scenarios at the first quartile
($\chi^2 = 132.6$, $df = 4$, $P < 0.001$; Fig 2a), median ($\chi^2 = 129.6$, $df = 4$, $P < 0.001$; Fig 2b) and third quartile
($\chi^2 = 120$, $df = 4$, $P < 0.001$; Fig 2c) thresholds. Across the three thresholds the most beneficial scenario
for black grouse in terms of increased habitat suitability across the landscape was the reduced grazing
265 scenario. It resulted in a significantly larger proportion of the landscape suitable for black grouse than
the next best scenario at the first quartile (increased open canopy forestry; $U = 900$, $n = 60$, $P < 0.001$)
and median (increased grouse moor; $U = 817$, $n = 60$, $P < 0.001$) thresholds. At the third quartile
threshold increasing grouse moor resulted in a significantly larger proportion of the landscape predicted
suitable for black grouse than the next best scenarios (reduced grazing; $U = 51$, $n = 60$, $p < 0.001$). The
270 lowest suitability at the first and third quartile thresholds was produced by the increased grazing
scenario. This predicted significantly smaller proportions of the landscape were suitable for black
grouse than the next lowest scenario at both thresholds (Q1 closed canopy forestry; $U = 607$, $n = 60$, P
 $= 0.02$, Q3 open canopy forestry; $U = 107$, $n = 60$, $P < 0.001$). Using the median threshold, the scenario
predicting the lowest suitability was increased closed canopy forestry. However, this did not predict

275 significantly lower suitability than the next lowest scenario (increased grazing; $U = 406$, $n = 60$, $P =$
0.52). Both the increased closed canopy forestry ($U = 2$, $n = 60$, $P < 0.001$) and increased grazing ($U =$
18, $n = 60$, $P < 0.001$) scenarios resulted in significantly lower suitability for black grouse than the next
worst scenario, increased open canopy forestry.

280 ### Table 2 approximately here ###

Fig 1 approximately here

Fig 2 approximately here

Effect of habitat placement

285 Changes to the area and position of closed-canopy forestry resulted in projections which could both
increase and decrease suitability for black grouse. Figs 3a-c represent simplified small sections of the
landscape showing 1994 habitat and altered habitat under an illustrative ‘one-off’ simulated land use
change, along with resultant differences in suitability. These figures are presented for reference only as
further analysis of this effect is beyond the scope of this paper. Below each map are selected landscape
290 metrics such as number of closed-canopy forestry patches, mean patch sizes and total woodland edge.
In most cases, the area covered by new forestry was much less suitable (Fig 3a), but in some situations,
the area immediately surrounding the new forest had improved suitability (Fig 3b). In other areas, a
more complex arrangement arose where habitat suitability had increased along some edges but not
others (Fig 3c).

295

Fig 3 approximately here

Effects of extent of land use change

Increases in the extent of open-canopy forestry resulted in a linear decrease in suitability (Fig 4a). For
300 the reduced grazing scenario, increasing the extent of land use change resulted in a nonlinear increase
in suitability (Fig 4b) which was disproportionately beneficial when 20% or more of grazed land was
converted (median 15% = 0.095 median 20% = 0.1; $U = 124$, $n = 100$, $P < 0.001$).

Fig 4 approximately here

Discussion

Fitting with our expectation, the most beneficial potential future scenario for the black grouse involved the conversion of grazed land to both grouse moor and open-canopy forestry and the most detrimental was an increase in grazing. This result reflected the detrimental impact of overgrazing on black grouse as well as their habitat requirements as woodland-edge species [49,50] although this demonstration of our method does not yet take into account fragmentation or the influence of habitat edge independent of habitat area. Consideration of changes to the amount of edge would be useful to include in future models, especially for species such as black grouse, as it would help to properly reflect patch shapes as well as sizes, especially in heavily-fragmented landscapes. Sheep grazing is a widespread feature of landscapes in the Scottish uplands and with added pressure from large red deer populations [51], overgrazing is a threat to many upland bird species [9]. Reduced grazing regimes, both in terms of fewer sheep and increased deer control, are likely to have widespread benefits for a number of species of conservation concern [52], including black grouse [50,53]. The numerical predictions of changes to the area suitable for black grouse should be regarded relative to the other scenarios rather than suggestions of the actual changes ‘on the ground’.

Under the increased closed-canopy forestry scenario, different model runs produced both increases and reductions in habitat suitability. Research has shown that canopy closure in areas of new plantation forestry has led to declines in black grouse populations [14]. Our example simulations using increased closed-canopy forestry (Fig 3) showed that the location of new patches of closed-canopy forestry was important in determining the resulting suitability for black grouse. That there is a landscape context to the effects of land use change on species is not surprising [54], especially in a species such as black

grouse, which is associated with habitat mosaics [12,13], and which can thrive in several rather
330 different combinations of land uses [13,24,55]. Indeed in our null models, we quantified the variability
in habitat suitability associated with moving habitat patches around rather than actually changing than
proportions of habitats. We then briefly explored some of the possible metrics, such as patch size and
edge effects, which may affect the impact of different patch placements on overall landscape suitability,
but it will take a more extensive modelling exercise to elucidate the multiple features that contribute to
335 landscape suitability. Our proposed method does, however, allow quantification of suitability changes
related to individual landscape changes, even if it does not identify the underlying causes of those
changes. At present, our scenarios are restricted to single land use changes (although an increase in one
land use results in loss of one or more other land uses) for ease of model demonstration. In reality,
changes may involve several land use types, as well as gradual changes in habitat characteristics
340 brought about natural habitat succession (e.g. canopy closure) and management interventions (e.g.
heather burning). Taken together with landscape context, this complexity has the potential to become
prohibitively computing-intensive to model accurately [56]. Again, such models make for powerful
ecological tools, and our illustration is a first practical step towards their realisation. As expected, the
interiors of closed-canopy plantations were consistently unsuitable [14,47], although the same was not
345 true for the areas surrounding new forest patches. Forest edges are a habitat feature preferred by black
grouse [49] and investigating the effect of patch shape on habitat suitability may be productive for land
managers.

At their most basic, increases or decreases in suitability would be proportionate to the amount of
350 habitat change (e.g. stone marten *Martes foina* and strawberry tree *Arbutus undoo* [57]) but more
complex, nonlinear relationships, perhaps depending on landscape structure or interacting effects (e.g.
models of invasive species [58]) or ecological thresholds (e.g. pine marten *Martes martes* in
fragmented forests [59]) are important in wildlife management. Increases in the area of grazed land

converted to grouse moor and open-canopy forestry was beneficial to black grouse, with a step change
355 at proportions above 15%. The spatial structure of the resulting landscapes is a likely cause for this
difference in response [58]. In one case, both moorland and grazed land were converted into open-
canopy forestry causing a homogenisation of the landscape which larger magnitudes of change
exacerbated. In contrast, reduced grazing resulted in a more heterogeneous landscape which contained
the mosaics attractive to black grouse [48,51]. Nonlinear responses to habitat management by species
360 can be related to edge effects [60]; as a forest edge species [49], black grouse might benefit from these
changes. Identifying these thresholds is important for species conservation as it highlights the potential
for rapid changes in abundance or distribution to occur [61,62].

Combined with knowledge of habitat management, scenario-led habitat suitability modelling could be
365 extremely useful both for agencies or consultants advising individual landowners on local
costs/benefits of land management changes [63], and as a basis for encouraging wider-scale changes
through appropriate policy or planning regulations [64]. Changes incorporated into the models could be
the results of changes in policy across landscapes, as we have demonstrated here, or specific spatial
changes of interest to land managers. Species of conservation concern which inhabit agricultural
370 landscapes could benefit greatly from changes to management practices (e.g. corn bunting *Miliaria
calandra* [65]) or from subsidies targeted to improving habitats (e.g. little bustard *Tetrax tetrax* [66] &
greater sage-grouse *Centrocercus urophasianus* [67]) both of which could be explored using landscape
simulation models. A natural progression would be to consider the position and shape of new land use
features as well as their location within the landscape. This would require an extension of the current
375 modelling framework and consideration of the computational requirements of creating realistic
representations of these complex situations. Landscape simulation models could be extended to work
on metrics such as species richness/diversity, or ecosystem function or services (e.g. abundance of

pollinators [68]). If the provision of ecosystem services or economic benefits is related to habitats or landscape structure then, by examining the effects of different land use change scenarios, a compromise
380 between economic activity and conservation can be achieved [6]. The inclusion of economic or sociological factors into landscape simulation models is another area of potential, with, for example, likelihood of land use change related to an ‘index of willingness’ which might be affected by differing financial rewards or levels of knowledge [69].

385 Black grouse in the Scottish uplands, like many species across the world, now exist largely within a working landscape [47]. Our models pointed to specific land use changes which are predicted to improve habitat for black grouse, finding a reduction in grazing, as field-based studies have [53], to be the most beneficial. Perhaps most importantly, our work indicates that it is not just the land use change itself that will determine whether species thrive or decline, but the extent of these changes and their
390 position in relation to other features in the landscape.

Acknowledgements

Thanks are due to the Perthshire Black Grouse Study Group for providing us with these data.

Thanks, also, to the experts surveyed for building future landscape scenarios.

References

1. Sala OE. Global Biodiversity Scenarios for the Year 2100. *Science*. 2000;287: 1770–1774. doi:10.1126/science.287.5459.1770
2. Fearnside PM. Deforestation in Brazilian Amazonia: History, Rates, and Consequences. *Conserv Biol*. 2005;19: 680–688.
3. Gillings S, Newson SE, Noble DG, Vickery JA. Winter availability of cereal stubbles attracts declining farmland birds and positively influences breeding population trends. *Proc R Soc B*. 2005;272: 733–9.
4. Gunilla E, Olsson A, Austrheim G, Grenne S ve N. Landscape change patterns in mountains, land use and environmental diversity, Mid-Norway 1960–1993. *Landsc Ecol*. 2000;15: 155–170.
5. Macchi L, Grau HR, Zelaya PV, Marinaro S. Trade-offs between land use intensity and avian biodiversity in the dry Chaco of Argentina: A tale of two gradients. *Agric Ecosyst Environ*. 2013;174: 11–20. doi:10.1016/j.agee.2013.04.011
6. Polasky S, Nelson E, Camm J, Csuti B, Fackler P, Lonsdorf E, et al. Where to put things? Spatial land management to sustain biodiversity and economic returns. *Biol Conserv*. 2008;141: 1505–1524. doi:10.1016/j.biocon.2008.03.022
7. Warren CR. *Managing Scotland’s Environment* [Internet]. Edinburgh: Edinburgh University Press; 2002. Available: <http://books.google.com/books?id=klpUoWztNL4C&pgis=1>
8. Scottish Government. *List of Rural Priorities Options* [Internet]. 2003 [cited 17 Sep 2014]. Available: <http://scotland.gov.uk/Topics/farmingrural/SRDP/RuralPriorities/Options>
9. Fuller RJ, Gough SJ. Changes in sheep numbers in Britain: implications for bird populations. *Biol Conserv*. 1999;91: 73–89.
10. Sim IMW, Gregory RD, Hancock MH, Brown AF. Recent changes in the abundance of British upland breeding birds: Capsule Breeding wader populations have more often shown declines than passerine populations during the last 10–20 years. *Bird Study*. 2005;52: 261–275. doi:10.1080/00063650509461399
11. Sim IMW, Eaton MA, Setchfield RP, Warren PK, Lindley P. Abundance of male Black Grouse *Tetrao tetrix* in Britain in 2005, and change since 1995–96. *Bird Study*. 2008;55: 304–313. doi:10.1080/00063650809461536
12. Baines D, Blake K, Calladine J. Reversing the decline: a review of some Black Grouse conservation projects in the United Kingdom. *Cah Ethologie Fondam Appl Anim Hum*. 2000;
13. Starling-Westerberg A. The habitat use and diet of Black Grouse *Tetrao tetrix* in the Pennine hills of northern England. *Bird Study*. 2001;48: 76–89.

- 430 14. Pearce-Higgins JW, Grant MC, Robinson MC, Haysom SL. The role of forest maturation in causing the decline of Black Grouse *Tetrao tetrix*. *Ibis*. 2007;149: 143–155. doi:10.1111/j.1474-919X.2006.00623.x
15. Austin M. Species distribution models and ecological theory: A critical assessment and some possible new approaches. *Ecol Model*. 2007;200: 1–19. doi:10.1016/j.ecolmodel.2006.07.005
16. Franklin J. Moving beyond static species distribution models in support of conservation biogeography. *Divers Distrib*. 2010;16: 321–330. doi:10.1111/j.1472-4642.2010.00641.x
- 435 17. Lahoz-Monfort JJ, Guillera-Arroita G, Milner-Gulland EJ, Young RP, Nicholson E. Satellite imagery as a single source of predictor variables for habitat suitability modelling: how Landsat can inform the conservation of a critically endangered lemur. *J Appl Ecol*. 2010;47: 1094–1102. doi:10.1111/j.1365-2664.2010.01854.x
- 440 18. Wilting A, Cord A, Hearn AJ, Hesse D, Mohamed A, Traeholdt C, et al. Modelling the species distribution of flat-headed cats (*Prionailurus planiceps*), an endangered South-East Asian small felid. *PloS One*. 2010;5: e9612. doi:10.1371/journal.pone.0009612
19. Knick ST, Hanser SE, Preston KL. Modeling ecological minimum requirements for distribution of greater sage-grouse leks: implications for population connectivity across their western range, U.S.A. *Ecol Evol*. 2013;3: 1539–1551. doi:10.1002/ece3.557
- 445 20. Hijmans RJ, Graham CH. The ability of climate envelope models to predict the effect of climate change on species distributions. *Glob Change Biol*. 2006;12: 2272–2281. doi:10.1111/j.1365-2486.2006.01256.x
21. Linnell J. Predators and people: conservation of large carnivores is possible at high human densities if management policy is favourable. *Anim Conserv*. 2001;4: 345–349.
- 450 22. Donald PF, Sanderson FJ, Burfield IJ, Bierman SM, Gregory RD, Waliczky Z. International Conservation Policy Delivers Benefits for Birds in Europe. *Science*. 2007;317: 810–813.
23. McRae BH, Schumaker NH, McKane RB, Busing RT, Solomon AM, Burdick CA. A multi-model framework for simulating wildlife population response to land-use and climate change. *Ecol Model*. 2008;219: 77–91. doi:10.1016/j.ecolmodel.2008.08.001
- 455 24. Geary M, Fielding AH, Marsden SJ. Designing mosaic landscapes for Black Grouse *Tetrao tetrix* using multi-scaled models. *Ibis*. 2013;155: 792–803. doi:10.1111/ibi.12090
25. Hancock M, Baines D, Gibbons D, Etheridge B, Shepherd M. Status of male Black Grouse *Tetrao tetrix* in Britain in 1995–96. *Bird Study*. 1999;46: 1–15.
- 460 26. Gilbert G, Gibbons DW, Evans J. *Bird Monitoring Methods: A Manual of Techniques for Key UK Species*. Pelagic Publishing Limited; 2011.

27. Cherrill AJ, McClean C, Lane A, Fuller RM. A comparison of land cover types in an ecological field survey in Northern England and a remotely sensed land cover map of Great Britain. *Biol Conserv.* 1995;71: 313–323. doi:10.1016/0006-3207(94)00053-S
28. Smith GM, Fuller RM, Sanderson JM, Hill RA, Thomson AG. Countryside Survey 2000
465 Module 7: land cover map 2000. Fifth quarterly progress report. 1999;
29. Biehl L, Landgrebe D. MultiSpec—a tool for multispectral–hyperspectral image data analysis. *Comput Geosci.* 2002;28: 1153–1159. doi:10.1016/S0098-3004(02)00033-X
30. Gowda CK. A feature reduction and unsupervised classification algorithm for multispectral data. *Pattern Recognit.* 1984;17: 667–676. doi:10.1016/0031-3203(84)90020-7
- 470 31. Fielding A. *Cluster And Classification Techniques for the Biosciences* [Internet]. Cambridge: Cambridge University Press; 2007. Available: <http://books.google.com/books?id=syHFxKjLSRgC&pgis=1>
32. Hijmans RJ, van Etten J. raster: Geographic analysis and modeling with raster data [Internet]. R package version 1.9-63.; 2012. Available: <http://cran.r-project.org/package=raster>
- 475 33. R Development Core Team. *R: A Language and Environment for Statistical Computing* [Internet]. Vienna, Austria; 2012. Available: <http://www.R-project.org/>
34. Cramp S. *Handbook of the Birds of Europe, the Middle East and North Africa: Hawks to bustards* [Internet]. Oxford: Oxford University Press; 1980. Available: <http://books.google.com/books?id=UDNftwAACAAJ&pgis=1>
- 480 35. Lindström J, Rintamäki PT, Storch I. Black grouse. - BWP Update. *J Birds West Palearct.* 1998;2: 173 – 191.
36. Phillips SJ, Anderson RP, Schapire RE. Maximum entropy modeling of species geographic distributions. *Ecol Model.* 2006;190: 231–259.
- 485 37. Hijmans RJ, Phillips S, Leathwick J, Elith J. dismo: Species distribution modeling [Internet]. R package version 0.7-17.; 2012. Available: <http://cran.r-project.org/package=dismo>
38. Elith J, Graham CH. Do they? How do they? WHY do they differ? On finding reasons for differing performances of species distribution models. *Ecography.* 2009;32: 66–77. doi:10.1111/j.1600-0587.2008.05505.x
- 490 39. Royle JA, Chandler RB, Yackulic C, Nichols JD. Likelihood analysis of species occurrence probability from presence-only data for modelling species distributions. *Methods Ecol Evol.* 2012;3: 545–554. doi:10.1111/j.2041-210X.2011.00182.x
40. Warren DL, Seifert SN. Ecological niche modeling in Maxent: the importance of model complexity and the performance of model selection criteria. *Ecol Appl.* 2011;21: 335–342.

41. Ficetola GF, Bonardi A, Mairota P, Leronni V, Padoa-Schioppa E. Predicting wild boar damages to croplands in a mosaic of agricultural and natural areas. *Curr Zool.* 2014;60: 170–179. 495
42. Jimenez-Valverde A, Lobo J. Threshold criteria for conversion of probability of species presence to either–or presence–absence. *Acta Oecologica.* 2007;31: 361–369. doi:10.1016/j.actao.2007.02.001
43. Fielding AH, Bell JF. A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environ Conserv.* 1997;24: 38–49. 500 doi:10.1017/S0376892997000088
44. Allouche O, Tsoar A, Kadmon R. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *J Appl Ecol.* 2006;43: 1223–1232. doi:10.1111/j.1365-2664.2006.01214.x
45. Geary M, Fielding AH, Marsden SJ. Both habitat change and local lek structure influence patterns of spatial loss and recovery in a black grouse population. *Popul Ecol.* In press; 505
46. VanDerWal J, Falconi L, Januchowski S, Shoo L, Storlie C. Species Distribution Modelling Tools: tools for processing data associated with species distribution modelling exercises. Package ‘SDMTools; 2012.
47. Watson A, Moss R. Grouse [Internet]. London: HarperCollins UK; 2008. Available: 510 <http://books.google.com/books?id=mFYAAQAACAAJ&pgis=1>
48. Geary M. The long-term viability of black grouse *Tetrao tetrix* populations in a changing upland landscape. PhD Thesis, PhD Thesis, Manchester Metropolitan University, Manchester, UK. 2012.
49. Baines D, Hudson PJ. The decline of Black Grouse in Scotland and northern England. *Bird 515 Study.* 1995;42: 122–131. doi:10.1080/00063659509477158
50. Calladine J, Baines D, Warren P. Effects of reduced grazing on population density and breeding success of black grouse in northern England. *J Appl Ecol.* 1996;39: 772–780.
51. Baines D. Seasonal differences in habitat selection by Black Grouse *Tetrao tetrix* in the northern Pennines, England. *Ibis.* 1994;136: 39–43.
52. Thompson DBA, Macdonald AJ, Marsden JH, Galbraith CA. Upland Heather Moorland in Great-Britain - a Review of International Importance, Vegetation Change and Some Objectives for Nature Conservation. *Biol Conserv.* 1995;71: 163–178. doi:10.1016/0006-3207(94)00043-P 520
53. Baines D. Seasonal variation in lek attendance and lekking behaviour by male black grouse *Tetrao tetrix*. *Ibis.* 1996;138: 177–180.
54. Schmiegelow FK, Mönkkönen M. Habitat Loss And Fragmentation In Dynamic Landscapes: Avian Perspectives From The Boreal Forest. *Ecol Appl.* 2002;12: 375–389. 525

55. Baines D, Warren P, Richardson M. Variations in the Vital Rates of Black Grouse *Tetrao tetrix* in the United Kingdom. *Wildl Biol.* 2007;13: 109–116. doi:10.2981/0909-6396(2007)13[109:VITVRO]2.0.CO;2
- 530 56. Kupfer JA. Landscape ecology and biogeography Rethinking landscape metrics in a post-FRAGSTATS landscape. *Prog Phys Geogr.* 2012;36: 400–420. doi:10.1177/0309133312439594
57. Cabezas-Díaz S, Lozano J, Mangas J, Virgós E. Spatial distribution models in a frugivorous carnivore, the stone marten (*Martes foina*): is the fleshy-fruit availability a useful predictor? *Anim Biol.* 2010;60: 423–436. doi:10.1163/157075610X523297
- 535 58. Didham RK, Tylianakis JM, Gemmell NJ, Rand TA, Ewers RM. Interactive effects of habitat modification and species invasion on native species decline. *Trends Ecol Evol.* 2007;22: 489–96. doi:10.1016/j.tree.2007.07.001
59. Caryl FM, Quine CP, Park KJ. Martens in the matrix: the importance of nonforested habitats for forest carnivores in fragmented landscapes. *J Mammal.* 2012;93: 464–474.
- 540 60. Hayward G. Response of Red-Backed Voles to Recent Patch Cutting in Subalpine Forest. *Conserv Biol.* 1999;13: 168–176. doi:10.1046/j.1523-1739.1999.97449.x
61. Ryall KL, Fahrig L. Response of predators to loss and fragmentation of prey habitat: a review of theory. *Ecology.* 2006;87: 1086–93.
62. Lindenmayer D, Hobbs RJ, Montague-Drake R, Alexandra J, Bennett A, Burgman M, et al. A checklist for ecological management of landscapes for conservation. *Ecol Lett.* 2008;11: 78–91. doi:10.1111/j.1461-0248.2007.01114.x
- 545 63. Ellis S, Wainwright D, Berney F, Bulman C, Bourn N. Landscape-scale conservation in practice: lessons from northern England, UK. *J Insect Conserv.* 2010;15: 69–81. doi:10.1007/s10841-010-9324-0
- 550 64. Hepinstall JA, Alberti M, Marzluff JM. Predicting land cover change and avian community responses in rapidly urbanizing environments. *Landsc Ecol.* 2008;23: 1257–1276. doi:10.1007/s10980-008-9296-6
65. Perkins AJ, Maggs HE, Watson A, Wilson JD. Adaptive management and targeting of agri-environment schemes does benefit biodiversity: a case study of the corn bunting *Emberiza calandra*. *J Appl Ecol.* 2011;48: 514–522. doi:10.1111/j.1365-2664.2011.01958.x
- 555 66. García J, Suárez-Seoane S, Miguélez D, Osborne PE, Zumalacárregui C. Spatial analysis of habitat quality in a fragmented population of little bustard (*Tetrax tetrax*): Implications for conservation. *Biol Conserv.* 2007;137: 45–56.

- 560 67. Aldridge CL, Nielsen SE, Beyer HL, Boyce MS, Connelly JW, Knick ST, et al. Range-wide patterns of greater sage-grouse persistence. *Divers Distrib*. 2008;14: 983–994. doi:10.1111/j.1472-4642.2008.00502.x
68. Polce C, Termansen M, Aguirre-Gutiérrez J, Boatman ND, Budge GE, Crowe A, et al. Species Distribution Models for Crop Pollination: A Modelling Framework Applied to Great Britain. *PLoS ONE*. 2013;8: e76308. doi:10.1371/journal.pone.0076308
- 565 69. Morris J, Mills J, Crawford IM. Promoting farmer uptake of agri-environment schemes: the Countryside Stewardship Arable Options Scheme. *Land Use Policy*. 2000;17: 241–254. doi:10.1016/S0264-8377(00)00021-1
70. Scottish Government. Abstract of Scottish Agricultural Statistics 1982 to 2013 [Internet]. 2014. Available: <http://www.scotland.gov.uk/Resource/0045/00455082.pdf>
- 570 71. Fraser of Allander Institute. *An Economic Study of Grouse Moors: An Update*. Game Conservancy Scottish Research Trust & Game Conservancy Ltd; 2010.
72. Pack B. *The road ahead for Scotland: Final report of the inquiry into future support for agriculture in Scotland*. Scottish Government; 2010.

575 **Figure Legends**

Fig 1. The proportion of the study area predicted suitable for black grouse under each of the land use change scenarios and their ‘null’ equivalent. Boxplots of the proportion of the study area predicted as suitable for black grouse using the median threshold (0.58) under each land use change scenario along with its ‘null’ equivalent where the configuration of land uses was changed but not the proportion of each land use type. The dotted line indicates the proportion of the study area predicted suitable using the original habitat.

580

Fig 2. The proportion of the study area predicted suitable for black grouse under each of the land use change scenarios. Boxplots of the proportion of the study area predicted as suitable for black grouse using the A) lower quartile (0.48), B) median (0.58); and, C) upper quartile (0.65) thresholds

585

using 30 m pixels. The dotted line indicates the proportion of the study area predicted suitable using the original habitat at each threshold.

590 **Fig 3. Examples of different outcomes resulting from closed canopy forestry placement within the landscape.** Examples of new closed-canopy forestry placement resulting in A) a reduction in habitat suitability within the new forest, B) a decrease in habitat suitability within the new forest coupled with an increase in habitat suitability around the new forest edge, and, C) a reduction of habitat suitability within the new forest with an increase in habitat suitability around some of the forest edge. Text under
595 individual habitat examples shows NP – Number of forestry patches, TE – Total forest edge (km), TA – total forest area (km²) and MA – mean forest area (km²).

Fig 4. The proportion of the study area predicted suitable for black grouse under different levels of habitat change. Boxplots representing the proportion of the study area predicted present under
600 scenarios different levels of habitat change (5-30%) using the median threshold (0.58) for A) the increased open-canopy forestry scenario, and, B) the reduced grazing scenario. The dotted line represents the proportion of the study area predicted suitable using the original habitat.

Supporting Information Captions

605 **S1 File. Validation of the habitat classification.** Validation of our own habitat classification with reference to the UK national land use classification.

S2 File. Modelling black grouse habitat suitability using MaxEnt. A description on the MaxEnt model used to assess habitat suitability under each scenario.

S3 File. Scenarios used in landscape simulation modelling. A description of the potential scenarios
610 indicating the scenarios chosen by experts as the most likely and the agreement between choices.

S4 File. R code for landscape simulation modelling. R code to run the models under the stated scenarios.

Tables

Table 1. Scenarios used in landscape simulation modelling. Scenarios used in landscape simulation

615 modelling, along with how the landscape is changed under each. Scenarios were chosen by upland experts as most likely to occur in the Scottish uplands from ten candidate scenarios. The ranked likelihood of each scenario, as decided by respondents to the questionnaire is also presented.

SCENARIO	RANK	CHANGES TO THE LANDSCAPE
Increased open-canopy forestry	1	Woodland creation grants larger than those for plantation forestry are available for the planting of native forestry under the (SRDP; http://www.scotland.gov.uk/Topics/farmingrural/SRDP/RuralPriorities/Options/). This will be reflected by a conversion of 7% of grazed land and 3% of grouse moor to open-canopy/mixed woodland. Grazed land was considered more likely to be converted to woodland than grouse moor.
Reduced grazing	2	Since 1982, sheep numbers in Scotland have decreased by 34% [70]. Grants encouraging a reduction in grazing are currently available through the Scottish Rural Development Programme (SRDP; http://www.scotland.gov.uk/Topics/farmingrural/SRDP/RuralPriorities/Options/). Under this scenario, a continued reduction in upland grazing will be reflected by a 10% reduction in grazed land 7% of which will be converted into moorland and 3% into open-canopy/mixed forestry.
Increased grouse moor	3	Economic analysis of the grouse shooting industry by the Fraser of Allander Institute [71] showed increased profitability in managed grouse shoots in 2010 and suggested this may lead to an increase in the area of moorland used for shooting. This will be reflected by a conversion of 5% of grazed land and 5% of open-canopy woodland to grouse moor.
Increased grazing	4	Both the Tenant farmers association and the Pack enquiry [72] have suggested a return to headage payments for upland farmers. In the past, this has resulted in an

increase in the number of sheep. This change will be reflected by converting 10% of moorland to grazed land.

Increased 5 Woodland creation grants available under the (SRDP;
closed-canopy <http://www.scotland.gov.uk/Topics/farmingrural/SRDP/RuralPriorities/Options/>)
forestry offer a financial incentive for the creation of plantation forestry. This will be reflected in the conversion of 5% of grouse moor and 5% of grazed land to plantation forestry.

620 **Table 2. Model selection comparing regularisation parameters (β) in candidate MaxEnt models.**

Regularisation parameter (β)	Log Likelihood	Parameters	AICc score	Δ AICc
1	-946.569	37	2052.35	111.38
2	-953.742	19	1960.39	19.41
3	-958.848	16	1959.77	18.80
5	-966.734	13	1965.85	24.88
7	-958.653	10	1940.97	0
9	-978.316	8	1974.95	33.98
10	-980.113	9	1981.18	40.20
11	-983.429	9	1987.81	46.84
13	-985.944	8	1990.21	49.24
15	-987.984	6	1989.28	48.31
17	-988.778	6	1990.87	49.90
19	-990.25	6	1993.81	52.84