

The Psychological Impact of Cancer (PIC) Scale: development and comparative psychometric testing against the Mini-MAC© Scale in UK and Australian cancer survivors.

Item Type	Article
Authors	Hulbert-Williams, Lee;Whelen, Liz;Mulcare, Hunter
Citation	Hubert-Williams, H. J., Hubert-Williams, L., Whelen, L. & Mulcare, H. (2019). The Psychological Impact of Cancer (PIC) Scale: development and comparative psychometric testing against the Mini-MAC© Scale in UK and Australian cancer survivors. Journal of Psychosocial Oncology Research and Practice, 1(2), e8.
DOI	10.1097/OR9.0000000000000008
Publisher	Wolters Kluwer Health
Journal	Journal of Psychosocial Oncology Research and Practice
Rights	Attribution-NonCommercial-NoDerivatives 4.0 International
Download date	2026-05-17 12:48:05
Item License	http://creativecommons.org/licenses/by-nc-nd/4.0/
Link to Item	http://hdl.handle.net/10034/622519

The Psychological Impact of Cancer (PIC) Scale: development and comparative psychometric testing against the Mini-MAC© Scale in UK and Australian cancer survivors.

Nicholas J Hulbert-Williams PhD CPsychol *
Centre for Contextual Behavioural Science, School of Psychology,
University of Chester, UK

Lee Hulbert-Williams PhD CPsychol
Centre for Contextual Behavioural Science, School of Psychology,
University of Chester, UK

Liz Whelen, PhD
School of Psychology, University of Chester, UK

Hunter Mulcare, D.Psych (Health)
Department of Psychology, Western Health, Victoria, Australia

*Full address for correspondence:

Prof Nick Hulbert-Williams, Centre for Contextual Behavioural Science, School of Psychology, University of Chester, Parkgate Road, Chester, UK, CH1 4BJ (n.hulbertwilliams@chester.ac.uk).

FUNDING: This work was supported by the North Wales Research Committee (Study 1) and The University of Chester (Study 2; Sample 1).

DISCLOSURE: The authors declare no conflicts of interest.

FOOTNOTES

AUTHORSHIP: NHW, LHW and HM were responsible for the design of this study. NHW and HM were responsible for original data collection, prior to this secondary analysis. All authors contributed to analysis. NHW was primary author of the manuscript. All authors contributed to manuscript re-drafting and approved the final manuscript.

EMAILS:

NHW: n.hulbertwilliams@chester.ac.uk
LHW: l.hulbertwilliams@chester.ac.uk
LW: e.whelen@chester.ac.uk
HM: huntermulcare@wh.org.au

TEXT: None

The Psychological Impact of Cancer (PIC) Scale: development and comparative psychometric testing against the Mini-MAC© Scale in UK and

Australian cancer survivors.

ABBREVIATIONS

AGFI	Adjusted Goodness of Fit Index
CFA	Confirmatory Factor Analysis
ECVI	Expected Cross Validation Index
MAC	Mental Adjustment to Cancer (Scale)
PIC	Psychological Impact of Cancer (Scale)
PNFI	Parsimony-adjusted Normal Fit Index
RMSEA	Root Mean Square Error of Approximation
SEM	Structural Equation Modelling

ABSTRACT

Background: Clinicians and researchers make considerable use of both the Mental Adjustment to Cancer (MAC) Scale, and the shorter Mini-MAC, to measure psychological adjustment in cancer patients. The length of the scale is problematic when used clinically, and its psychometric properties have been criticized. This paper presents two studies leading to the development of a novel scale the Psychological Impact of Cancer (PIC) Scale using items drawn from the MAC.

Methods: Study 1 used standard item-reduction techniques to shorten the Mini-MAC in a sample of 160 cancer patients of mixed diagnosis, recruited an average 46 days post-diagnosis. This resulted in a 12-item scale with a four-factor structure, similar to that derived from a 2012 re-analysis of the Mini-MAC. Study 2 presents confirmatory factor analysis (CFA) of this new measure and tests its construct validity and test-retest reliability in a sample of 183 mixed cancer survivors.

Results: This study indicated that the shorter scale performed well on CFA indicators (RMSEA= .083; ECVI= .923; PNFI= .604; AGFI .857) and tests of internal consistency (all $>.623$); and comparable concurrent validity with longer versions. The four factors were labeled cognitive distress, cognitive avoidance, emotional distress and fighting spirit.

Conclusions: Given its shorter length and acceptable psychometrics, the PIC offers a useful clinical and research tool to assess the psychological impact of cancer. Psychometric properties of one subscale (fighting spirit) remain poor, but no worse than in the original scale; directions for further development of the scale are described.

KEYWORDS

Cancer, Adjustment, Distress, Quality of Life, Measurement, Psychometric Validation, Structural Equation Modelling, Factor Structure, Validity, Reliability,

BACKGROUND

A cancer diagnosis often results in distress [1,2]. Indeed, distress is such a pressing concern that recommendations and clinical practice guidance have been developed to advise oncology services in the screening, management and intervention to ensure that distress is recognised and treated as efficiently as possible [3]. Reliable and valid psychometric tools are needed to enable screening, to facilitate psychosocial treatment, and for research on the psychological impact of cancer [4].

One of the most commonly used measures in psychosocial oncology is the Mental Adjustment to Cancer (MAC) Scale [5] (which is now also available in 20 languages (see www.ipos-society.org) and the shorter Mini-MAC [6]. Though there is some debate as to the precise nature of the constructs measured (i.e. whether it assess coping behaviours, cognitions, or some other underlying individual difference), the various sub-scales (of both the MAC and Mini-MAC) correlate with numerous psychosocial outcome measures [7-9], and are predictive of other important factors associated with illness such as information needs and information-seeking behaviours [10,11].

Watson and Homewood [12] published one of the largest re-analyses of the full MAC scale to date (n=1255), suggesting two higher order factors (positive and negative adjustment) as a more useful scoring algorithm compared with the 6 subscales in the original. Though there are some foreign-language validations of the shorter Mini-MAC – for example into Chinese, Greek, Italian, Korean, Norwegian, Portuguese, and Spanish [13-19] – which demonstrate overall reasonable psychometric performance, there are few robust psychometric analyses of the original English-language format [20]. Previous published applications of the scale have suggested that poor internal consistency of some sub-scales necessitates further scale development [15,17,20]. Indeed, a number of authors suggest that when using the foreign-language translations of the scale, items perform psychometrically better when grouped according to alternative factor structures [13-20].

A recently published factor analysis of the English-language version of the Mini-MAC [21] similarly suggested an alternative factor structure. Removal of five items yielded equivalent internal consistency and test-retest reliability, but markedly improved convergent validity with anxiety, depression and quality of life. Furthermore, this analysis suggested that the four scales—cognitive distress, emotional distress, cognitive avoidance, and fighting spirit—were more parsimonious and were more easily interpreted in the context of modern psychological interventions. From a practical perspective, however, a 25-item scale is still lengthy and participant burden may reduce utility as both a clinical

and research tool. Whilst there is a need to confirm this alternative factor structure, it is equally important that attempts are made to create as brief a measure as possible for maximised use in clinical practice.

AIMS

This paper comprises results from two studies. The first aimed to create a reliable, but much briefer measure of the psychological impact of cancer using items taken from the Mini-MAC. This was based on our previous work which had resulted in a revised four-factor structure of the measure [21]. The second study aimed to compare the psychometric properties of this newer, brief measure (The Psychological Impact of Cancer Scale) against the previously validated Mini-MAC scoring algorithms (both original 5 factor and revised 4 factor). This study pooled data from UK and Australian cancer patients for secondary statistical analysis.

STUDY 1

METHOD

This study was a secondary analysis of a previously published dataset in which an alternative four-factor structure for the Mini-MAC was proposed [*reference excluded for anonymous peer-review*]. The same data were re-analysed to create a new, brief measure of the psychological impact of cancer.

Recruitment and data collection

Following university and health service ethical approval, participants were recruited (by postal invitation from their cancer nurse specialist) from three clinical sites in North Wales, UK. Patients were excluded where life expectancy was less than six months, cancer diagnosis was recurrent, English language comprehension was poor, or where they were considered too distressed (as assessed by clinical care teams). Of 902 patients diagnosed, 554 (51%) met study entry criteria; of these, 160 participated (response rate = 35%). The sample comprised colorectal ($n=44$), breast ($n=69$), prostate ($n=19$) and lung ($n=28$) cancer patients recruited a mean of 46 days ($SD=24.8$) post diagnosis. The sample included both male ($n=63$) and female ($n=97$) participants with a mean age of 64 years ($SD=9.97$).

Informed consent was obtained using written information sheets. The Clinical Nurse Specialist (CNS) assessed patient eligibility using a standardised eligibility flowchart and subsequently distributed information packs, consent forms and questionnaires via the post. Questionnaire packs comprised a brief socio-demographic questionnaire and the Mini-MAC [6].

Analysis

Using [*ref removed for anonymity*] revised four-factor structure of the Mini-MAC as a starting point, item-reduction techniques were used to create brief versions of each sub-scale [22]. Our goal was to reduce the number of items in each subscale without causing too much deterioration in psychometric properties. Simulation studies show that it is difficult to cross-validate factor structures when fewer than three items are included in each sub-scale [23], therefore 3 items were chosen to be the lower limit for each subscale. As the *fighting spirit*

scale contained only three items, all were retained.

Some psychometricians recommend the use of Cronbach's alpha as a reference criterion during item reduction [22]. Robust criticisms have been levelled against coefficient alpha, in particular because it is biased by the number of items in the scale and must, therefore, be interpreted in the context of scale length [24]. We therefore chose two pragmatic criteria. Items were removed so long as the mean inter-item correlation was not reduced by more than .05 and a reduction no greater than .1 was seen in Cronbach's alpha.

Analyses for both studies were conducted using SPSS v19 with AMOS add-on (IBM Corporation, Somers, NY USA).

RESULTS

Through item reduction, a 12-item scale was developed:

Cognitive Distress (9 items deleted). Three items cross-loaded with other factors and so were eliminated. Though this negatively impacted Cronbach's alpha initially, this was improved through the elimination of six further items. The remaining three items had a Cronbach's alpha of .733 (lower than the fuller scale, but still adequate) and mean inter-item of correlation of $r=.493$ (substantially improved from the full scale).

Cognitive Avoidance (2 items deleted). Marginal improvement in Cronbach's alpha was achieved through elimination of one item. Removing one further item caused a substantial improvement in mean inter-item correlation ($r=.452$), but caused a slight dip in Cronbach's alpha (.709).

Emotional Distress (2 items deleted). This was initially the weakest performing subscale. A substantially improved Cronbach's alpha was achieved by eliminating one item, though this was attenuated slightly by the removal of a second item (.830). Overall, this resulted in an improved mean inter-item correlation for the scale ($r=.619$).

Fighting Spirit. This subscale was not submitted to item-reduction as it contained only three items.

With the exception of the *fighting spirit* sub-scale, this brief measure compares favourably with the 25-item version of the Mini-MAC from which it was developed [reference excluded for anonymous peer review]. Only for two sub-scales is Cronbach's alpha smaller, and even so it surpasses the .70 cut-off for adequacy recommended by Kline [25]. Mean inter-item correlation is substantially improved for all sub-scales. Table 1 presents items included in the PIC Scale. As the measure deviated considerably from the full Mini-MAC, permission was given by the original authors to publish this as a new measure: the Psychological Impact of Cancer (PIC) Scale.

INSERT TABLE 1 ABOUT HERE

STUDY 2

METHOD

Relevant data from three pre-existing clinical studies [*references excluded for anonymous peer review*] were combined to produce a sample of sufficient size to power Structural Equation Modelling (SEM). Bentler & Chou [26] recommend 5 participants per observed variable as the lower limit for confirmatory factor analysis.

Sample and procedure of data collection

An outline of the sampling strategy and procedure of data collection for each of the three samples is presented below (see also table 2). Ethical approval was provided and participants were recruited using a fully informed procedure.

Sample 1: A cross-sectional sample of adult cancer patients (breast, colorectal, lung and prostate) between two and twelve months post-diagnosis, were recruited by postal invitation from a regional cancer centre in the UK (distributed directly by hospital cancer services). From 500 patients approached, 130 returned completed questionnaires (26% response rate); these were of mixed gender and had a mean age of 63.7 years (SD=11.6). A small proportion (14.3%) had been diagnosed with palliative illness.

Sample 2: Australian adults with primary diagnosis of lung cancer (mainly non-small cell and/or late stage disease) were recruited at first radiation oncology appointment. Eligible patients were identified in conjunction with the treating oncologists. Consenting patients were provided with study materials (in person, during routine clinical appointments) and followed up by phone if they had not mailed the questionnaire back within two weeks. Of 125 eligible patients, 73 returned completed questionnaires (58% response rate). The sample were mixed gender and had a mean age of 65.7 years (SD=12.0). Participants were recruited at an early time-point from diagnosis: over 50% were yet to commence treatment (42.5% with palliative intent). Fifty-nine participants also completed the Mini-MAC at a second time-point, one month later.

Sample 3: Fifteen Australian adults with a primary diagnosis of lung cancer were recruited into a pilot study prior to data collection for sample 2; these were recruited in the same manner as sample 2 though they were not invited to participate in a follow-up time 2 questionnaire. This sample was of mixed gender and had a mean age 70.6 years (SD=11.1).

From each dataset we extracted Mini-MAC responses from all participants who had completed the questionnaire at the point of recruitment. Additionally, equivalent clinical and demographic variables were extracted for all three samples. From samples 1 and 2 we were able to match data on anxiety and depression (using the Hospital Anxiety and Depression Scales (HADS [27])). Sample 1 also included data on Quality of Life (FACT-G [28]), Perceived Stress (PSS-10 [29]) and Benefit Finding (Perceived Benefit Finding for Cancer Scale [30]).

Analysis

Where less than 10% of data on the Mini-MAC were missing, this was replaced by expectation-maximization method; participants with higher proportions of missing data were excluded listwise. Three scoring algorithms were compared using confirmatory factor analysis: the original five-factor Mini-MAC model (Hopelessness/Helplessness, Anxious Preoccupation, Cognitive Avoidance, Fighting Spirit, Fatalism) [6]; the revised four-factor Mini-MAC model (Cognitive Distress, Cognitive Avoidance, Fighting Spirit, Emotional Distress) [21]; and, the briefer PIC model derived in Study 1.

A Confirmatory Factor Analysis (CFA) was conducted to compare the level of fit for each model. As our aim was to obtain the most parsimonious model capable of explaining the observed associations we report Parsimonious Normed Fit Index (PNFI) and Adjusted Goodness of Fit Index (AGFI), which are adjusted for parsimony [28]. We further report Expected Cross Validation Index (ECVI) and Root Mean Squared Error of Approximation (RMSEA), which involve estimation of population criteria, and thus better estimate the extent to which the tested models will be confirmed in other samples [31].

Cronbach's alpha and inter-item correlation were calculated for each factor to allow for comparison of internal consistency. Test-retest reliability was analysed by correlating baseline and one-month follow-up responses for sample two participants only. Convergent validity was analysed by correlating Mini-MAC scores with self-reported anxiety, depression, perceived stress, perceived benefit finding and quality of life: based on the broad psychosocial oncology literature, scores more indicative of positive adjustment and psychological well-being were hypothesised to correlate with higher quality of life and benefit finding; and lower stress, anxiety and depression.

RESULTS

Sample description

After exclusion of missing data, 183 participants remained in the study. There was an almost equal gender split. The sample was biased towards a greater proportion of lung cancer patients (55%). There was a good spread of participants at each disease stage and a quarter of the sample were being treated with palliative intent. Mean age was 64.8 years (SD=12; range 32 to 89), and mean time from diagnosis to recruitment was 163 days (SD=115; range 30 to 577) (see table 2).

INSERT TABLE 2 ABOUT HERE.

Demographic and clinical characteristics were compared to ensure that these samples were suitable to be combined using Chi-Square and ANOVA tests. Sample 1 included proportionally more females (mainly breast cancer patients) whilst Sample 2 recruited more males. There were no significant differences in participants' age between samples. Sample 2 had a higher proportion of participants being treated with palliative intent, and, the two lung cancer samples (samples 2 and 3) had higher proportions at more advanced disease stage ($\chi^2=66.202$, $df=2$, $p<.01$). Mean time from diagnosis was significantly different between samples; Sample 1 participants were recruited, on average,

later after diagnosis, though Sample 2 included a wider range of time interval. Whilst these differences indicate heterogeneity within our pooled sample, we are encouraged that the result is greater representativeness of broader cancer populations.

Confirmatory Factor Analysis

The results of the CFA and tests of internal reliability for each sub-scale are shown in table 3.

INSERT TABLE 3 ABOUT HERE

Confirmatory Factor Analysis. For a well-fitting model, both RMSEA and ECVI (related statistics which estimate population parameters) should be small. For RMSEA, 0.05 is usually taken as indicating good fit, whilst 0.10 is suggestive of adequate fit: no similar cut-offs are well accepted for ECVI, though the general interpretation is that a smaller value indicates a better fit [32]. Both the five-factor Mini-MAC and PIC models demonstrate comparable RMSEA, with the four-factor Mini-MAC performing less well. ECVI is considerably better for the PIC model; much better than both versions of the Mini-MAC. PNFI and AGFI should be closer to 1.00 for a well-fitting model; again, the PIC model is considerably better on both statistics than either of the longer scoring formats.

Internal Consistency. Both Cronbach's alpha and mean inter-item correlation were examined as indicators of internal consistency as Cronbach's alpha is known to be affected by the length of a scale [24].

Highest Cronbach's alpha emerged for the five-factor Mini-MAC, though this is not necessarily unexpected given the larger number of items in each subscale [33]. The four-factor Mini-MAC model scores equally high for *cognitive distress* and *cognitive avoidance*, but *fighting spirit* falls slightly below the .7 level usually considered good; *emotional distress* is considerably lower. The PIC performed slightly worse for *cognitive distress* but both *cognitive avoidance* and *emotional distress* demonstrate higher internal reliability than in the four-factor Mini-MAC model. Though these are lower than the five-factor Mini-MAC model, this is

more likely a product of reduced item numbers rather than poorer performance of the scales *per se*. Again, we would draw readers' attention to Streiner [34], Cortina [24] and Cronbach's original exposition of the use of alpha [35] to see why alpha tends to lend an appearance of acceptability to long scales, even where it is constructed from items which in truth share little variance. We suspect the length of the original Mini-MAC scale may be masking its low internal consistency.

Examining mean inter-item correlation, the five-factor Mini-MAC structure performs well with the exception of the *fatalism* sub-scale. In all respects the PIC statistically out-performs the four-factor Mini-MAC from which it was derived, in one case achieving a higher mean inter-item correlation than also any of the five-factor Mini-MAC sub-scales.

Test-retest reliability and construct validity

Data from Sample 2 included a one-month follow-up of data collection; these data were used to calculate test-retest reliability. Construct validity is assessed by testing correlation with a range of commonly-used psychosocial outcome variables with data taken from Samples 1 and 3 (see table 4).

INSERT TABLE 4 ABOUT HERE

Time-lagged Spearman's correlation coefficients were significant for all sub-scales, indicating good test-retest agreement, though effect sizes varied; for test-retest reliability, correlations greater than .7 are considered good, but those less than .6 are considered weak and should be regarded with caution [25]. *Anxious preoccupation* ($r=.789$) and *fatalism* ($r=.889$) within the original five-factor Mini-MAC indicated highest test-retest reliability; these factors did not emerge in the four-factor models and so direct comparison is not possible. The five-factor structure also presents the scale with poorest test-retest reliability statistic—that for *hopelessness/helplessness* ($r=.502$)—though it may be that this construct is more sensitive to mood changes over time, and so retest stability may not be an appropriate psychometric dimension. The PIC model presents the most consistent set of test-retest reliabilities; in creating this brief

measure, three sub-scales remain largely equivalent with regard to test-retest reliability, but *cognitive distress* decreased from $r=.663$ to $.504$. Though this falls below the $.6$ level of acceptability this may not be a true reflection of weakness within the measure, but simply that the variable, like *hopelessness/helplessness* itself is not stable over time.

Regarding concurrent validity, the PIC subscales are associated with many of the same psychological outcomes as the original Mini-MAC, and indeed the four-factor Mini-MAC. Benefit finding positively correlated with *fighting spirit* in each model. Each measure has at least one subscale that correlates positively with Quality of Life, though these subscales vary in construction due to the different factor solutions offered for each scoring algorithm. The PIC retains subscales which correlate with anxiety, depression and perceived stress, though the strength of these relationships are slightly attenuated as one might expect when working with shorter psychometrics optimized for clinical utility. Based on these data, neither the Mini-MAC (five or four factor) or the PIC stands out as superior. What is perhaps more pertinent is that the pattern of significant correlation is largely identical for the full five-factor and shorter four-factor version of the Mini-MAC and the much briefer PIC scale: concurrent validity was not substantially affected by the item-reduction process.

DISCUSSION

This paper details the development and psychometric validation of a short measure of the psychological impact of cancer: the PIC Scale. This was developed using items previously forming the Mini-Mental Adjustment to Cancer Scale [6]. Although a previous psychometric validation study of the Mini-MAC [21] achieved somewhat improved psychometric properties, it was still a lengthy scale for clinical use (25 items) and confirmatory factor analysis was required. This study addressed both of these issues. Firstly by resulting in the construction of a briefer (12-item) measure of the psychological impact of cancer with a simplified four factor structure (fighting spirit, cognitive distress, cognitive avoidance and emotional distress) as recommended by [reference removed for anonymous peer review] [21]. Secondly it provided confirmatory psychometric analysis of the revised four-factor Mini-MAC against the original scoring structure of the Mini-MAC.

Study 1 involved item-reduction analysis to create the PIC from the revised, four-factor Mini-MAC. The original *fighting spirit* sub-scale only contained three items and so was excluded from item-reduction development. Lower Cronbach's alphas for *cognitive distress*, *cognitive avoidance* and *emotional distress* resulted from item reduction, however, these are still within acceptable ranges. The observed reduction in Cronbach's alpha is likely a bi-product of including fewer items than a true indication of internal consistency [24,35]. Analysis of mean inter-item correlation, an alternative indicator of internal consistency, supported this interpretation as item-reduction resulted in substantially improved mean inter-item correlation for all three sub-scales.

Pooled data from three studies assessing the Mini-MAC were then analysed to compare the psychometric performance of the PIC against both the 25-item four-factor version, and the original 29-item five-factor version of the Mini-MAC. The results indicate that the PIC performs somewhat better than its longer counterparts on both confirmatory factor analysis indicators, and internal consistency. Convergent validity of the PIC is comparable with that of both scoring versions of the Mini-MAC here reported. These results indicate that whilst none of the scoring models are superior, the revised four-factor Mini-

MAC, and more-so the PIC, performs no poorer than the original five-factor scoring model.

Though test-retest reliability was slightly worse for the PIC, there is questionable utility of such a statistic for this particular measure. Psychological adjustment to cancer is complex [13,36] and previous data shows that psychological aspects are in constant flux through this period of stressor adaptation [37]. It may be that lower test-retest reliability reflects temporal instability of underlying constructs rather than measurement error.

It was previously suggested that the four-factor version of the Mini-MAC offered a conceptually improved sub-scale model, where re-organised items group into variables that are more theoretically meaningful [21]. On the basis of the data presented in this paper we maintain this position and concur that the four factors may hold more meaningful implementation as screening or outcome assessments in both the clinical setting and research use. Factor analysis did not offer clear confirmatory data for this 25-item model. Rather, these data suggest that the PIC, which contains these same four factors, offers a psychometrically improved measure than its longer counter-part. Indeed, on most criteria assessed, the PIC outperformed the original five-factor scoring model of the Mini-MAC. We would encourage those seeking a measure of psychological adjustment to consider this version for their own use. In a move towards shorter and more parsimonious assessments in the psychosocial oncology setting, the PIC offers a valid and reliable assessment of various sub-types of psychological distress and adaptation observed within clinical settings.

Whilst this paper offers a definitive comparison on the statistical utility of the measure, there are still areas for further development. Of the four components included, the *fighting spirit* sub-scale is the most weakly performing. As the only sub-scale phrased in terms of positive adjustment to cancer, the sub-scale seems to offer a unique construct not addressed by the remaining items. However, in both [reference removed for anonymous peer review] [21] previous paper, and this latest analysis, the reliability and validity data suggest scope for improvement. Further work exploring the importance of this construct within

overall adjustment processes would be useful, as would a careful analysis of whether the performance of the sub-scale could be improved through revised wording of the question items. A more generic wording moving away from a focus on fighting spirit and towards a broader measure of positive adaptation and acceptance may be more useful and psychometrically valid [38].

These analyses suggest room for improvement with regard to convergent validity. It is important to consider, however, that with a few exceptions [39], attempts to validate the various versions of the Mini-MAC use other self-report measures of psychosocial outcome as the marker for convergent validity. Whilst we are not suggesting this is an inappropriate technique, we wish to highlight that such measures are themselves proxy indicators of well-being and clinically-observable co-morbidities. It may, therefore, be appropriate to consider exploring convergent validity of this measure within a clinical setting where scores (and change over time) can be analysed alongside real-world referral to, and treatment outcomes from, psychological services within the clinical oncology setting.

Though this study represents a useful advance on the science of measuring the psychological impact of cancer, it is not without flaws. As is often the case with survey research in psychosocial oncology [40], our response rates were somewhat disappointing; this is probably consequential of the impersonalized nature of postal recruitment in two out of four included samples in this paper. It is likely that with these lower response rates to the research invitation, the samples were relatively self-selecting and we hypothesise that this may have excluded those who were highly distressed who may perceive research participation as too additionally burdensome [41], a factor that we know to be problematic in broader clinical trial research [42]. Whilst this is not a problem from a psychometrics point of view (there was still sufficient variance in scores to conduct the type of analysis intended), it would be interesting to explore whether the factor structure, and especially convergent validity, alter in a sample more highly distressed. Improving on previous validation work, we were able to gain a more varied sample of participants, including improved gender split, and higher proportions of more physically-unwell cancer patients. This

sample has the added value of including data from multiple countries, and therefore varied health-care settings (UK and Australia), though in non-English speaking countries further validation is warranted.

In conclusion: From the original Mini-MAC items, a new, brief measure of the Psychological Impact of Cancer (PIC) was created. This initial analysis demonstrates that the PIC has good psychometric properties whilst being brief and conceptually easy to understand. Whilst confirmatory psychometric analysis of this new scale should be undertaken when data permit, the results from this study should provide some level of confidence for clinicians and researchers alike to use this tool to assess an individual's psychological response to cancer diagnosis and treatment.

ACKNOWLEDGEMENTS

We are very grateful to the authors of the Mini-MAC© for giving us copyright permission to use items from the scale as a basis for developing this brief measure of psychological impact of cancer. We are thankful to Maggie Watson for commenting on an early draft of this work. This work was funded by the North Wales Research Committee (Study 1) and The University of Chester (Study 2; Sample 1).

REFERENCES

- [1] Dunn J, Ng SK, Holland J, Aitken J, Youl P, Baade PD et al. Trajectories of psychological distress after colorectal cancer. *PsychoOncology* 2013; 22(8): 1759-1765.
- [2] Kwak M, Zebrack BJ, Meeske KA, Embry L, Auilar C, Block R et al. Trajectories of psychological distress in adolescent and young adult patients with cancer: A 1-year longitudinal study. *J Clin Onc* 2013; 31(17): 2160-2166.
- [3] Holland,JC, Andersen B, Breitbart WS, Buchmann LO, Compas B, Deshields TL et al. Distress Management. *J Natl Compr Canc Netw* 2013; 11: 190-209.
- [4] Stanton AL, Leucken LJ, MacKinnon DP & Thompson EH. Mechanisms in psychosocial interventions for adults living with cancer: Opportunity for integration of theory, research, and practice. *J Consult Clin Psychol* 2012; 81(2): 318-335.
- [5] Watson M, Greer S, Young Q, Burgess C & Robertson B. Development of a questionnaire measure of adjustment to cancer: the MAC scale. *Psychol Med* 1988; 18: 203-209.
- [6] Watson M, Law M, Dos Santos M, Greer S, Baruch J & Bliss J. The Mini-MAC: further development of the Mental Adjustment to Cancer Scale. *J Psychosoc Oncol* 1994; 12(3): 33-46.
- [7] Greer S. Psychological response to cancer and survival. *Psychol Med* 1991; 21(1): 43-49.
- [8] Kugaya A, Akechi T, Okamura H, Mikami I & Uchitomi Y. Correlates of depressed mood in ambulatory head and neck cancer patients. *PsychoOncology* 1999; 8: 494-499.
- [9] Watson M, Homewood J, Haviland J & Bliss JM. Influence of psychological response on breast cancer survival: 10-year follow –up of a population-based cohort. *Eur J Cancer*, 2005; 41: 1710-1714.
- [10] Mulcare H, Schofield P, Kashima Y, Milgrom J, Wirth A, Bishop M, et al. Adjustment to cancer and the information needs of people with lung cancer. *PsychoOncology* 2010; 20(5): 488-496.
- [11] Mulcare H, Kashima Y, Milgrom J, Wheeler G, Wirth A, Bishop M et al.

Avoidant adjustment predicts lower information seeking in people with lung cancer. *PsychoOncology* 2012; 22(3): 540-547.

- [12] Watson M & Homewood J. Mental Adjustment to Cancer Scale: Psychometric properties in a large cancer cohort. *PsychoOncology* 2008; 17(11): 1146-1151.
- [13] Ho S, Fung W, Chan C, Watson M & Tsui K. Psychometric properties of the Chinese version of the Mini-Mental Adjustment to Cancer (MINI-MAC) scale. *PsychoOncology* 2003; 12(6): 547-556.
- [14] Anagnostopoulos F, Kolokotroni P, Spanea E & Chrysoschoou M. The Mini-Mental Adjustment to Cancer (Mini-MAC) scale: construct validation with a Greek sample of breast cancer patients. *PsychoOncology*, 2006; 15(1): 79-89.
- [15] Grassi L, Buda P, Cavana L, Annunziata M, Torta R & Varetto A. Styles of coping with cancer: The Italian version of the Mini-Mental Adjustment to Cancer (Mini-MAC) scale. *PsychoOncology* 2004; 14(2): 115-124.
- [16] Kang JI, Chung HC, Kim SJ, Choi HY, Ahn BJ, Jeung H & Namkoong K. Standardization of the Korean version of the Mini-Mental Adjustment to Cancer (K-Mini-MAC) scale: factor structure, reliability and validity. *PsychoOncology* 2007; 17: 592-597.
- [17] Bredal I. The Norwegian version of the Mini-Mental Adjustment to Cancer Scale: factor structure and psychometric properties. *PsychoOncology* 2010; 19(2): 216-221.
- [18] Pereira FMP & de Brito Santos CSV. Initial validation of the Mini-Mental Adjustment to Cancer (Mini-MAC) scale: Study of Portuguese end-of-life cancer patients. *Nursing* 2014; 18(5): 534-539.
- [19] Costa-Requena G & Gil F. The mental adjustment to cancer scale: a psychometric analysis in Spanish cancer patients/ *PsychoOncology* 2009; 18(9): 984-991
- [20] Hulbert-Williams N, Neal R, Morrison V, Hood K & Wilkinson C. Anxiety, depression and quality of life after cancer diagnosis: What psychosocial variables best predict how patients adjust? *PsychoOncology* 2012; 21(8): 857-867.

- [21] *Reference removed to facilitate anonymous peer-review*
- [22] Anastasi A & Urbina S. *Psychological Testing (7th ed.)*. New Jersey: Prentice Hall, 1997.
- [23] Fabrigar LR, Wegener DT, MacCallum RC & Strahan EJ. Evaluating the use of exploratory factor analysis in psychological research. *Psychol Methods* 1999; 4(3): 272.
- [24] Cortina J. What is coefficient alpha? An examination of theory and applications. *J Appl Psychol* 1993; 78(1): 98-104.
- [25] Kline P. *The handbook of psychological testing (2nd ed.)*. London: Routledge 1999.
- [26] Bentler PM & Chou CP' Practical issues in structural modeling. *Sociol Methods Res* 1987;16: 78-117.
- [27] Zigmond A & Snaith R. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 1983; 67: 361-370.
- [28] Cella DF, Tulsky DS, Gray G, Sarafian B, Lloyd S, Linn E, et al. The Functional Assessment of Cancer Therapy (FACT) scale: Development and validation of the general measure. *J Clin Onc* 1993; 11(3): 570-579
- [29] Cohen S, Kamarck T & Mermelstein R. A global measure of perceived stress. *J Health Social Behav* 1986; 24: 385-396.
- [30] Carver CS & Antoni MH. Finding benefit in breast cancer during the year after diagnosis predicts better adjustment 5 to 8 years after diagnosis. *Health Psychol* 2004; 26: 595-598.
- [31] Loehlin JC. *Latent variable models*. London: Lawrence Erlbaum Associates 1998.
- [32] Loehlin JC & Beaujean AA. *Latent variable models: An introduction to factor, path, and structural equation analysis*. New York: Taylor & Francis 2016
- [33] Nunnally JC & Bernstein IH. *Psychometric theory*. New York: McGraw-Hill 1994.
- [34] Streiner DL. Starting at the Beginning: An Introduction to Coefficient Alpha and Internal Consistency. *J Pers Assess* 2003; 80(1): 99–103.
- [35] Cronbach,LJ. Coefficient alpha and the internal structure of tests. *Psychometrika* 1951; 16(3): 297–334.

- [36] Brennan J. Adjustment to cancer – coping or personal transition? *PsychoOncology* 2001; 10: 1-18.
- [37] Hulbert-Williams N, Morrison V, Wilkinson C & Neal R. Investigating the cognitive precursors of emotional response to cancer stress: Re-testing Lazarus's transactional model. *Brit J Health Psychol* 2013; 18(1): 97-112.
- [38] Hulbert-Williams N, Storey L & Wilson K. Psychological interventions for patients with cancer: Psychological flexibility and the potential utility of Acceptance and Commitment Therapy. *Eur J Cancer Care* 2015; 24(1): 15-27.
- [39] Greer S, Moorey S & Watson M. Patients' adjustment to cancer: The Mental Adjustment to Cancer (MAC) scale vs clinical ratings. *J Psychosom Res* 1989; 33(3): 373-377.
- [40] Wakefield CE, Fardell JE, Doolan E, Aaronson NK, Jacobsen PB, Cohn RJ & King M. Participation in psychosocial oncology and quality-of-life research: a systematic review. *Lancet Oncol* 2017; 18(3):153-165.
- [41] Hulbert-Williams NJ, Pendrous R, Hulbert-Williams L & Swash. Recruiting cancer survivors into research studies: piloting online recruitment strategies. *eCancer*, submitted.
- [42] Ross S, Grant A, Counsell C, Gillespie W, Russell I & Prescott R. Barriers to participation in randomised controlled trials: A systematic review. *J Clin Epid* 1999; 52(12): 1143-1156.

Table 1. Items included in the PIC (numbers indicate their respective question number in the original Mini-MAC).

Factor	Question	Item-total correlation
Cognitive Distress	6.I feel completely at a loss about what to do.	.532
	15.I feel there is nothing I can do to help myself.	.537
	16.I think it is the end of the world.	.628
Cognitive Avoidance	17.Not thinking about it helps me cope.	.392
	26.I make a positive effort not to think about my illness.	.423
	27.I deliberately push all thoughts of cancer out of my mind.	.453
Fighting Spirit	2.I see my illness as a challenge.	.337
	10.I try to fight the illness.	.434
	23.I am determined to beat this disease.	.426
Emotional Distress	9.I worry about the cancer returning or getting worse.	.648
	13.I am apprehensive.	.687
	29.I am a little frightened.	.732

Table 2. Overall demographic and clinical characteristics for the combined sample, including inferential tests comparing samples.

Characteristics	N (%)	M (SD)	Differences between sub-samples
Gender (n=183)			$\chi^2=9.188$, df=2, p=.01
Male	90 (49.2)		
Female	92 (50.3)		
Missing	1 (0.5)		
Cancer type			
Breast	42 (23.0)		
Prostate	17 (9.3)		
Lung	101 (55.2)		
Colorectal	21 (11.5)		
Missing	1 (0.5)		
Disease Stage			$\chi^2=66.202$, df=2, p<.01
Not determined	11 (6.0)		
Stage I	21 (11.5)		
Stage II	39 (21.3)		
Stage III	40 (21.9)		
Stage IV	28 (15.3)		
Limited / Localised	8 (4.4)		
Extensive	14 (7.7)		
Missing	22 (12.0)		
Treatment intent			$\chi^2=14.016$, df=2, p<.01
Curative	119 (65.0)		
Palliative	45 (24.6)		
Missing	19 (10.4)		
Days from diagnosis to consent		163.3 (115)	F(2, 158)=33.083, p<.01

Table 3. Results of the CFA, including internal consistency and mean inter-item correlations ($n=183$).

	Cronbach's alpha	Inter-item correlation	Confirmatory Factor Analysis			
			RMSEA	ECVI	PNFI	AGFI
Five-factor Mini-MAC model (29 items)			.084	5.020	.582	.727
Hope/Helplessness	.851	.383				
Anxious	.885	.454				
Preoccupation	.770	.456				
Cognitive Avoidance	.728	.412				
Fighting Spirit	.572	.217				
Fatalism						
Four-factor Mini-MAC model (25 items)			.097	4.254	.568	.699
Cognitive Distress	.832	.316				
Cognitive Avoidance	.738	.373				
Fighting Spirit	.697	.445				
Emotional Distress	.583	.221				
Psychological Impact of Cancer Scale (12 items)			.083	.923	.604	.857
Cognitive Distress	.623	.359				
Cognitive Avoidance	.752	.517				
Fighting Spirit	.697	.445				
Emotional Distress	.713	.456				

RMSEA=Root Mean Square Error of Approximation; ECVI=Expected Cross Validation Index; PNFI=Parsimony-adjusted Normal Fit Index; AGFI=Adjusted Goodness of Fit Index)

Table 4. Summary of results from test-retest reliability analyses, and correlation with commonly used psychosocial outcome measures (sample size indicated in parentheses).

	Test-retest reliability	Construct Validity				
		Quality of Life	Anxiety	Depression	Perceived Stress	Benefit Finding
Five-factor Mini-MAC model (29 items)						
Hope/Helplessness	.502** (n=55)	-.009 (n=81)	.387** (n=164)	.356** (n=164)	.267** (n=94)	-.137 (n=92)
Anxious Preoccupation	.789** (n=52)	.158 (n=81)	.563** (n=164)	.299** (n=164)	.413** (n=94)	-.165 (n=92)
Cognitive Avoidance	.667** (n=54)	.029 (n=81)	.234** (n=164)	.101 (n=164)	.050 (n=94)	.111 (n=92)
Fighting Spirit	.628** (n=52)	.149 (n=81)	-.083 (n=164)	-.245** (n=164)	-.035 (n=94)	.380** (n=92)
Fatalism	.889** (n=47)	.325** (n=81)	-.027 (n=164)	-.148* (n=164)	.191* (n=94)	.508** (n=92)
Four-factor Mini-MAC model (25 items)						
Cognitive Distress	.633** (n=58)	.074 (n=81)	.514** (n=164)	.347** (n=164)	.350** (n=94)	-.171 (n=92)
Cognitive Avoidance	.691** (n=54)	.052 (n=81)	.331** (n=164)	.160* (n=164)	.175* (n=94)	.108 (n=92)
Fighting Spirit	.670** (n=52)	.150 (n=81)	-.010 (n=164)	-.184** (n=164)	.033 (n=94)	.316** (n=92)
Emotional Distress	.638** (n=55)	.344** (n=81)	.497** (n=164)	.214** (n=164)	.488** (n=94)	-.071 (n=92)
Psychological Impact of Cancer Scale (12 items)						
Cognitive Distress	.504** (n=56)	.043 (n=81)	.382** (n=164)	.259** (n=164)	.302** (n=94)	-.118 (n=92)
Cognitive Avoidance	.648** (n=56)	-.017 (n=81)	.155* (n=164)	.066 (n=164)	-.020 (n=94)	.100 (n=92)
Fighting Spirit	.670** (n=52)	.150 (n=81)	-.010 (n=164)	-.184** (n=164)	.033 (n=94)	.316** (n=92)
Emotional Distress	.620** (n=55)	.228* (n=81)	.549** (n=164)	.238** (n=164)	.458** (n=94)	-.166 (n=92)

* $p < .05$, ** $p < .01$