

## Prevalence, Risk Factors and Self-awareness for Hypertension and Diabetes: Rural-Urban and Male-Female Dimensions from a cross-sectional study in Ghana

Item Type	Article
Authors	Ellahi, Basma;Dikmen, Derya;Faith, Agbozo;Francis, Zotor;Amanda, Aitken;Bilge, Seyhan-Erdoğan;Omer, Faruk Karabulut
Citation	Ellahi, B., Dikmen, D., Seyhan-Erdoğan, B., Karabulut, O. F., Aitken, A., Agbozoc, F., & Zotor, F. B. (2023). Prevalence, risk factors and self-awareness for hypertension and diabetes: Rural-urban and male-female dimensions from a cross-sectional study in Ghana. <i>International Journal of Diabetes in Developing Countries</i> , 43, 694–708. <a href="https://doi.org/10.1007/s13410-022-01141-9">https://doi.org/10.1007/s13410-022-01141-9</a>
DOI	<a href="https://doi.org/10.1007/s13410-022-01141-9">10.1007/s13410-022-01141-9</a>
Publisher	Springer
Journal	<i>International Journal of Diabetes in Developing Countries</i>
Download date	2026-05-19 15:26:24
Item License	<a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Link to Item	<a href="http://hdl.handle.net/10034/627345">http://hdl.handle.net/10034/627345</a>

**Prevalence, Risk Factors and Self-awareness for Hypertension and Diabetes: Rural-Urban and Male-Female Dimensions from a cross-sectional study in Ghana**

Basma Ellahi<sup>a</sup>, Derya Dikmen<sup>b</sup>, Bilge Seyhan-Erdoğan<sup>b</sup>, Omer Faruk Karabulut<sup>b</sup>, Amanda Aitken<sup>a</sup> Faith Agbozo<sup>c</sup> and Francis B. Zotor<sup>c</sup>

<sup>a</sup> Faculty of Health and Social Care, University of Chester, Chester UK

<sup>b</sup>Hacettepe University, Faculty of Health Sciences, Department of Nutrition and Dietetics, Ankara, Turkey

<sup>c</sup>Department of Family and Community Health, School of Public Health, University of Health and Allied Sciences, Ho Ghana

\*Corresponding author: Professor Francis Zotor

Contact details: [francisfirst@gmail.com](mailto:francisfirst@gmail.com)

## **Abstract**

**Background:** Hypertension and diabetes remain the primary cause of non-communicable disease (NCD) related morbidity and mortality globally. Rural-urban transitions in developing countries might aggravate the risk factors and prevalence of these conditions. The study aims to determine prevalence, demographic, anthropometric and diet-related predisposing factors for hypertension and diabetes among urban and rural dwellers and assess participants' self-awareness of their hypertension and diabetes status.

**Methods:** This cross-sectional survey involved 850 adult males and females age  $\geq 18$  years residing in urban and rural areas in the Hohoe Municipality of Ghana, randomly sampled using probability proportional to size. Data included demographic, anthropometric, physiologic (blood pressure, fasting blood glucose) and dietary information. Nutrient quantities were analysed using the Research to Improve Infant Nutrition and Growth (RIING Nutrient Database Software. All other data was analysed in SPSS (v25). Risk factors for hypertension were estimated through ordinal logistic regression and the odds ratio (OR) with the corresponding 95% confidence level (CI) documented.

**Results:** More females participated than males (58.4% vs 41.6%), similarly rural compared to urban inhabitants (53.5% vs 47.5%,  $p=0.002$ ) with a mean age of  $47.3 \pm 16.1$  years. Females generally had higher adiposity, rural dwellers had higher BMI whereas urban dwellers had higher waist and hip circumferences. Overall, 4.4% and 18.5% were diabetics and pre-diabetics; while 20.4% and 12.1% were overweight and obese respectively. Of the 36.8% hypertensives, only 18.2% were aware of their status, with significant male (40.3%) female (59.7%); and urban (43.5%) rural (56.6%) differences. Males had higher intakes of energy and nutrients of public health importance to hypertension, similar as rural inhabitants except for

cholesterol. Advancing age (95% CI: 0.02-0.05), being male (OR: 1.56 95% CI:0.12-0.81) and increased BMI (95% CI: 0.01-0.11) were independently associated with hypertension.

**Conclusions:** A third of the population were hypertensives but were unaware. Findings challenge the assumption of lower predisposition among rural inhabitants as we observe lifestyle habits consistent with increasing urbanisation. Efforts to mitigate rising NCD trend requires intensified community-based screening, awareness creation, and lifestyle interventions to improve diet, physical activity and health seeking.

**Keywords:** Elevated blood pressure; elevated blood glucose; non-communicable diseases; adiposity; body mass index; urbanization; undiagnosed

## Background

Globally, nutrition related non-communicable diseases (NCDs) including cardiovascular diseases, hypertension and diabetes mellitus (DM) are the primary cause of morbidity and mortality (1). According to the World Health Organization (WHO), NCDs are responsible for 71% (41 million) of the 57 million deaths globally and 85% of NCD deaths are reported in low- and middle-income countries (LMIC) (2). Self-reported high blood pressure has increased from 594 million (in 1975) to 1.3 billion (in 2015) with the increase mostly in LMIC (2). Hypertension is a risk factor for many NCDs worldwide, with the highest prevalence of 27% reported in African (2). In sub-Saharan Africa, 19.4 million adults aged 20-79 years had diabetes in 2019 with 6.8% of all-cause mortality attributed to diabetes (3).

NCDs are estimated to account for 43% of all deaths (4) with hypertension and diabetes ranked seventh and ninth respectively among the ten most common causes of mortality in Ghana, in 2016 (5). The current situation of hypertension in many LMIC is a burden on resource limited health care and is already higher than in high-income countries (6). In Ghana, a systematic review reported an increase in hypertension prevalence from 19 to 48% from 1970 to 2007 (7). Diabetes in Ghana represents 2% of all deaths and prevalence is around 4.8% but is predicted to rise (8). Meanwhile, related risk factors of overweight and obesity are at 30.8% and 10.9% respectively (9), albeit the absence of a national survey data assessing blood glucose levels.

Worldwide, demographic, socio-economic, dietary, lifestyle and behavioural factors are associated risk factors for hypertension and this is similar for Africa (10). NCDs has increased notably from the growth in urbanisation in many LMICs (11). Some differences have been observed between urban and rural areas in Ghana in relation to prevalence and occurrence of risk factors, in that urban populations have a higher prevalence and risk factors for hypertension. (12-15). Despite this, there is limited data on the prevalence and risk factors in

rural areas as these areas are given less research attention (12, 14). For instance, in urban Accra in Ghana, reported prevalence of hypertension was 28.3% and most alarmingly, 92.6% of participants were not aware of their condition. Of those that were aware, only 3.9% of participants accessed treatment (13).

Urbanisation is considered a determinant of health and one of the key drivers of NCDs, especially in LMICs (16), although industrialisation varies from country to country and differences between urban and rural regions of the same geographical area are evident (17). Whilst this has been demonstrated in previous research in many areas of Ghana where economic shifts have resulted in increasing urbanisation, the impact of urbanisation in transitional areas of the country is rarely reported. This applies to largely rural areas now undergoing development and urbanisation as a consequence of economic development. Ghana is experiencing growth in its cities and districts due to urbanisation and these areas are being redefined as peri-urban and urban areas. Furthermore, occupation and gender roles of women particularly in rural areas is often different from urban areas where many women find employment outside the home and area of residence.

Determining the prevalence and risk factors of hypertension in peri-urban and rural areas is important to guide public health policy. Furthermore, assessing the awareness is critical not only to inform health promotion activities, but also to prevent development of related NCDs such as diabetes (18). Thus, we need to assess the prevalence of hypertension and diabetes in the context of wider risk factors for NCDs among adult males and females residing in urban and rural areas in Ghana. We aimed to undertake a cross-sectional survey in a rural and peri-urban setting in Ghana. We assessed prevalence and associated risk factors for hypertension and diabetes. We also compared the rates among males and females in rural and developing

peri-urban sub-districts of the Hohoe municipality in Ghana and included an additional focus on diabetes to assess the current situation.

## **Methods**

### **Study Design and Location**

The cross-sectional survey was conducted in the Hohoe Municipality, with data collection completed in 2018. The Hohoe municipality is one of the 17 administrative districts in the newly re-demarcated Volta region, one of 16 regions in Ghana. Hohoe municipality, the second largest municipality in the region, is located in the north-eastern end of the region, about 78 and 218 kilometre (km) from Ho, the Volta regional Capital and Accra, the national capital respectively. The municipality shares its boundaries with the Republic of Togo to the East, Jasikan District to the North, Kpando District to the West and Afadjato South District to the South. It covers an area of 1,403 km<sup>2</sup> consisting of 174 communities with a population of 188,963 comprising 52.1% females and 47.9% males. (19). The municipality is youthful (35.9% of the population are under age 15 years) and also largely urban (52.6%) (19).

### **Study Population**

Inclusion criteria: Free-living adult males and self-reporting non-pregnant females age 18 years and above were recruited from urban and rural areas of the Hohoe municipality (n=850).

Exclusion criteria: Males and females under 18 and not residing in the Hohoe municipality were excluded from the study. Individual written informed consent was obtained from all participants.

Study communities were selected by random sampling using probability proportional to size. This method was carried out by creating a cumulative list of communities with the respective

populations in the six sub-districts of the Hohoe municipality (Hohoe, Gbi rural, Alavanyo, Agumatsa, Likpe, Akpafu/Santrokofi) (Table 1) after which a systematic sample was drawn from a random start. Based on the WHO’s Expanded Programme on Immunization (EPI) sampling method, 30 clusters representing 30 communities were chosen and the total municipality population of 195,856 inhabitants divided by the 30 clusters to obtain the sampling interval (20). This generated a sample size of 850 participants. Thereafter, proportional allocation was employed to derive a representative sample size from each community identified and bias was eliminated.

**Table 1: Distribution of participants by sub-districts and geographical location**

<b>Sub-district</b>	<b>Classification of geographical location</b>	<b>n</b>	<b>%</b>
Hohoe	Urban	405	47.6
Gbi-rural	Rural	52	6.1
Alavanyo	Rural	36	4.2
Akpafu/Sanko	Rural	77	9.1
Agumatsa	Rural	39	4.6
Likpe	Rural	241	28.4
Total		850	100

n: participants

### **Data Collection Tools and Procedures**

A pre-piloted structured quantitative questionnaire with mainly closed questions was used to collect the data. Information elicited included demographic, anthropometric and physiologic measurements. Dietary data was collected using a multiple-pass 24-hour recall method and common household measures used as visual tools to assist in quantifying the portion sizes. Participants were asked to provide a list of all foods, drinks, beverages and snacks eaten the day prior to the survey from when they awake in the morning to when they go to sleep. Data was collected by trained health professionals pursuing a top-up Bachelor degree in Public Health Nutrition at the School of Public Health, University of Health and Allied Sciences, Hohoe Ghana (F.Z, F.A and B.E). Data was obtained through face-to-face interview using the questionnaire and measurements tools. Data collected on a daily basis was checked by

members of the research team to check for missing data and address this (O.F.K, D.D, and B.E). Data was inputted into the databases by two assistants and checked by other members of the research team.

### **Anthropometric Measurements**

Weight, height, waist and hip measurements were recorded according to the Centre for Disease Control standardised anthropometric (21) protocols by trained researchers with participants wearing lightweight clothing without shoes. All measurements were taken in triplicate and the two nearest measurements recorded to achieve consistency of data collected. Height was measured with a SECA stadiometer (SECA 875) to the nearest 0.1 centimetres (cm). Weight was measured with a SECA digital scale (SECA 217) to the nearest 0.1 kilogram. Waist and hip circumference were measured using a non-extensible measuring tape to the nearest 0.1 cm. Waist circumference was measured an inch above the upper hip bone with the tape measure drawn around the waist parallel to the floor. The hip was identified at the widest part around the buttocks.

### **Physiological Measurements**

Resting upper arm arterial blood pressure (BP) was measured using Omron digital sphygmomanometer recorded in mmHg. Repeated measurements were taken in triplicates at five minutes intervals and the average of the two nearest measurements recorded. Fasting blood glucose level was measured from a capillary thumb prick using OneTouch digital glucometer and recorded in mmol/L. Only subjects who were fasting at the point of data collection (before 9 am) had their fasting blood glucose levels measured.

## **Data Handling and Statistical Analysis**

Dietary information was analysed using the Research to Improve Infant Nutrition and Growth (RIING) Project Nutrient Database Software developed by the Department of Nutrition and Food Science, University of Ghana. The focus was on energy and nutrients of public health importance to hypertension. These included protein, carbohydrate, total fat, cholesterol, magnesium, potassium, sodium, and fibre. Nutrient values were automatically generated onto an Excel sheet and checked by a second researcher. Out-of-range values were flagged and cross-checked to identify possible coding errors (O.F.K). The Excel output was exported into Statistics Package for Social Sciences (SPSS, 24.0 IBM, USA) for analysis. Energy and nutrient values were compared with the Public Health England dietary reference values (22). Under or over-reporting of dietary intakes were calculated as the ratio of energy intake (EI) to Basal Metabolic Rate (BMR) (23). The EI to BMR ratio was calculated for each individual and the cut off values used to identify under-reporters ( $<1.14-1.34$ ), normal reporters ( $1.35-2.39$ ) and over-reporters ( $\geq 2.4$ ).

Waist-to-hip ratio (WHR) was calculated by dividing waist circumference (WC) by hip circumference (HC). Cut-offs for obesity defined by WHR and WC was  $>0.8$  and  $>88$  cm respectively as recommended by the WHO (24). Body mass index (BMI) was calculated based on WHO (25) criteria as weight in kg divided by height in meters squared, and the corresponding BMI defined as:  $<18.5$  kg/m<sup>2</sup> underweight;  $18.5 - 24.9$  kg/m<sup>2</sup> normal weight;  $25.0-29.9$  kg/m<sup>2</sup> overweight; and  $>30$  kg/m<sup>2</sup> obese.

Hypertension was defined as a systolic BP of at least 140 mmHg and a diastolic BP of at least 90 mmHg according to National Institute for Health and Care Excellence (NICE) criteria (26). Also, BP  $<130/85$  mmHg was considered as normal and 130 to 139/85 to 89 mmHg as high normal (pre-hypertension).

Fasting blood glucose less than 5.6 mmol/L was considered normal, 5.6 to 6.9 mmol/L was considered pre-diabetic (impaired fasting glucose) and 7 mmol/L or higher was defined as diabetes according to the WHO criteria (27).

Data analysis focused on the relationship between hypertension and diabetes status of individuals and their anthropometric measurements, dietary intakes and their awareness thereof. All statistical analyses were performed using the Statistics Package for Social Sciences (SPSS IBM version 25.) Kolmogorov-Smirnov was used to test the normality of variables. Comparisons between groups were made using one-way analysis of variance (ANOVA) if the data were normally distributed and Kruskal-Wallis test was used for the non-parametric distribution. Post-hoc Bonferonni correction and Dunn's test were used for ANOVA. Differences in nutrient intakes were examined with Mann-Whitney U test. Categorical data were examined using Chi-square test and reported as frequencies and percentages. Continuous data were presented as means with the corresponding standard deviation (SD). The relationship between hypertension, diabetes and its risk factors were examined through ordinal logistic regression by computing the unadjusted odds ratio at 95% confidence level (CI). Differences  $p < 0.05$ . were considered significant.

## **Results**

### *Demography of participants*

Demographic characteristics of study participants are presented in Table 1 and 2. Overall, 41.6% were males and 58.4% were females ( $p=0.002$ ); 47.5% resided in urban areas and 53.5% in rural areas. Mean age of participants was  $47.3 \pm 16.1$  years. In the rural and urban areas, the primary occupations were farming and trading respectively. In terms of educational level, the highest level of completed education among the majority was junior high school. Table 3

represents the demographic analysis of participants categorised according to gender and place of residence.

**Table 2. Demographic status of participants according to gender and place of residence**

Variables	All n (%)	Male n (%)	Female n (%)	Urban n (%)	Rural n (%)
<b>Gender</b>					
Male	354 (41.6)			168(47.5)	186(52.5)
Female	496 (58.4)			237(47.8)	259(52.2)
<b>Age groups</b>					
<20	12 (1.4)	8 (2.3)	4 (0.8)	8 (2.0)	4 (0.9)
20-29	129 (15.2)	70 (19.8)	59 (11.9)	86 (21.2)	43 (9.7)
30-39	148 (17.4)	66 (18.6)	82 (16.5)	91 (22.5)	57 (12.8)
40-49	162 (19.1)	68 (19.2)	94 (19.0)	76 (18.8)	86 (19.3)
50-59	184 (21.6)	58 (16.4)	126 (25.4)	75 (18.5)	109 (24.5)
60-69	130 (15.3)	42 (11.9)	88 (17.7)	39 (9.6)	91 (20.4)
70-79	68 (8.0)	35 (9.9)	33 (6.7)	24 (5.9)	44 (9.9)
80-89	17 (2.0)	7 (2.0)	10 (2.0)	6 (1.5)	11 (2.5)
<b>Occupation</b>					
Trading	208 (24.5)	23(6.5)	185(37.3)	133(32.8)	75(16.9)
Farming	262(30.8)	113(31.9)	149(30.0)	37(9.1)	225(50.6)
Artisan	146(17.2)	92(26.0)	54(10.9)	95(23.5)	51(11.5)
Formal Job	81(9.5)	48(13.6)	33(6.7)	55(13.6)	26(5.8)
Schooling	38(4.5)	27(7.6)	11(2.2)	26(6.4)	12(2.7)
Retired	30(3.5)	21(5.9)	9(1.8)	11(2.7)	19(4.3)
Unemployed	61(7.2)	15(4.2)	46(9.3)	30(7.4)	31(7.0)
Others*	24(2.8)	15(4.2)	9(1.8)	18(4.4)	6(1.3)
<b>Education</b>					
No education	78(9.2)	14(4.0)	64(12.9)	47(11.6)	31(7.0)
Primary school	117(13.8)	25(7.1)	92(18.5)	56(13.8)	61(13.7)
Junior High school	397(46.7)	149(42.1)	248(50.0)	175(43.2)	222(49.9)
Senior High school	149(17.5)	91(25.7)	58(11.7)	63(15.6)	86(19.3)
Tertiary	109(12.8)	75(21.2)	34(6.9)	64(15.8)	45(10.1)
<b>Sub-district</b>					
Urban	405(47.5)	168(47.5)	237(47.8)		
Rural	445(53.5)	186(52.5)	259(52.2)		
<b>Total</b>	<b>850</b>	<b>354</b>	<b>496</b>	<b>405</b>	<b>445</b>

\*Not included in any other group such as watchman

**Table 3. Anthropometry and dietary intakes according to gender and place of residence**

Variables	Mean (SD)			p-value	Mean (SD)		P-value
	All	Male (n=354)	Female (n=496)		Urban (n=405)	Rural (n=445)	
Age (years)*	47.3 (16.1)	45.4 (17.2)	48.6 (15.2)	<b>0.002</b>	43.2 (16.0)	51.0 (15.4)	<b>&lt;0.001</b>
Body mass index (kg/m <sup>2</sup> )*	24.6 (5.2)	22.5 (3.6)	26.1 (5.6)	<b>&lt;0.001</b>	22.5 (5.5)	23.8 (4.8)	<b>&lt;0.001</b>
Waist circumference (cm)*	83.2 (14.8)	79.8 (10.4)	85.6 (14.1)	<b>&lt;0.001</b>	84.9 (12.6)	81.6 (13.2)	<b>&lt;0.001</b>

Hip circumference (cm)*	96.5 (12.3)	91.9 (8.04)	99.8 (13.63)	<b>&lt;0.001</b>	98.9 (11.3)	94.4 (12.8)	<b>&lt;0.001</b>
Waist hip ratio*	0.86 (0.08)	0.86 (0.08)	0.86 (0.08)	0.113	0.86 (0.08)	0.86 (0.08)	0.083
Fasting glucose (mmol/L)*	5.35 (1.41)	5.42 (1.46)	5.29 (1.37)	0.200	5.46 (1.57)	5.23 (1.20)	0.158
Diabetes <sup>z</sup> (n %)	61 (9.2)	27 (9.6)	34 (8.8)	0.464	46 (10.1)	15 (7.2)	0.173
Diastolic BP (mmHg)*	82.35 (14.2)	81.7 (14.3)	82.8 (15.2)	0.188	81.99 (15.0)	82.68 (14.7)	0.510
Systolic BP (mmHg)*	132.2 (23.9)	132.6 (22.6)	131.7 (24.9)	0.598	130.0 (22.9)	134.0 (24.8)	0.054
Energy (kcal)	1898.21 (818.02)	2032.41 (863.81)	1802.43 (770.40)	<b>&lt;0.001</b>	1707.86 (706.79)	2071.45 (872.85)	<b>&lt;0.001</b>
Protein (g)	97.66 (72.29)	98.66 (74.21)	96.94 (70.95)	0.732	72.90 (49.35)	120.18 (81.91)	<b>&lt;0.001</b>
Carbohydrate (g)	248.17 (118.42)	275.85 (126.58)	228.42 (108.10)	<b>&lt;0.001</b>	240.59 (114.27)	255.07 (121.79)	0.117
Total Fat (g)	52.93 (34.54)	56.06 (36.57)	50.70 (32.88)	<b>0.028</b>	49.87 (31.63)	55.72 (36.81)	<b>0.037</b>
Cholesterol (mg)	32.83 (84.26)	37.51 (88.9)	29.49 (80.70)	0.178	41.48 (89.46)	24.96 (78.5)	<b>&lt;0.001</b>
Fibre (g)	18.37 (12.74)	21.2 (14.27)	16.33 (11.10)	<b>&lt;0.001</b>	18.09 (12.90)	18.61 (12.60)	0.381
Magnesium(mg)	58.35 (68.35)	66.03 (74.92)	54.58 (62.89)	<b>0.016</b>	66.03 (74.92)	54.58 (62.89)	<b>&lt;0.001</b>
Potassium (mg)	375.36 (464.4)	450.10 (564.74)	322.02 (368.36)	<b>&lt;0.001</b>	450.10 (564.74)	322.02 (368.36)	<b>&lt;0.001</b>
Sodium (mg)	498.29 (1493.26)	604.01 (1992.56)	422.84 (989.66)	0.081	604.01 (1992.56)	422.84 (989.66)	<b>&lt;0.001</b>
Smoking <sup>z</sup> (Yes) (n %)	49 (5.8)	46 (13)	3 (0.6)	$\chi^2=58.36$ , <b>p&lt;0.001</b>	13 (3.2)	36 (8.1)	$\chi^2=9.295$ , <b>p=0.002</b>
Alcohol consumption <sup>z</sup> (Yes) (n %)	354 (41.6)	207 (58.5)	147 (29.6)	$\chi^2=70.68$ , <b>p&lt;0.001</b>	130 (32.1)	224 (50.3)	$\chi^2=29.02$ , <b>p=0.001</b>
Physical activity <sup>z</sup> (Yes) (n %)	631 (74.2)	297 (83.9)	334 (67.3)	$\chi^2=30.80$ , <b>p&lt;0.001</b>	268 (66.2)	363 (81.6)	$\chi^2=26.291$ , <b>p&lt;0.001</b>

\*Mann-Whitney U test reporting the mean. <sup>z</sup> Chi-square test reporting the proportions. Values with mean and standard deviations were continuous and number and percentage were categorical. Fasting glucose and diabetes was measured for n=665 participants who were still in a fasting state at the time of data collection. BP = blood pressure

### *Anthropometric indicators*

Females had a higher BMI than males ( $p<0.001$ ) with a mean BMI of  $26.1 \pm 5.6$  kg/m<sup>2</sup> compared to  $22.5 \pm 3.6$  kg/m<sup>2</sup> for their male counterparts. Rural inhabitants had higher BMI than urban inhabitants ( $p<0.05$ ). Also, females had higher waist and hip circumferences than males ( $p<0.001$ ).

### *Dietary intakes and lifestyle habits*

Dietary intakes are presented in Table 3. The mean energy intake was  $2032.41 \pm 863.81$  kcal for males and  $1802.43 \pm 770.40$  kcal for females. Mean protein intake for males and females was  $98.66 \pm 74.21$  and  $96.94 \pm 70.95$  g, respectively. Mean carbohydrate intake was  $275.85 \pm 126.58$  and  $228.42 \pm 108.10$  g for males and females respectively. Public Health England recommends protein intakes of 55.5g/day for males and 45.0g/day for females, and carbohydrate intakes of at least 333g/day and 267 g/day for males and females respectively (22). Our findings show higher protein and lower carbohydrate intake than the recommendations in both males and females. Energy, carbohydrate, total fat, fibre, magnesium and potassium intakes of males were significantly higher than females. Regarding the accuracy of reporting dietary intakes, 43.5% were under-reporters, 44% reported as actually consumed, and 12.3% were over-reporters. There was no statistical difference between gender, BMI and WC classifications among under- or over-reporters of dietary intake (data not shown).

When we compared urban and rural dwellers, we identified higher intakes of energy, protein, total fat among rural dwellers. Mean energy intake was  $1707.86 \pm 706.79$  kcals, and  $2071.45 \pm 872.85$  kcals, for urban and rural dwellers, respectively. In rural areas, mean protein intake was higher ( $120.18 \pm 81.91$  g) compared to urban areas ( $72.90 \pm 49.35$  g). The significant differences in protein intake might be due to rural communities having access to more nutritious foods of much higher dietary quality probably because they are more inclined to work in animal rearing. But the consumption of plant-based protein is common in rural areas. In urban areas, mean cholesterol intake was  $41.48 \pm 89.46$  g while in rural areas, it was lower ( $24.96 \pm 78.5$  g). Although unsubstantiated, a plausible reason for the high cholesterol levels of urban inhabitants might be due to the availability of cooked and processed foods high in saturated fatty foods. Furthermore, urban inhabitants had higher intakes of magnesium, potassium and sodium compared to rural inhabitants ( $p < 0.05$ ). On a weekly basis, 5.8% of participants smoked cigarettes and 41.6% drank alcohol (obtained as a yes or no binary

question). In relation to frequency of physical activity, there was a significant difference based on gender; males were more physically active than females ( $p < 0.001$ ), and the physical activity took the form of walking.

### *Physiological parameters*

Overall, 665 eligible participants had their FBG measured (the rest had already taken breakfast). Mean FBG was  $5.35 \pm 1.41$  mmol/L with no significant relationship between males and females nor urban and rural dwellers. Mean BP was 133/82 mmHg (SD 22.6/14.3) for males and 132/83 mmHg (SD 24.9/15.2) for females. Diastolic BP was slightly higher in females ( $82.8 \pm 15.2$  mmHg) than males ( $81.7 \pm 14.3$  mmHg) ( $p = 0.188$ ). Mean systolic BP was  $132.1 \pm 23.9$  mmHg in all participants with no statistical difference between males and females. Figure 1 presents participants' educational status and geographical location categorised according to BP values.

Mean diastolic and systolic BP was higher among participants who had only primary or no formal education and also among rural inhabitants compared to urban participants. Systolic BP was significantly higher among rural inhabitants. This finding suggests a relationship with isolated systolic hypertension (ISH). ISH is defined as systolic BP (SBP)  $\geq 140$  mmHg and diastolic BP (DBP)  $< 90$  mmHg (28). ISH is the most common type of hypertension and is associated with older age (28). We noticed a higher mean age in the rural sub-districts (Table 2), yet participants were largely younger than 65 years. Thus, we find this elevated SBP at an earlier age profile among rural communities. Exploring the possible relationships between urban and rural areas as well as both DBP and SBP, we observed a significant relationship between SBP and rural residency ( $p < 0.005$ ).

INSERT FIGURE 1

**Figure 1. Mean diastolic and systolic blood pressure values classified according to educational level and place of residence**

Table 4 presents blood pressure levels according to fasting blood glucose and anthropometric status. Overall, 36.8% had hypertension of which 40.3% were males and 59.7% females. Furthermore, 43.5% of urban dwellers and 56.6% of rural dwellers had hypertension. Fasting blood glucose levels of the hypertensive participants were the highest when compared with those with high-normal and normal measurements ( $p < 0.05$ ). Hypertensive participants had higher BMI than non-hypertensives ( $p < 0.001$ ) with a mean BMI of  $25.32 \pm 5.78 \text{ kg/m}^2$ . According to BMI groups, a significant proportion of the hypertensives were overweight (24.7%) and obese (19.6%) (totalling 44.3%) in comparison to the normal group (totalling 32.5%). Hypertensive participants had the highest WC and WHR than the other participants ( $p < 0.001$ ).

**Table 4. Hypertension status of participants classified according to demographic, glycaemic and anthropometric status**

Variables	Normal** (n=387)	High-Normal** (n=150)	Hypertension** (n=313)	p-value
Allz (%)	45.5	17.7	36.8	$\chi^2=0.398,$ p>0.05
Male (%)	42.4	42.7	40.3	
Female (%)	57.6	57.3	59.7	
Urban (%)	49.9	50.7	43.5	$\chi^2=3.525,$ p=0.172
Rural (%)	50.1	49.3	56.5	
Fasting blood level* (mean±SD)	5.17±1.19	5.47±1.39	5.51±1.62	<b>0.011</b>
Diabetes (%)***	4.4	8.5	10	$\chi^2=10.015,$ p<0.05
IFG (%)***	18.5	20.3	23.3	
Normal (%)***	77.2	71.2	66.7	
BMI* (mean±SD)	23.82±4.85	25.05±4.51	25.32±5.78	<b>&lt;0.001</b>
Underweight (%)	11.4	3.3	6.7	$\chi^2=20.633,$ p<0.05
Normal (%)	56.1	52.7	49.0	
Overweight (%)	20.4	28.0	24.7	
Obese (%)	12.1	16.0	19.6	

WC* (mean±SD)	80.67±11.33	83.81±12.30	86.02±14.00	<0.001
WHR* (mean±SD)	0.85±0.80	0.86±0.67	0.88±0.86	<0.001

\*ANOVA,  $\chi$  Chi-square. IFG = Impaired fasting glucose, BMI = body mass index, WC = waist circumference, WHR = waist-to-hip ratio.

\*\*Normal: Blood pressure <130/85 mmHg; high-normal (pre-hypertension): 130 to 139/85 to 89 mmHg; and hypertension:  $\geq$ 140 mmHg; [25].

\*\*\*Normal: fasting blood glucose <5.6 mmol/L; impaired fasting glucose (pre-diabetic): 5.6-6.9 mmol/L; and diabetes:  $\geq$ 7 mmol/L [26].

#### *Risk factors for hypertension and diabetes*

Age, BMI, gender and educated to SHS/VocTech between education categories were found significant covariates for hypertension (significance level 95%) (Table 5). Aging (95% CI, 0.0251–0.0464;  $p < 0.001$ ) increases the risk for hypertension. Being male is also a risk factor (OR, 1.586, 95% CI, 0.1174–0.8083;  $p = 0.0086$ ) as is an increase in BMI (95% CI, 0.0120–0.1065;  $p = 0.014$ ). Men were about 1.59 times more likely to have high blood pressure than women. SHS/Voc/Tech education level was at lower risk of having high blood pressure than those in the “none” education level. Those who have no education were 2.03 times more likely to have high blood pressure than those SHS/Voc/Tech education level. However, neither energy, dietary magnesium, fat, fibre, calcium, sodium intakes were significantly associated with hypertension.

**Table 5. Ordinal logistic regression showing the risk factors for high blood pressure**

High Blood Pressure							
		$\beta$	Std. Error	p	95% Confidence Interval		Odds ratio
					Lower Bound	Upper Bound	
	Age (years)	0.0358	0.0054	<0.001	0.0251	0.0464	-
	Waist circumference (cm)	0.0144	0.0091	0.1144	-0.0035	0.0323	-
	BMI (kg/m <sup>2</sup> )	0.0592	0.0241	0.0140	0.0120	0.1065	-

	Energy	0.0001	0.0006	0.8336	-0.0011	0.0014	-
	Carbohydrate	0.0009	0.0027	0.7320	-0.0043	0.0062	-
	Fat	-0.0025	0.0058	0.6596	-0.0138	0.0087	-
	Fiber	-0.0083	0.0072	0.2497	-0.0224	0.0058	-
	Sodium	4.16x10 <sup>-6</sup>	0.0001	0.9388	-0.0001	0.0001	-
	Potassium	0.0001	0.0002	0.6021	-0.0004	0.0006	-
	Magnesium	-0.0014	0.0016	0.3885	-0.0046	0.0018	-
	Calcium	-9.11x10 <sup>-6</sup>	0.0001	0.9515	-0.0003	0.0003	-
Gender	Male/female	0.4629	0.1763	0.0086	0.1174	0.8083	1.5886
	Urban/Rural	0.0644	0.1566	0.6808	-0.2426	0.3714	1.0666
Education level	Primary	-0.1292	0.2913	0.6573	-0.7002	0.4417	0.8788
	JHS	-0.3944	0.2659	0.1380	-0.9155	0.1267	0.6741
	SHS/Voc/Tech	-0.7093	0.2913	0.0149	-1.2802	-0.1384	0.4920
	Middle School	-0.3483	0.2745	0.2044	-0.8863	0.1896	0.7059
	Tertiary	-0.5241	0.3570	0.1420	-1.2238	0.1755	0.5921
	Others	2.1502	1.1655	0.0650	-0.1340	4.4345	8.5866
Occupation	Trading	0.6627	0.4245	0.1185	-0.1693	1.4947	1.9401
	Farming	0.5574	0.4233	0.1879	-0.2722	1.3869	1.7460
	Artisan	0.4444	0.4230	0.2935	-0.3847	1.2735	1.5596
	Formal job	0.6348	0.4801	0.1861	-0.3063	1.5758	1.8866
	Schooling	0.8081	0.5202	0.1203	-0.2114	1.8275	2.2435
	Retired	1.0511	0.5878	0.0738	-0.1010	2.2031	2.8607
	Unemployed	0.7824	0.4779	0.1016	-0.1542	1.7190	2.1867
	Alcohol	0.2038	0.1462	0.1633	-0.0827	0.4904	1.2261
	Smoking	-0.1695	0.3011	0.5735	-0.7597	0.4207	0.8441
Physical Activity	Never/none	-0.2669	0.2543	0.2939	-0.7653	0.2315	0.7657
	Once a while	-0.0433	0.2502	0.8627	-0.5336	0.4471	0.9577
	Quite often	-0.2839	0.3142	0.3662	-0.8997	0.3319	0.7529
	Very often-regularly	-0.1185	0.2436	0.6267	-0.5960	0.3590	0.8883
Goodness-of-fit test of overall model (Likelihood Ratio): Chi-square = 154.878, df = 32, p-value = 0.000, Pseudo R-Square=0.167							
Test of parallel lines (Likelihood Ratio): Chi-square = 44.572, df = 32, p-value = 0.069							
Rural, gender:Female, Education level:None educated, Occupation: Others, Alcohol usage: No, Smoking: No, physical activity: daily exercising were used as reference category							

Gender is the only risk factor with high blood glucose level at significance level 90% for this population group (Table 6). Men are at a higher risk of having high fasting glucose than women.

Men are about 1.45 times more likely to have high fasting glucose than women (OR, 1.4457, 90% CI, -0.0450–0.7822;  $p>0.05$ ). Dietary factors were not associated with diabetes in our study group.

**Table 6. Ordinal logistic regression showing risk factors for high fasting glucose**

		<b>High Fasting Glucose</b>					
		$\beta$	S.E.	p	<b>95% Confidence Interval</b>		<b>Odds ratio</b>
					<b>Lower Bound</b>	<b>Upper Bound</b>	
	Age	0.0102	0.0064	0.1106	-0.0023	0.0227	-
	Waist circumference	0.0183	0.0113	0.1068	-0.0039	0.0405	-
	BMI	0.0097	0.0286	0.7350	-0.0464	0.0657	-
	Energy	0.0005	0.0007	0.5044	-0.0009	0.0018	-
	Carbohydrate	-0.0019	0.0029	0.5115	-0.0075	0.0037	-
	Fat	-0.0014	0.0062	0.8241	-0.0136	0.0108	-
	Fiber	0.0090	0.0083	0.2745	-0.0072	0.0253	-
	Sodium	-0.0001	0.0001	0.4410	-0.0002	0.0001	-
	Magnesium	-0.0025	0.0017	0.1447	-0.0058	0.0009	-
	Calcium	-0.0002	0.0002	0.1405	-0.0006	0.0001	-
Gender	Male/Female	0.3686	0.2110	0.0807	-0.0450	0.7822	1.4457
	Urban/Rural	0.2459	0.1945	0.2062	-0.1354	0.6272	1.2788
Education level	Primary	0.1964	0.3236	0.5438	-0.4377	0.8306	1.2171
	JHS	-0.1676	0.3106	0.5894	-0.7764	0.4412	0.8457
	SHS/Voc/Tech	-0.2417	0.3345	0.4700	-0.8973	0.4139	0.7853
	Middle school	-0.1098	0.3074	0.7208	-0.7123	0.4926	0.8960
	Tertiary	-0.3081	0.3927	0.4327	-1.0778	0.4616	0.7348
	Other	1.0834	0.6690	0.1054	-0.2278	2.3945	2.9546
Occupation	Trading	0.2953	0.5850	0.6136	-0.8512	1.4418	1.3436
	Farming	-0.1244	0.5905	0.8331	-1.2818	1.0330	0.8830
	Artisan	0.4008	0.5781	0.4881	-0.7322	1.5338	1.4930
	Formal	0.4069	0.6250	0.5150	-0.8180	1.6319	1.5022
	Schooling	-0.2104	0.7434	0.7771	-1.6674	1.2465	0.8103
	Retired	0.8520	0.6779	0.2088	-0.4767	2.1807	2.3443
	Unemployed	0.2053	0.6334	0.7459	-1.0362	1.4468	1.2279
	Alcohol	0.1682	0.1669	0.3134	-0.1588	0.4953	1.1832
	Smoking	-0.0943	0.3429	0.7833	-0.7664	0.5778	0.9100
Physical Activity	Never/none	-0.1386	0.2917	0.6347	-0.7104	0.4332	0.8706
	Once a while	-0.0022	0.2796	0.9938	-0.5501	0.5457	0.9978

	Quite often	-0.3059	0.3706	0.4091	-1.0323	0.4205	0.7365
	Very often-regularly	-0.2603	0.2878	0.3658	-0.8245	0.3038	0.7708
Goodness-of-fit test of overall model (Likelihood Ratio): Chi-square = 59.830, df = 31, p-value = 0.001, Pseudo R-Square=0.086							
Test of parallel lines (Likelihood Ratio): Chi-square = 11.658, df = 31, p-value = 0.999							
S.E = Standard Error, CI = Confidence Interval C.I. for EXP(B)							
Rural, gender: female, Education level: none, Occupation: others, Alcohol usage:no, Smoking: no, physical activity: daily exercising were used as reference category							

### *Self-awareness of hypertension and diabetes status*

In this study, 36.8% of participants were hypertensive (Table 4), but when asked whether they were diagnosed in a healthcare facility, only 49.5% of those we detected to be hypertensive (from our measurements) were aware of being hypertensive (data not shown). As a consequence, the unaware participants were not under any treatment. Furthermore, 10% of participants had concomitant hypertension and diabetes, but only 29.0% were diagnosed in a healthcare facility.

Figure 2a presents diagnosis by a health professional vis-à-vis the actual measures of hypertension according to gender and place of residence. In total, 173 participants were aware of being hypertensive as diagnosed by a health professional, (45 males and 128 females). However, from our assessment, 126 males and 187 females were hypertensive, and an additional 64 males and 86 females had high-normal hypertension without knowing. Meanwhile, 24 participants (5 males and 19 females) reported being hypertensive but were instead found to have high-normal hypertension. This may reflect a positive impact of their medication in controlling their BP, thus reducing it to high-normal levels (pre-hypertensive) (Figure 2b).

INSERT FIGURE 2

**Figure 2. Distribution of self-awareness on hypertension status classified by gender and place of residence**

Exploring awareness on diabetes, Figure 3a presents diagnosis of diabetes by a health professional and the actual measures of diabetes according to gender and place of residence. Twelve participants reported having been diagnosed as diabetic by a health professional (six males and six females). However, from our results, 27 males and 34 females were diagnosed with diabetes, while another 48 participants (24 males and 24 females) had impaired fasting blood glucose (IFG). Furthermore, five participants (two males and three females) reported having diabetes but were found to have IFG. This might be due to a positive treatment outcome (Figure 3b).

INSERT FIGURES 3

**Figure 3. Distribution of awareness on diabetes status by gender and place of residence**

Table 7 presents an analysis of questions on knowledge of NCDs and their management. Participants were asked if they could identify one cause and one action to manage and reduce HT and DM. Regarding knowledge on hypertension, there was a significant relationship identified between correct responses and place of residence. However, there is no significant relationship between gender and correct responses for hypertension. According to our results, more females were able to state correctly one action to reduce and manage hypertension ( $p < 0.005$ ). For similar questions on DM, more females were able to mention one action to prevent and manage diabetes compare to males ( $p < 0.05$ ) and rural dwellers were able to state this compared to urban dwellers ( $p < 0.05$ ).

**Table 7. Knowledge of participants on the causes, prevention and management of hypertension and diabetes**

Questions	Correct Respond	Male n (%)	Female n (%)	P-value <sup>z</sup>	Urban n (%)	Rural n (%)	P-value <sup>z</sup>
Able to identify at least one cause of HT correctly	Y	200 (56.5)	312 (62.9)	0.060	226 (55.8)	286 (64.3)	<b>0.012</b>
	N	154 (43.5)	184 (37.1)		179 (44.2)	159 (35.7)	

Able to state at least one action to prevent/manage HT	Y	160 (45.2)	273 (55.0)	<b>0.005</b>	193 (47.7)	240 (53.9)	0.067
	N	194 (54.8)	223 (45.0)		212 (52.3)	205 (46.1)	
Able to identify at least one cause of DM correctly	Y	196 (55.4)	252 (50.8)	0.189	210 (51.9)	238 (53.5)	0.634
	N	158 (44.6)	244 (49.2)		195 (48.1)	207 (46.5)	
Able to state at least one action to prevent/manage DM	Y	154 (43.5)	179 (36.1)	<b>0.025</b>	143 (35.3)	190 (42.7)	<b>0.023</b>
	N	200 (56.5)	317 (63.9)		262 (64.7)	255 (57.3)	
<b>Total</b>		<b>354(100.0)</b>	<b>496 (100.0)</b>		<b>445 (100.0)</b>	<b>405(100.0)</b>	

<sup>a</sup>Chi-square test, HT, Hypertension; DM, Diabetes Mellitus; Y, Yes; N, No.

## Discussion

This study assessed the prevalence of hypertension and diabetes in the context of wider risk factors for NCDs among adult males and females residing in urban and rural areas in Ghana. We addressed the hypothesis that the prevalence of these risk factors and associated diseases are higher than expected for areas which are considerably less urbanised, finding this to be the case. Increasing age, being male and having a BMI in the overweight and obese category were associated with being hypertensive. The findings have good internal validity and also external validity and are therefore generalisable.

Other Ghanaian studies have reported advancing age and overweight/obesity as risk factors for increasing BP and thus HT (14, 15). In urban Accra and a peri-urban area, males were found to have an increased risk for HT. Likewise we found that being male presents a higher risk (15, 29) . We found in our study the only risk factor for diabetes was being male. There was no association between dietary intake (such as energy, fat, micronutrients etc.) and hypertension and diabetes. There are other factors known to be related to NCDs which we analysed for our study participants, however none of these were risk factors for the participants.

We visited participants in their home's early morning when household members of working age (males and younger females) may already have left for work. In rural settings, this

assumption corroborates the observation that men usually leave for the farm much earlier than females and allow the females to follow much later as they were responsible for preparing the children for school and also preparing food to take along to the farm. This could, therefore, be a plausible reason why more females were recruited compared to males.

We also found urban-rural differences. Our mean diastolic ( $82.35 \pm 14.82$  mmHg) and systolic BP ( $132.12 \pm 23.9$  mmHg) were similar as reported by Obirikorang et al. (80.0 vs 133 mmHg) (12). They also found that among urban dwellers, diastolic BP was significantly higher and they were more likely to have cardiometabolic risk factors such as higher BMI and WC (12). We found a significant relationship between the systolic BP of rural inhabitants. Meanwhile, research on cardiovascular risk factors in relation to rural-urban transition shows a significantly higher prevalence of hypertension among urban inhabitants (11).

In our findings, BMI was higher in rural areas, yet like other studies on obesity and cardiometabolic risk factors, WC and WHR were higher in urban areas (12). Compared to lateral obesity, (where the fat is distributed around the body), central obesity imposes a higher risk for cardiometabolic diseases and could reflect unhealthy dietary habits and sedentary lifestyle in urban areas.

Overweight and obesity are risk factors for NCDs, including hypertension, diabetes and cardiovascular diseases (12). Increasing BMI is a known risk factor for higher BP, thereby predisposing overweight and obese individuals to hypertension compared to normal-weight individuals (30). We observed a significant relationship between higher BMI and the category of hypertension but this was not surprising as similar results have been reported (14, 15, 29, 31). While we observed a 1.7-fold likelihood of developing high BP with increasing BMI, 2.6-fold (95% CI =2.3–2.9) increased likelihood for hypertension has been reported among

overweight and obese persons (32). Furthermore, a significant relationship was observed between WC and hypertension; a finding confirmed by Jayawardana et al. (31).

Research shows a 35% prevalence of hypertension in Ghana (33), which is similar to our findings of 36.8%. However, our results support a direct relationship between increasing age and hypertension despite the majority of our sample being below 65 years of age. Other studies in Ghana have similarly reported aging as a risk factor for hypertension with ages 22-59 (34), 18-45 (15) and  $\geq 40$  years (35) being the most associated with hypertension (31). Hypertension leads to changes in the vasculature which includes endothelial dysfunction, vessel walls thickening, reduced flexibility and arterial stiffening as a consequence of aging (36) thereby affecting the development and progression of micro and macro complications of diabetes unless interventions are instituted (37). Concomitant hypertension and diabetes increase the risk of metabolic syndrome (37). Like our findings, a recent study in Ghana found a positive association between fasting blood glucose and BP in urban areas but with only diastolic BP in rural areas (12). Elsewhere in Ghana, the prevalence of hypertension was 32.4%, but only 46.2% of the hypertensive individuals were aware of their status (33).

Few studies have reported on the combined awareness of hypertension and diabetic status in African communities (35, 38). In our study, 10.0%, of hypertensives had concurrent diabetes, whereas 20.3% of hypertensives were pre-diabetics. We noticed large numbers of participants being unaware of either their hypertension or diabetic status, which could increase the incidence of stroke (39). This finding indicates a public health problem which needs to be addressed through enhanced health screening in communities. In relation to awareness, more females correctly identified factors which trigger hypertension as well as their management. Despite this, females were more hypertensive. This could be linked to their availability and willingness to participate in research, complemented by their positive health-seeking behaviour. We identified more hypertension among rural inhabitants and a significant

relationship with their knowledge. Similarly, there was a significant relationship between urban dwelling and knowledge on the management of diabetes. However, we identified less awareness of diabetes in urban areas.

Excessive intake of dietary salt is related with increased risk of hypertension (40). The Dietary Approach to Stop Hypertension (DASH) study shows that a reduction in sodium intake decreases BP (34). In line with this, we expected a significant association between consumption of dietary sodium and hypertension, but that was not observed. Williams et al. reported a significant association between salt and alcohol consumption and diastolic BP, especially in males (14), similar as reported by Mohammed et al. (15). We, however, did not obtain any association between alcohol consumption and BP.

Magnesium also regulates BP (41), with an inverse relationship between dietary magnesium intake and hypertension risk. In a recent systematic review, 100 mg/day increase in dietary magnesium intake decreased hypertension risk by 5% (42). Public Health England 's recommendations for magnesium intake for males and females are 300 mg/day and 270 mg/day, respectively (22). We noticed low consumption of magnesium (males:66.03±74.92 vs females:54.58±62.89 mg/day). Increasing dietary intakes of magnesium is essential for health and also to reduce hypertension in this community as intakes are below the reference nutrient intake. Contrary to expectations, no significant association was observed between dietary magnesium intake and hypertension. Establishing optimal intakes of magnesium for this population is worthy of further research. Moreover, this an observational study. To observe an effect, an analytical design is required. Effect of dietary fibre on decreasing BP has been reported (43) but we did not find any significant difference between participants' consumption of dietary fibre and hypertension.

Hypertension is a major risk factor contributing to increased morbidity and mortality, particularly for NCDs. Hypertension and its related co-morbidities impact on health systems in many sub-Saharan African countries (44). Currently, 70% of individuals with diagnosed hypertension are not accessing treatment, while only 13% have controlled BP. In terms of diabetes, the reported prevalence is 9% (5). Nonetheless, as our study shows, this may be an underestimation especially as accessibility and affordability of health service for diabetes detection, management, and control are limited, especially in the rural areas (45).

There was an apparent shift in the health status of the rural inhabitants, particularly women, who were less active, thus predisposing them to greater risk for NCDs. The assumption that there is lower prevalence of these risk factors in rural areas is now challenged, as we observed an epidemiological shift consistent with increasing urbanisation. However, we found no evidence of nutrition transition affecting the diets of rural inhabitants. Rural-to-urban transitions are often associated with marked changes in behaviour and lifestyle, such as diminished physical activity. However, our data suggests a level of inactivity already exists [11].

In Ghana, rising levels of NCDs are mainly attributed to inadequate health care and interventions (46). However, the rapid changes, particularly in rural areas, are changing the landscape and exacerbating the NCD problem (47) (48). Our findings support the need for community-based screening using inexpensive rapid test kits and educational interventions in rural areas (using a variety of media and social media) to address risk factors for hypertension and diabetes in populations who are clearly predisposed but unaware of their risk status. Improvements in physical activity and dietary intakes that provide key micronutrients are necessary (49). Health professionals working in these areas need to be mindful of health literacy and ensure that behaviour change communication takes cognisance of the low education of rural communities (50).

## **Limitations**

Despite the generalisability of the study findings, some limitations are worth noting. Being a cross-sectional study, we are limited in assessing causation and studying prevalence and incidence over time. Low levels of physical activity, including its under- and over-reporting, may influence the findings. Although participants asserted undertaking physical activity, the nature and extent is unknown. Also, data on the intensity of physical activity was not collected. Dietary data is self-reported, which is often subject to reporting and memory biases. Using a one-day diet recall method rather than a three-day diet diary may have resulted in the lack of association between dietary factors and HT and diabetes. Use of digital dietary methodologies may be a useful future approach. The questions used for alcohol consumption, smoking and physical activity were crude measures of exposure thereby lacking internal validity to reflect real exposures. To provide information on the nature and extent, more valid measures would have been appropriate. Our methodology may also have affected the prevalence of diabetes because not all participants were eligible for BP testing. Furthermore, type 1 and type 2 diabetes could not be differentiated despite the varying prevalence and risk factors among older adults. Cholesterol levels were not measured due to financial constraints. It would have enhanced the findings in relation to cardio-metabolic risk. Lastly, we did not investigate some of the known risk factors of diabetes, such as genetic predisposition. Despite these limitations, there are a number of strengths of this study. Based on published data, the RIING software is a validated dietary database for Ghanaian foods and has enhanced the value of the dietary data. With a powerful sample size of 850 participants, our findings provide valuable data for health policymakers to initiate public health interventions. Our study is the first of its nature in the Volta region and Hohoe, Ghana.

## **Conclusions**

We conclude there is an epidemiological shift consistent with increasing urbanisation in these rural areas which is concerning. Our prevalence study, showed males, older persons and those who are overweight/obese have greater odds/risk for hypertension, consistent with other literature. Pre-diabetes, and pre-hypertension, in addition to undiagnosed diabetes and hypertension were common with high proportions of these individuals unaware of their health status. Poor diet underpins the rates.

We conclude that efforts to mitigate the rising levels of hypertension and diabetes require concerted action, intensified testing, awareness creation on NCDs, lifestyle interventions to improve dietary intakes, physical activity and health-seeking behaviours that involve rural inhabitants as well.

## **Declarations**

## **Consent for publication**

Not applicable

## **Ethical approval**

This research was conducted in accordance with the Declaration of Helsinki. The study was approved by the University of Chester's Faculty of Health and Social Care Ethics Sub-Committee (RESC0514-518) and the Ghana Health Service Ethical Review Committee (GHS-ERC: 060714).

## **Data Availability**

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

**Competing interests:** None exist

**Funding:** University of Chester (HSC 14/220).

**Authors' contributions:** BE, FA and FBZ conceived the study, drafted the protocol and designed the instruments for data collection. BE acquired funding. FA, FZ and BE trained, collected and supervised the data collection process. OFK cleaned and managed the data obtained and DD undertook the analysis supervised by BE. BE, DD and BS wrote the first draft of the manuscript. BE and DD undertook manuscript redrafting. All authors critically reviewed the manuscript and also interpreted the findings for intellectual soundness.

**Acknowledgements:** We appreciate the cohort of students who pursued Bachelor in Public Health Nutrition programme at the School of Public Health, University of Health and Allied Sciences Hohoe campus. They supported immensely with data collection. Special thanks to Samuel Darrah and Carl Agbittor who were pivotal in the data collection process and Gemma Rogers (University of Chester) for her assistance with the submission of this paper. We also thank all the individuals who agreed to participate in the study.

## **References**

1. World Health Organisation (WHO). Global status report on noncommunicable diseases 2010. 2011.
2. World Health Organisation (WHO). Non Communicable Diseases Country Profiles 2018. Switzerland; 2018.
3. International Diabetes Federation (IDF). IDF Diabetes Atlas Tenth Edition. 2021.
4. World Health Organisation (WHO). Noncommunicable Diseases (NCD) Country Profiles, Ghana, 2018.; 2018.

5. Ghana Health Service. Annual Report 2016. 2017.
6. Cappuccio FP, Miller MA. Cardiovascular disease and hypertension in sub-Saharan Africa: burden, risk and interventions. *Intern Emerg Med*. 2016;11(3):299-305.
7. Bosu WK. Epidemic of hypertension in Ghana: a systematic review. *BMC Public Health*. 2010;10:418.
8. Ama de-Graft Aikins SA-M, Charles Agyemang. Diabetes in Ghana: a review of research on prevalence, experiences and healthcare. In: Ama de-Graft Aikins SA-M, Charles Agyemang, editor. *Chronic Non-communicable Diseases in Ghana: Multidisciplinary Perspectives*: Sub Sahara Publishers; 2018. p. 41-55.
9. World Health Organization (WHO). Ghana Diabetes country profile 2016.
10. Bosu WK, Aheto JMK, Zucchelli E, Reilly S. Prevalence, awareness, and associated risk factors of hypertension in older adults in Africa: a systematic review and meta-analysis protocol. *Syst Rev*. 2017;6(1):192.
11. Kodaman N, Aldrich MC, Sobota R, Asselbergs FW, Poku KA, Brown NJ, et al. Cardiovascular Disease Risk Factors in Ghana during the Rural-to-Urban Transition: A Cross-Sectional Study. *PLoS One*. 2016;11(10):e0162753.
12. Obirikorang C, Osakunor DN, Anto EO, Amponsah SO, Adarkwa OK. Obesity and Cardio-Metabolic Risk Factors in an Urban and Rural Population in the Ashanti Region-Ghana: A Comparative Cross-Sectional Study. *PLoS One*. 2015;10(6):e0129494.
13. Awuah RB, Anarfi JK, Agyemang C, Ogedegbe G, Aikins A. Prevalence, awareness, treatment and control of hypertension in urban poor communities in Accra, Ghana. *J Hypertens*. 2014;32(6):1203-10.
14. Williams EA, Keenan KE, Ansong D, Simpson LM, Boakye I, Boaheng JM, et al. The burden and correlates of hypertension in rural Ghana: a cross-sectional study. *Diabetes Metab Syndr*. 2013;7(3):123-8.



15. Mohammed H, Ghosh S, Vuvor F, Mensah-Armah S, Steiner-Asiedu M. Dietary intake and the dynamics of stress, hypertension and obesity in a peri-urban community in Accra. *Ghana Med J.* 2016;50(1):16-21.
16. Angkurawaranon C, Wattanatchariya N, Doyle P, Nitsch D. Urbanization and non-communicable disease mortality in Thailand: an ecological correlation study. *Trop Med Int Health.* 2013;18(2):130-40.
17. Rani R. MV, Gupta R. K., Sharma H. K. Hypertension and Its Risk Factors – A Cross Sectional Study in an Urban Population of a North Indian District. *Public Health Research.* 2015;5(3):67-72.
18. World Health Organisation (WHO). *Health in 2015: from MDGs, Millennium Development Goals to SDGs, Sustainable Development Goals 2015.*
19. Ghana Statistical Service. *Facts and Figures.* 2015.
20. Chao LW, Szrek H, Peltzer K, Ramlagan S, Fleming P, Leite R, et al. A Comparison of EPI Sampling, Probability Sampling, and Compact Segment Sampling Methods for Micro and Small Enterprises. *J Dev Econ.* 2012;98(1):94-107.
21. Center for Disease Control and Prevention [CDC] NHANES. *Anthropometry Procedures Manual.*; 2007.
22. Public Health England. *Government Dietary Recommendations. Government recommendations for energy and nutrients for males and females aged 1 – 18 years and 19+ years.* 2016.
23. Johansson L, Solvoll K, Bjorneboe GE, Drevon CA. Under- and overreporting of energy intake related to weight status and lifestyle in a nationwide sample. *Am J Clin Nutr.* 1998;68(2):266-74.
24. World Health Organisation (WHO). *Waist Circumference and Waist-Hip Ratio Report of a WHO Expert Consultation.* Geneva; 2008.

25. World Health Organisation (WHO). Body mass index - BMI [Available from: <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>].
26. National Institute for Health and Care Excellence (NICE). Hypertension in adults: diagnosis and management 2016 [Available from: <https://www.nice.org.uk/guidance/cg127/chapter/1-Guidance>].
27. World Health Organisation (WHO) IDF. Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycemia. 2006.
28. Kobalava ZD, Kotovskaya YV. [Isolated Systolic Hypertension in Different Ages]. *Kardiologiya*. 2015;55(9):84-90.
29. Afrifa-Anane E, Agyemang C, Codjoe SN, Ogedegbe G, de-Graft Aikins A. The association of physical activity, body mass index and the blood pressure levels among urban poor youth in Accra, Ghana. *BMC Public Health*. 2015;15:269.
30. Tsukinoki R, Murakami Y, Huxley R, Ohkubo T, Fang X, Suh I, et al. Does body mass index impact on the relationship between systolic blood pressure and cardiovascular disease?: meta-analysis of 419 488 individuals from the Asia Pacific Cohort Studies Collaboration. *Stroke*. 2012;43(6):1478-83.
31. Jayawardana N, Jayalath W, Madhujith WMT, Ralapanawa U, Jayasekera RS, Alagiyawanna S, et al. Aging and obesity are associated with undiagnosed hypertension in a cohort of males in the Central Province of Sri Lanka: a cross-sectional descriptive study. *BMC Cardiovasc Disord*. 2017;17(1):165.
32. Nuertey BD, Alhassan AI, Nuertey AD, Mensah IA, Adongo V, Kabutey C, et al. Prevalence of obesity and overweight and its associated factors among registered pensioners in Ghana; a cross sectional study. *BMC Obes*. 2017;4:26.

33. Lamptey P, Laar A, Adler AJ, Dirks R, Caldwell A, Prieto-Merino D, et al. Evaluation of a community-based hypertension improvement program (ComHIP) in Ghana: data from a baseline survey. *BMC Public Health*. 2017;17(1):368.
34. Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, et al. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med*. 2001;344(1):3-10.
35. Asamoah-Boaheng M, Sarfo-Kantanka O, Tuffour AB, Eghan B, Mbanya JC. Prevalence and risk factors for diabetes mellitus among adults in Ghana: a systematic review and meta-analysis. *Int Health*. 2019;11(2):83-92.
36. Harvey A, Montezano AC, Lopes RA, Rios F, Touyz RM. Vascular Fibrosis in Aging and Hypertension: Molecular Mechanisms and Clinical Implications. *Can J Cardiol*. 2016;32(5):659-68.
37. Mogre V, Apala P, Nsoh JA, Wanaba P. Adiposity, hypertension and weight management behaviours in Ghanaian type 2 diabetes mellitus patients aged 20-70 years. *Diabetes Metab Syndr*. 2016;10(1 Suppl 1):S79-85.
38. Gatimu SM, Milimo BW, Sebastian MS. Prevalence and determinants of diabetes among older adults in Ghana. *BMC Public Health*. 2016;16(1):1174.
39. Sarfo FS, Mobula LM, Plange-Rhule J, Ansong D, Ofori-Adjei D. Incident stroke among Ghanaians with hypertension and diabetes: A multicenter, prospective cohort study. *J Neurol Sci*. 2018;395:17-24.
40. Rust P, Ekmekcioglu C. Impact of Salt Intake on the Pathogenesis and Treatment of Hypertension. *Adv Exp Med Biol*. 2017;956:61-84.



41. Lennon SL, DellaValle DM, Rodder SG, Prest M, Sinley RC, Hoy MK, et al. 2015 Evidence Analysis Library Evidence-Based Nutrition Practice Guideline for the Management of Hypertension in Adults. *J Acad Nutr Diet.* 2017;117(9):1445-58 e17.
42. Han H, Fang X, Wei X, Liu Y, Jin Z, Chen Q, et al. Dose-response relationship between dietary magnesium intake, serum magnesium concentration and risk of hypertension: a systematic review and meta-analysis of prospective cohort studies. *Nutr J.* 2017;16(1):26.
43. Aleixandre A, Miguel M. Dietary fiber and blood pressure control. *Food Funct.* 2016;7(4):1864-71.
44. Tenkorang EY, Kuuire V, Luginaah I, Banchani E. Examining risk factors for hypertension in Ghana: evidence from the Study on Global Ageing and Adult Health. *Glob Health Promot.* 2017;24(1):14-26.
45. Petti CA, Polage CR, Quinn TC, Ronald AR, Sande MA. Laboratory medicine in Africa: a barrier to effective health care. *Clin Infect Dis.* 2006;42(3):377-82.
46. Ofori-Asenso R, Agyeman AA, Laar A, Boateng D. Overweight and obesity epidemic in Ghana-a systematic review and meta-analysis. *BMC Public Health.* 2016;16.
47. Boateng GO, Adams EA, Odei Boateng M, Luginaah IN, Taabazuing MM. Obesity and the burden of health risks among the elderly in Ghana: A population study. *PLoS One.* 2017;12(11):e0186947.
48. Lloyd-Sherlock P, Beard J, Minicuci N, Ebrahim S, Chatterji S. Hypertension among older adults in low- and middle-income countries: prevalence, awareness and control. *Int J Epidemiol.* 2014;43(1):116-28.
49. Chomistek AK, Chiuve SE, Eliassen AH, Mukamal KJ, Willett WC, Rimm EB. Healthy lifestyle in the primordial prevention of cardiovascular disease among young women. *J Am Coll Cardiol.* 2015;65(1):43-51.

50. Ellahi B. Practical evaluation of Public Health Interventions in diverse communities  
In: Judith L. Buttriss AAW, John M. Kearney, Dr. Susan A. Lanham-New, editor. Public  
Health Nutrition, 2nd Edition: Nutrition Society, Wiley-Blackwell; 2017. p. 406-12.





**Figure 1. Mean diastolic and systolic blood pressure values classified according to educational level and place of residence**

BP: Blood pressure



**Figure 2. Distribution of self-awareness on hypertension status classified by gender and place of residence**

HPT: Hypertension



**Figure 3. Distribution of awareness on diabetes status by gender and place of residence**

IFG: Impaired fasting glucose; DM: Diabetes